		SEMESTER II
18BTU201	GENETICS	<b>4H - 4C</b>
Total hours/week: L:4 T:0 P:0	Marks: Internal: 40 External: 60	Total: 100

**Course Objectives:** The objective of this course is to present basic concepts of heredity and various in living organism and be able to articulate connection between the multiple level of organization, molecular mechanism the expression of DNA at individual level and the transmission of DNA across generation at the individual and population level.

**Course Outcomes:** upon successful completion of this course student should be able to demonstrate the following competencies; an understanding the central theories and methodologies traditional, molecular and population genetics.

### UNIT- I

**Introduction:** Historical developments in the field of genetics. Organisms suitable for genetic experimentation and their genetic significance. Prokaryotic genetics. Cell Cycle: Mitosis and Meiosis: Control points in cell-cycle progression in yeast. Role of meiosis in life cycles of organisms.

### UNIT-II

**Mendelian genetics :** Mendel's experimental design, monohybrid, di-hybrid and tri hybrid crosses, Law of segregation & Principle of independent assortment. Verification of segregates by test and back crosses, Chromosomal theory of inheritance, Allelic interactions: Concept of dominance, recessiveness, incomplete dominance, co-dominance, semi-dominance, pleiotropy, multiple allele, pseudo-allele, essential and lethal genes, penetrance and expressivity.

#### UNIT-III

**Chromosome and genomic organization:** Eukaryotic nuclear genome nucleotide sequence composition –unique & repetitive DNA, satellite DNA. Centromere and telomere DNA sequences, repetitive transposed sequences- SINEs & LINEs. Genetic organization of prokaryotic and viral genome. Structure and characteristics of bacterial and eukaryotic chromosome, chromosome morphology, concept of euchromatin and heterochromatin. Concept of cistron, exons, introns, genetic code, gene function.

#### UNIT-IV

**Chromosome and gene mutations:** Definition and types of mutations, causes of mutations, Ames test for mutagenic agents, screening procedures for isolation of mutants and uses of mutants, variations in chromosomes structure - deletion, duplication, inversion and translocation (reciprocal and Robertsonian). Sex determination, sex linkage, sex linked diseases: Mechanisms of sex determination, Fragile-X-syndrome and chromosome, sex influenced dominance, sex limited gene expression, sex linked inheritance.

#### UNIT-V

**Genetic linkage, crossing over and chromosome mapping:** Linkage and Recombination of genes in a chromosome crossing over. Extra chromosomal inheritance: Rules of extra nuclear inheritance, maternal effects, maternal inheritance, cytoplasmic inheritance, organelle heredity, genomic imprinting. Evolution and population genetics: In breeding and out breeding, Hardy Weinberg law

(prediction, derivation), allelic and genotype frequencies, changes in allelic frequencies, systems of mating, evolutionary genetics, natural selection.

#### References

- 1. Gardner, E.J., Simmons, M.J., & Snustad, D.P. (2006). *Principles of Genetics* (8th ed.). John Wiley & Sons.
- 2. Snustad, D.P., & Simmons, M.J. (2009). *Principles of Genetics* (5th ed.). USA: John Wiley and Sons Inc.
- 3. Russell, P. J. (2009). Genetics- A Molecular Approach (3rd ed.). Benjamin Cummings.
- 4. Griffiths, A.J.F., Wessler, S.R., Lewontin, R.C., & Carroll, S.B. (2007). *Introduction to Genetic Analysis* (9th ed.). W. H. Freeman & Co.



Enable | Enlighten | Enrich (Deemed to be University) (Under Section 3 of UGC Act 1956)

### **KARPAGAM ACADEMY OF HIGHER EDUCATION**

(Deemed to be University Established Under Section 3 of UGC Act 1956) Coimbatore – 641 021.

### LECTURE PLAN DEPARTMENT OF BIOTECHNOLOGY

STAFF NAME: Dr Anilkumar & Dr.Nishu Sekar SUBJECT NAME: Genetics SEMESTER: II

SUB.CODE:18BTU201 CLASS: I B.Sc (BT)

S.No	Lecture Duration Period	Topics to be Covered	Support Material/Page Nos	
		UNIT-I		
1	1	Historical development in the field of genetics	T1:15-16	
2	1	Organisms suitable for genetic experimentation	W1: R1 28-3	
3	1	Prokaryotic genetics		
4	1	Cell cycle mitosis and meiosis	W1	
5	1	Control points in cell cycle progression in yeast	W1	
6	1	Role of meiosis in life cycle of organism	W1	
7	1	Revision of unit-I		
8	1	Class test of unit-I		
	Total No of Hou	otal No of Hours Planned For Unit 1=08		
		UNIT-II		
1	1	Mendels experiment design	R1:73-81	
2	1	Mono hybrid, di-hybrid and tri- hybrid crosses	T121-24	
3	1	Law of segregation & principle of independent assortment	T121-24	
4	1	Verification of segregates by test and back crosses	R1:106-119	
5	1	Chromosomal theory of inheritance	W1	

8	-2021
ch	í .

6	1	allelic interaction concepts of	R1:106-119
		dominance, recessiveness	
7	1	Concepts of incomplete	R1:106-119
		dominance, co-dominance	
8	1	Concepts of semi dominance,	R1120-123
		pleiotropy	
	Total No of H	ours Planned For Unit II=08	
		UNIT-III	
1	1	Eukaryotic nuclaear genome	R1:93-95
2	1	Nucleotide sequence	R195:T1:36-54
		composition-unique & repetitive	
		DNA	
3	1	Satellite DNA	T1: 536-542
	1		<b>T</b> 1 <b>C</b> 2 <b>C C 1</b> 2
4		Centromere and telomere DNA	11: 536-542
5	1	Respective transposed sequences	
6	1	Genetic organization of	
		prokaryotic and viral genome	
7	1	Structure and characteristics of	T1: 536-542
		bacterial and eukaryotic	
		chromosomes	
8	1	Chromosome morphology	T1: 536-542
	Total No of Hours Planned For Unit III=08		
		UNIT-IV	
1	1	Definition and types of mutation	T1 167-176
2	1	Causes of mutations	R1:214-215
3	1	Ames test for mutagenic agents	T1:167-176
4	1	Screening procedure for	T1:167-176
		isolations of mutants and use of	
		mutants	
5	1	Variations in chromosomes	R1:212-214; W1
		structure deletion, duplication	
6	1	Inversion and translocation	R1:212-214
7	1	Sex determination	R1:212; W1
8	1	Sex linkage, sex linked diseases	R1:212W1
	Total No. of H	ours Planned For Unit IV-08	

Prepared by Dr Anilkumar and Dr. Nishu Sekar, Department of Biotechnology, KAHE

)18	-2	0	2	1
atch	l			

	1		
		UNIT-V	
1	1	Linkage and recombination of	T1:317-318
		genes in a chromosome crossing	W1
2	1	Rules of extra nuclear inheritance	T1:318-319
3	1	Material inheritence	T1:427-428
4	1	Cytoplasmic inheritance,	T1:428-429
		organism heredity, genomic	
		imprinting	
5	1	Evolution and population	T1:429-432
		genetics, inbreeding and out	
		breeding	
6	1	Hardy Weinberg law	T1:440-441
7	1	Alleic and genotype frequency	T1:459,460
8	1	Changes in allelic frequency	T1:463-466,R2
	Total No of Hours Planned for unit V=40		
Total	40		
Planned			
Hours			

### **TEXT BOOK**

 Gardener, G.,2001, principles of genetics john wiley and sons Inc, New york (Page Nos. : 24-54, 56-77, 80-92, 103-118, 204-215, 236-256, 427-443, 458-463, 490-495, 502-505)

### REFERENCES

- 1. P.S.Verma and V.K aggarwal., 2000 Genetics, S.chand and company ltd New Delhi.
- 2. Monoroe w.stricberger, 1985, genetics 3<sup>rd</sup> edition, macmillan

### **WEBSITES**

1. WWW.nptel.com

CLASS: I B Sc BT **COURSE CODE: 18BTU201** 

**COURSE NAME: GENETICS** UNIT: I (Introduction)

BATCH-2018-2021

### UNIT-I

### **SYLLABUS**

GENETICS: Introduction The Basics: Historical development in the field of genetics, Organisms suitable for genetic experimentation, Prokaryotic genetics, Cell cycle mitosis and meiosis, Control points in cell cycle progression in yeast, Role of meiosis in life cycle of organism

### Unit – I

#### **1. INTRODUCTION**

Genetics is the study of genes, and tries to explain what they are and how they work. Genes are how living organisms inherit features from their ancestors; for example, children usually look like their parents because they have inherited their parents' genes. Genetics tries to identify which features are inherited, and explain how these features are passed from generation to generation.

#### **SCOPE OF GENETICS**

"The experiments which will here be discussed took their origin from artificial fertilizations which have been carried out on ornamental plants in order to obtain new color variants." This opening sentence of Gregor Mendel's paper points to one of the humble origins of genetics: horticulture. Significantly, as late as 1901, it was the Royal Horticultural Society of London that sponsored the English translation of the then newly rediscovered classic. But the practical desires of the lovers of flowers were not the only incentives to the birth of genetics. All through the eighteenth and nineteenth centuries the species problem occupied the minds of biologists. On the one hand, Linnaeus had defined species in a way which lacked operational usefulness: There are as many species as there were diverse forms created at the beginning by the Infinite Being. On the other hand, Linnaeus and other plant hybridizers attempted to probe into the nature of species by experimental means. Were hybridizations between different species possible? Could they be induced artificially? Did they ever occur in nature? What kind of organism was a species hybrid?

Was it sterile like the mule? If fertile, what kind of offspring did it produce? In 1859, Darwin's Origin of Species proved to be the successful break through the confines of a static conception of species within which biology had accumulated so much of its unordered bulk of empirical facts. Then plant hybridization became a potential tool for progress in a new range. Mendel himself, who began his studies before the appearance of Darwin's book but wrote his report six years after that event, by then had recognized clearly the role of genetics in this respect. The detailed analysis of successive generations of hybrids "seems to be the only right way," he wrote, "by which we can finally reach the solution of a question the importance of which should not be underestimated in connection with the history of the evolution of organic forms." Genetics and evolution were indeed to become intimately and fruitfully associated with one another; but the scope of genetics was destined to embrace many other regions. From variations in color and form, and from the species problem and hybridization, the emphasis soon shifted to the permanent units which underlie the externally visible phenomena. Mendel's essential discovery had been the recognition that the hereditary contributions of two parents were not each undefined wholes which in the offspring mingled into a new whole. On the contrary, he showed that the hereditary makeup consists of many separable elements which recombine in many ways in the sequences of generations and always reappear in unchanged nature. Bateson called these units factors, "makers." Johannsen, in 1909, coined the term "gene." For him, "the word gene was ... completely free from any hypothesis." It was meant "solely to express the fact that characters of the organism are conditioned by special, at least partially separable and thus in a sense independent 'states,' 'factors,' 'units' or 'elements' in the constitution of the germcells."

The scope of genetics as a central biological discipline rests on the fact that genes are those parts of biological systems which give continuity to them. The primitive experiment of cutting an amoeba in two and finding that only the nucleated part survives and multiplies has been refined until we know that it is the genic material on which survival and multiplication of the organism depends. Most of the cellular material may be removed without permanent harm. Regeneration soon takes care of replacement. A gene, however, once removed from a cell, is not regenerated. Within a cell, a gene always comes from a gene. Again, this attribute of the gene is part of the organization of the whole. The material structure of the gene seems to carry within it the potentiality of its replication, but actual genic multiplication depends on the presence of a whole

assembly of structures and processes. The discovery of the cyclical nature of many biochemical sequences in cellular metabolism has been one of the significant events of our times. Genes are agents in these cycles not only because they determine the existence of enzymes, without which the reactions would not proceed sufficiently, but because some of these reactions themselves must lead to the making of more genes. We admire a jongleur who tosses and keeps rotating several rings moving in space. Figuratively speaking, a cell consists of thousands of such rings functionally bound together not by an outside mover but by themselves. The rings are not all of equal rank. Destroy some, and the rest will re-create them. Destroy a single agent in others, and the whole system will disintegrate. These single agents are the genes. Through them, cellular physiology is brought within the scope of genetics. Rightly, more than thirty years ago, Muller called the gene "the basis of life."

Evolution early was close to the mind of the geneticist. Its study is now unthinkable without the framework of genetics. The differences between species are largely if not exclusively due to differences in their genes. Evolution is based on mutability, the origin of new varieties of genes, on the distribution and recombination's of these varieties within the members of populations, and on the effects which these diverse genotypes have on the fitness of individuals and of populations as whole entities. The New Systematic and the Modern Synthesis in evolution, to borrow phrases from Julian Huxley, are witness to the part which genetics has taken in the transformation of these subjects. Evolution is a historical field. Just as the history of the earth, or that of astronomical systems, is based on the properties of elementary physical particles, so the history of life on our planet is based on the properties of its genetic material. These sciences of their own spheres of endeavor. To say-as it has been said that a lion or a palm is only some genes' way of making more genes expresses deep insights which link the lion and the palm to genetics-although little work on the genetics of lions or palms has been carried out. But the lions and palms remain valid topics for biological study as lions and palms. How evolution brought forth these specific species is not explained simply by referring to their genes. The analysis of animal behavior has rightly been re-recognized as another central discipline within biology. The inheritance of behavior patterns and the inheritance of plasticity in the range of possible behavioral phenomena are being rediscovered in general and analyzed in detail. Genes determine the degree of exploratory behavior in rats, intelligence or dullness in running through a maze,

fear reactions in birds to an object resembling in outline a specific predator, or the details of a complex song. This is amazing and intelligible at the same time. If chemicals can change intimately one's reactions, if under the influence of drugs spiders change the pattern according to which they weave their webs, why should not genes as biochemical cellular agents do the same? If genes can help to organize the differential development of a complex nervous system, why should they not be able to incorporate in it the subtleties of inborn reactions? The wonder is that there is behavior. Genetics provides one of the analytical tools for its study.

The scope of genetics, with its enlightenment of all branches of biology, makes it an obvious partner in the pursuit of applied biology. From horticulture to plant and animal breeding in general, the concepts and methods of genetics-be it individual or strain genetics in terms of hybridization and segregation, or population genetics in terms of equilibria and selection-form indispensable parts of procedure. Problems of pest control involve the existence of innately sensitive strains and the origin of resistant ones. Host-parasite relations depend on a balance of genetically determined interactions between the two components of the combating teams.

Human genetics is both a basic and an applied science. Knowledge of the genetic makeup of the species Homo sapiens, in its individuals and varieties, is the latest gain in the long history in which anatomy and physiology, in ever more penetrating modes, have provided insight into the biological nature of Man and anthropology into that of men. Eugenics, the planning toward rational improvement of the genetic makeup of human populations, has been regarded as a utopian dream and a dangerous one in addition. Many of our concerns have been recognized as being excessive or even unnecessary, and our eagerness to act precipitously has rightly decreased. But, while the attempt at improvement beyond the present average could well be relegated to the future, even the most determined opponents of positive eugenics have now realized with the fervor of converts that negative eugenics, the attempt to inhibit the increase of harmful genotypes, is an immediate necessity.

A good understanding of the principles of Mendelian inheritance is a prerequisite to the conceptual understanding of evolutionary theory. Indeed, though Darwin himself subscribed to

### KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B Sc BTCOURSE NAME: GENETICS

### COURSE CODE: 18BTU201 UNIT: I (Introduction)

BATCH-2018-2021

the "blending" theory of inheritance, it has since been determined that evolution by natural selection requires discrete genes.

### HISTORY OF GENETICS

Year	Scientist(s)	Discovery
1858	Charles Darwin Alfred Russel Wallace	Joint announcement of the theory of natural selection-that members of a population who are better adapted to the environment survive and pass on their traits.
1859	Charles Darwin	Published The Origin of Species.
1866	Gregor Mendel	Published the results of his investigations of the inheritance of "factors" in pea plants.
1900	Carl Correns Hugo de Vries Erich von Tschermak	Mendel's principles were independently discovered and verified, marking the beginning of modern genetics.
1902	Walter Sutton	Pointed out the interrelationships between cytology and Mendelism, closing the gap between cell morphology and heredity.
1905	Nettie Stevens Edmund Wilson	Independently described the behavior of sex chromosomes-XX determines female; XY determines male.
1908	Archibald Garrod	Proposed that some human diseases are due to "inborn errors of metabolism" that result from the lack of a specific enzyme.

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 5/5

CLASS: I B Sc BTCOURSE NAME: GENETICSCOURSE CODE: 18BTU201UNIT: I (Introduction)BATCH-2018-2021

1910	Thomas Hunt Morgan	Proposed a theory of sex-linked inheritance for the first mutation discovered in the fruit fly, Drosophila, white eye. This was followed by the gene theory, including the principle of linkage.
1927	Hermann J. Muller	Used x-rays to cause artificial gene mutations in Drosophila.
1928	Fred Griffith	Proposed that some unknown "principle" had transformed the harmless R strain of Diplococcus to the virulent S strain.
1931	Harriet B. Creighton Barbara McClintock	Demonstrated the cytological proof for crossing-over in maize.
	George Beadle	Irradiated the red bread mold, Neurospora, and proved that the
1941	Edward Tatum	gene produces its effect by regulating particular enzymes.
1944	Oswald Avery Colin MacLeod Maclyn McCarty	Reported that they had purified the transforming principle in Griffith's experiment and that it was DNA.
1945	Max Delbruck	Organized a phage course at Cold Spring Harbor Laboratory which was taught for 26 consecutive years. This course was the training ground of the first two generations of molecular biologists
late 1940s	Barbara McClintock	Developed the hypothesis of transposable elements to explain color variations in corn.
1950	Erwin Chargaff	Discovered a one-to-one ratio of adenine to thymine and guanine to cytosine in DNA samples from a variety of organisms.
1951	Rosalind Franklin	Obtained sharp X-ray diffraction photographs of DNA.
1952	Martha Chase	Used phages in which the protein was labeled with 35S and the

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 6/5

CLASS: I B Sc BT COURSE CODE: 18BTU201

**COURSE NAME: GENETICS** UNIT: I (Introduction)

BATCH-2018-2021

	Alfred Hershey	DNA with 32P for the final proof that DNA is the molecule of heredity.
1953	Francis Crick James Watson	Solved the three-dimensional structure of the DNA molecule.
1958	Matthew Meselson Frank Stahl	Used isotopes of nitrogen to prove the semiconservative replication of DNA.
1958	Arthur Kornberg	Purified DNA polymerase I from E. coli, the first enzyme that made DNA in a test tube.
1966	Marshall Nirenberg H. Gobind Khorana	Led teams that cracked the genetic code- that triplet mRNA codons specify each of the twenty amino acids.
1970	Hamilton Smith Kent Wilcox	Isolated the first restriction enzyme, HindII, that could cut DNA molecules within specific recognition sites.
1972	Paul Berg Herb Boyer	Produced the first recombinant DNA molecules.
1973	Joseph Sambrook	Led the team at Cold Spring Harbor Laboratory that refined DNA electrophoresis by using agarose gel and staining with ethidium bromide.
1973	Annie Chang Stanley Cohen	Showed that a recombinant DNA molecule can be maintained and replicated in E. coli.
1977	Fred Sanger	Developed the chain termination (dideoxy) method for sequencing DNA.
1983	James Gusella	Used blood samples collected by Nancy Wexler and her co-

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 7/5

CLASS: I B Sc BT COURSE I COURSE CODE: 18BTU201 UNIT: I (Introduction)

COURSE NAME: GENETICS roduction) BATCH-2018-2021

		workers to demonstrate that the Huntington's disease gene is on chromosome 4.
1985	Kary B. Mullis	Published a paper describing the polymerase chain reaction (PCR), the most sensitive assay for DNA yet devised.
1988		The Human Genome Project began with the goal of determining the entire sequence of DNA composing human chromosomes.
1989	Alec Jeffreys	Coined the term DNA fingerprinting and was the first to use DNA polymorphisms in paternity, immigration, and murder cases.
1989	Francis Collins Lap-Chee Tsui	Identified the gene coding for the cystic fibrosis transmembrane conductance regulator protein (CFTR) on chromosome 7 that, when mutant, causes cystic fibrosis.
1990		First gene replacement therapy-T cells of a four-year old girl were exposed outside of her body to retroviruses containing an RNA copy of a normal ADA gene. This allowed her immune system to begin functioning.
1993		FlavrSavr tomatoes, genetically engineered for longer shelf life, were marketed.

### **DEFINITIONS AND TERMS**

- Allele: One alternative of a pair or group of genes that could occupy a specific position on a chromosome.
- Chromosome: A linear strand of DNA harboring many genes.
- **DNA:** Deoxyribonucleic acid; the molecule in which genetic information is encoded.

### CLASS: I B Sc BTCOURSE NAME: GENETICSCOURSE CODE: 18BTU201UNIT: I (Introduction)BATCH-2018-2021

- **Dominant:** An allele producing the same phenotypic effect whether inherited heterozygously or homozygously; an allele that "masks" a recessive allele.
- Gene: A unit of genetic information that occupies a specific position on a chromosome and comes in multiple versions called alleles.
- **Genotype:** The genetic constitution of an organism.
- Heterozygous: Having a genotype with two different and distinct alleles for the same trait.
- Homozygous: Having a genotype with two of the same alleles for a trait.
- **Phenotype:** The physical or observable characteristics of an organism.
- **Recessive:** An allele producing no phenotypic effect when inherited heterozygously and only affecting the phenotype when inherited homozygously; an allele "masked" by a dominant allele.

### **Model organisms**

A model organism is a non-human species that is extensively studied to understand particular biological phenomena, with the expectation that discoveries made in the organism model will provide insight into the workings of other organisms. Model organisms are in vivo models and are widely used to research human disease when human experimentation would be unfeasible or unethical. This strategy is made possible by the common descent of all living organism, and the conservation of metabolic and developmental pathways and genetic material over the course of <u>evolution</u>Studying model organisms can be informative, but care must be taken when extrapolating from one organism to another

In researching human disease, model organisms allow for better understanding the disease process without the added risk of harming an actual human. The species chosen will usually meet a determined taxonomic equivalency to humans, so as to react to disease or its treatment in a way that resembles human physiology as needed. Although biological activity in a model organism does not ensure an effect in humans, many drugs, treatments and cures for human diseases are developed in part with the guidance of animal models. There are three main types of disease models: homologous, isomorphic and predictive. Homologous animals have the same causes, symptoms and treatment options as would humans who have the same disease. Isomorphic

### KARPAGAM ACADEMY OF HIGHER EDUCATION CLASS: I B Sc BT COURSE NAME: GENETICS COURSE CODE: 18BTU201 UNIT: I (Introduction) BATCH-2018-2021

animals share the same symptoms and treatments. Predictive models are similar to a particular human disease in only a couple of aspects, but are useful in isolating and making predictions about mechanisms of a set of disease features.

### Model organisms in genetics

- Many organisms are used in genetic research. Some of the qualities that make an organism a good model for genetic experimentation include:
  - well known genetic history, allowing extrapolation to other related organisms.
  - short life cycle.
  - matings produce large amount of offspring.
  - easy to handle
  - genetic variation must exist between individuals in a population. Without genetic variation, genetic crosses are uninformative.
- Model organisms include representatives from all the main lineages of the tree of life
  - Eukaryote model organisms: Saccharomyces cereviceae, Drosophila melanogaster, Caenorhabditis elegans, Arabidopsis thaliana, Mus musculus, Zebra fish, Neurospora crassa, Tetrahymena, paramecium, Chlamydomonas reinhardtii, Pisum sativum, Zea maize, and Gallus.
  - Prokaryotic model organisms: Most common is *E. coli*, but many others as well, including members of Archaea.
  - 0



#### **Prokaryotic Genetics**

#### Bacterial plasmids and conjugation

Bacteria are typical representatives of prokaryotes. Prokaryotes have cell structure and their DNA is concentrated in a so-called nucleoid region. However, different from eucaryotes, bacteria have neither meiosis nor mitosis.

There are three principal mechanisms of genetic recombination (genetic material transfer into a HOST organism) in bacteria occurring in nature and also in lab conditions: conjugation, transformation and transduction.

CONJUGATION This is the most important mechanism for widespread transfer of genetic information between bacteria. It is the direct transfer of bacterial DNA between organisms and requires CELL-TO-CELL CONTACT.Most conjugation is PLASMID- MEDIATED.

### KARPAGAM ACADEMY OF HIGHER EDUCATION CLASS: I B Sc BT COURSE NAME: GENETICS COURSE CODE: 18BTU201 UNIT: I (Introduction) BATCH-2018-2021

PLASMIDS are small, circular, ds-DNA molecules, measuring about 2-10 kilobases. They all have the ability to replicate in bacteria. Multiple copies of plasmids can be found in a host cell; however, not all plasmids can transfer themselves.

A CONJUGATIVE plasmid has the codes for the genes allowing transfer between cells.

A NONCONJUGATIVE plasmid requires the help of a conjugative plasmid to transmit itself to another bacteria.

NARROW-HOST RANGE PLASMIDS exist only within a single species.

BROAD-HOST RANGE PLASMIDS exist in different genera of organisms.

Plasmids contain the following:

- 1. 1. An ORI or origin of replication that allows autonomous replication.
- 2. 2. One or more genes that confer specific antibiotic resistance.
- 3. 3. Sites for restriction endonucleases which are used for inserting specific DNA fragments.
- 4. 4. Code for virulence factors such as toxins or adhesions



Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 12/5

Plasmids capable of mediating conjugation carry genes coding for the production of a PILUS, a protein appendage which is found on the surface of the donor cells. Bacteria which carry these plasmids are called F(+) cells, making them the donor cells. The tip of the pilus attaches to the surface of the recipient cell, and DNA transfer occurs. It is unclear whether the DNA is transferred through the pilus, or if the pilus serves as the mechanism for holding the two cells together. Either way, this is the reason why it has also been termed BACTERIAL SEX.

Once the recipient cell acquires the F plasmid, it changes from a F(-) to a F(+) cell, thus allowing it to become a donor cell now.

In a very small proportion of cells, the F plasmid is incorporated into the bacterial chromosome. Once inserted, the entire bacterial chromosome acts like a F plasmid. These are then called high frequency recombination strains (Hfr strains). This leads to 2 processes:

- 1. 1. The F plasmid and the entire bacterial DNA undergoes conjugation with an F(-) cell.
- 2. 2. The F plasmid is excised along with a portion of the bacterial DNA; hence the F plasmid is termed F prime (F').

### **Bacterial transformation**

Transformation is a process in which genetic material taken in from the environment is added to a part of the bacterial DNA. The DNA may also replace an existing gene or part of it from the genome of the bacteria, thus resulting in loss of the activity of that gene.

For the first time transformation was demonstrated in 1928 by <u>Frederick Griffith</u>, an English bacteriologist searching for a vaccine against bacterial pneumonia. Griffith discovered that a non-virulent strain of <u>Streptococcus pneumoniae</u> could be transformed into a virulent one by exposure to strains of virulent S. pneumoniae that had been killed with heat. In 1944 it was demonstrated that the transforming factor was genetic, when <u>Oswald Avery</u>, <u>Colin MacLeod</u>, and <u>Maclyn McCarty</u> showed gene transfer in S. pneumoniae. Avery, Macleod and McCarty called the uptake and incorporation of DNA by bacteria "transformation."

Bacterial transformation may be referred to as a stable genetic change brought about by taking up <u>naked DNA</u> (DNA without associated cells or proteins), and <u>competence</u> refers to the state of

being able to take up exogenous DNA from the environment. Two different forms of competence should be distinguished: natural and artificial.

Natural competence Some bacteria (around 1% of all species) are naturally capable of taking up DNA under laboratory conditions; many more may be able to take it up in their natural environments. Such species carry sets of genes specifying machinery for bringing DNA across the cell's membrane or membranes. Natural competence Artificial competence is not encoded in the cell's genes. Instead it is induced by laboratory procedures in which cells are passively made permeable to DNA, using conditions that do not normally occur in nature.

### **Bacterial transduction**

As abovementioned, transduction is one of three mechanisms of genetic recombination in bacteria occuring in nature and also in lab conditions. In other words, transduction is a process, in which nucleic acid from one organism can be transfered into another living organim by means of a intermediate virus.

In other words transduction is the phage-mediated transfer of bacterial DNA.

Phages are bacterial viruses with a DNA molecule enclosed in a protein shell. The structure of the protein shell imposes a limitation on the amount of DNA that can be enclosed by the virus, usually about 50 kilobases.

Most phages carry ds-DNA coiled up inside a protein coat. There may be species of phages which have a ss-DNA or ss-RNA as well.

There are two kinds of transductions:

### Generalized transduction

When phages enter a bacterial cell and replicate, each phage head usually has a copy of the replicated phage genome. Sometimes, an empty phage head is produced. Bacterial DNA may be mistakenly packaged into that empty phage head, then transferred to another bacterial cell. Thus, the DNA that enters the second bacterial cell is a short segment of the chromosome of the first

bacterial cell. As with transformation, recombination must occur for transduction to be successful.

It is termed generalized since the phage will pick up any portion of the bacterial chromosome at random. (It is not specific for a particular chromosomal segment.)

The genes can be transferred only within a certain species because phages usually attack a limited range of bacteria.

Phages can also pick up and transfer plasmid DNA. For example, the penicillinase gene of Staphlococci is located in a plasmid, and the phages can pick up that plasmid and through transduction, pass it on to another cell, thus transferring the penicillanase gene to other staphylococci species.

#### Specialized transduction

Bacteriophages that lyse the host cell are known as virulent phages, and cause a lytic cycle of infection. In contrast, phages that can infect a bacterial cell without causing its death are called temperate phages.

The bacterial cell which survives after phage infection is called a lysogen, and the integrated temperate phage inside of it is termed a prophage.

The phage DNA is then inserted into the bacterial cell DNA and is also replicated as part of the host cell chromosome.

When a bacterial cell is infected by one phage, no other phage can attack the bacteria, because once the phage is integrated into the bacterial cell DNA, it imparts immunity against superinfection by other phages.

The lysogen phase is stable but not permanent. If the prophage is excised during the process of replication, the bacterial cell may eventually be lysed. This can be replicated in the lab by exposure of bacteria to UV light. (Eg., hospital sterilization of rooms with UV.)

In contrast to generalized transduction, temperate phages normally have a specific insertion site on the chromosome and can pick up only a short length of DNA containing a few genes on either side of this site. Thus, it is termed specialized or restrictive transduction.

### The Cell Division Cycle



Cells that stop dividing exit the G1 phase of the cell cycle into a so-called G0 state.

Cells reproduce genetically identical copies of themselves by cycles of cell growth and division. The cell cycle diagram on the left shows that a cell division cycle consists of 4 stages:

- G1 is the period after cell division, and before the start of DNA replication. Cells grow and monitor their environment to determine whether they should initiate another round of cell division.
- S is the period of DNA synthesis, where cells replicate their chromosomes.
- G2 is the period between the end of DNA replication and the start of cell division. Cells check to make sure DNA replication has successfully completed, and make any necessary repairs.
- M is the actual period of cell division, consisting of prophase, metaphase, anaphase, telophase, and cytokinesis.

#### Chromosomes

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 16/5

Chromosomes were first named by cytologists viewing dividing cells through a microscope. The modern definition of a chromosome now includes the function of heredity and the chemical composition. A chromosome is a DNA molecule that carries all or part of the hereditary information of an organism. In eukaryotic cells, the DNA is packaged with proteins in the nucleus, and varies in structure and appearance at different parts of the cell cycle.

Chromosomes condense and become visible by light microscopy as eukaryotic cells enter mitosis or meiosis. During interphase (G1 + S + G2), chromosomes are fully or partially decondensed, in the form of chromatin, which consists of DNA wound around histone proteins (nucleosomes).



In G1, each chromosome is a single chromatid. In G2, after DNA replication in S phase, as cell enter mitotic prophase, each chromosome consists of a pair of identical sister chromatids, where each chromatid contains a linear DNA molecule that is identical to the joined sister. The sister chromatids are joined at their centromeres, as shown in the image below. A pair of sister chromatids is a single replicated chromosome, a single package of hereditary information.

### Ploidy

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 17/5

Humans are **diploid**, meaning we have two copies of each chromosome. We inherited one copy of each chromosome from other mother, and one copy of each from our father. Gametes (sperm cells or egg cells) are **haploid**, meaning that they have just one complete set of chromosomes.

Chromosomes that do not differ between males and females are called **autosomes**, and the chromosomes that differ between males and females are the sex chromosomes, X and Y for most mammals. Humans most commonly have 22 pairs of autosomes and 1 pair of sex chromosomes (XX or XY), for a total of 46 chromosomes. We say that humans have 2N = 46 chromosomes, where N = 23, or the haploid number of chromosomes.

Cells with complete sets of chromosomes are called **euploid**; cells with missing or extra chromosomes are called **aneuploid**. The most common aneuploid condition in people is variation in the number of sex chromosomes: XO (having just one copy of the X), XXX, or XYY. Having no X chromosome results in early embryonic death.

The two copies of a particular chromosome, such as chromosome 1, are called **homologous**. The karyotype image above shows the homologous pairs for all the autosomes. Homologous chromosomes are not identical to each other, unlike sister chromatids. They frequently have different variants of the same hereditary information – such as blue eye color vs brown eye color, or blood type A versus blood type B.

### Mitosis

<u>Mitosis</u> produces two daughter cells that are genetically identical to each other, and to the parental cell. A diploid cell starts with 2N chromosomes and 2X DNA content. After DNA replication, the cells is still genetically diploid (2N chromosome number), but has 4X DNA content because each chromosome has replicated its DNA. Each chromosome now consists of a joined pair of identical sister chromatids. During mitosis the sister chromatids separate and go to opposite ends of the dividing cell. Mitosis ends with 2 identical cells, each with 2N chromosomes and 2X DNA content. All eukaryotic cells replicate via mitosis, except **germline** cells that undergo meiosis (see below) to produce **gametes** (eggs and sperm).

### CLASS: I B Sc BTCOURSE NAME: GENETICSCOURSE CODE: 18BTU201UNIT: I (Introduction)BATCH-2018-2021

- prophase chromosomes condense; each chromosome consists of a pair of identical sister chromatids joined at the centromere.
- metaphase chromosomes line up at the middle of the cell, along the plane of cell division, pushed and pulled by microtubules of the spindle apparatus
- anaphase *sister chromatids separate* and migrate towards opposite ends of the cell
- telophase chromatids cluster at opposite ends of the cell and begin to decondense
- cytokinesis the membrane pinches in to divide the two daughter cells

Here is a simplified diagram illustrating the overall process and products of mitosis:



#### Meiosis

This is a special sequence of 2 cell divisions that produces haploid gametes from diploid germline cells. It starts with a diploid cell that has undergone chromosomal DNA replication: 2N chromosomes, 4X DNA content. Two successive divisions, with no additional DNA replication, results in 4 haploid gametes: 1N chromosomes, 1X DNA content.

NOVA has a good interactive side-by-side comparison of mitosis and meiosis on this page: <u>How</u> <u>cells divide</u>

Meiosis sets the stage for Mendelian genetics. Students need to know that most of the genetics action occurs in the **first** meiotic division:

• homologous chromosomes pair up and align end-to-end (synapsis) in prophase I

- crossing over occurs between homologous chromosomes in prophase I, *before*chromosomes line up at the metaphase plate
- homologous chromosomes separate to daughter cells (sister chromatids do not separate) in the first division, creating haploid (1N) cells
- the separation of each pair of homologous chromosomes occurs independently, so all
  possible combinations of maternal and paternal chromosomes are possible in the two
  daughter cells this is the basis of Mendel's Law of Independent Assortment
- the first division is when daughter cells become functionally or genetically haploid

The last point appears to be the most difficult for students to grasp. Consider the X and Y chromosomes. They pair in prophase I, and then separate in the first division. The daughter cells of the first meiotic division have either an X or a Y; they don't have both. Each cell now has only one sex chromosome, like a haploid cell.

One way of thinking about ploidy is the number of *possible* alleles for each gene a cell can have. Right after meiosis I, the homologous chromosomes have separated into different cells. Each homolog carries one copy of the gene, and each gene could be a different allele, but these two homologs are now in two different cells. Though it looks like there are two of each chromosome in each cell, these are *duplicated* chromosomes; ie, it is one chromosome which has been copied, so there is only one possible allele in the cell (just two copies of it).

The second meiotic division is where sister (duplicated) chromatids separate. It resembles mitosis of a haploid cell. At the start of the second division, each cell contains 1N chromosomes, each consisting of a pair of sister chromatids joined at the centromere.

Here is a simplified diagram illustrating the overall process and products of meiosis:



Meiosis Overview from Wikipedia by Rdbickel. In prophase I, homologous chromosomes pair and separate in the first division (Meiosis I). In Meiosis II, sister chromatids separate.

### It is very important that you recognize how and why cells become haploid after meiosis I.

To confirm for yourself that you understand meiosis, work through one or more of these interactive tutorials:

- The U. Arizona Cell Biology Project's <u>Meiosis tutorial</u> has a click-through animation of meiosis, with 10 thought-provoking problem questions.
- Jung Choi's interactive flash tutorial, programmed by Pearson, uses human chromosome 7, with wild type and cystic fibrosis alleles for CFTR, to track segregation through meiosis, with and without crossing over: <u>Meiotic Segregation tutorial</u>

Chromosomes, chromatids, what is the difference and how many chromosomes are there at different times of the cell cycle and after mitosis and meiosis?

Chromosomes by definition contain the DNA that makes up the fundamental genome of the cell. In a prokaryote, the genome is usually packaged into one circular chromosome consisting of a circular DNA molecule of a few million base pairs (Mbp). In eukaryotes, the genome is packaged into multiple linear chromosomes, each consisting of a linear DNA molecule of tens or hundreds of Mbp. Chromosomes exist at all different phases of the cell cycle. They condense and become visible to light microscopy in prophase of mitosis or meiosis, and they decondense during interphase, in the form of chromatin (DNA wrapped around nucleosomes, like "beads on a string").

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 21/5

### KARPAGAM ACADEMY OF HIGHER EDUCATION CLASS: I B Sc BT COURSE NAME: GENETICS COURSE CODE: 18BTU201 UNIT: I (Introduction) BATCH-2018-2021

The chromosome number, N, in eukaryotes, refers to the number of chromosomes in a haploid cell, or gamete (sperm or egg cell). Diploid cells (all the cells in our body except our gametes) have 2N chromosomes, because a diploid organism is created by union of 2 gametes each containing 1N chromosomes. In terms of chromosome number (ploidy), it's useful to think of chromosomes as packages of genetic information. A pair of sister chromatids is one chromosome because it has genetic information (alleles) inherited from only one parent. A pair of homologous chromosomes, each consisting of a single chromatid in a daughter cell at the end of mitosis, has alleles from the father and from the mother, and counts as 2 chromosomes.

This chromosome number stays the same after chromosome replication during S phase: each chromosome entering cell division now consists of a pair of sister chromatids joined together at the centromere. Then in mitosis, the sister chromatids of each chromosome separate, so each daughter cell receives one chromatid from each chromosome. The result of mitosis is two identical daughter cells, genetically identical to the original cell, all having 2N chromosomes. So during a mitotic cell cycle, the DNA content per chromosome doubles during S phase (each chromosome starts as one chromatid, then becomes a pair of identical sister chromatids during S phase), but the chromosome number stays the same.

A chromatid, then, is a single chromosomal DNA molecule. The number of chromatids changes from 2X in G1 to 4X in G2 and back to 2X, but the number of chromosomes stays the same.

The chromosome number is reduced from 2N to 1N in the first meiotic division, and stays at 1N in the second meiotic division. Because homologous chromosomes separate in the first division, the daughter cells no longer have copies of each chromosome from both parents, so they have haploid genetic information, and a 1N chromosome number. The second meiotic division, where sister chromatids separate, is like mitosis. Chromosome number stays the same when sister chromatids separate.

Using the information above, compare these two simplified diagrams of mitosis and meiosis to visualize why cells are haploid after meiosis I. Specifically, compare the chromosomes in cells at the end of mitosis vs the end of meiosis I, recognizing that the diagram of mitosis tracks just



### Cell cycle and check points

Introduced the cell cycle and the role of microtubules therein. This lecture will discuss the regulatory mechanisms and biochemical checkpoints throughout the cell cycle. Disclaimer: these notes are not my finest work – a lot of this is just a collection of random facts.

Homologous Chromosomes Meiosis II

Meiosis I

### cyclins and cyclin-dependent kinases (CDKs)

CDKs are important master regulators of the cell cycle. Their role is to phosphorylate proteins on either S or T amino acids and thereby regulate the activity of those proteins. Yeast have just

KARPAGAM AG	CADEMY OF HIGHER EDU	JCATION
CLASS: I B Sc BT	COURSE NA	ME: GENETICS
COURSE CODE: 18BTU201	UNIT: I (Introduction)	BATCH-2018-2021

one CDK (Cdk1), while 'metazoans' (animals) like us have nine, of which four are really critical to the cell cycle and will be introduced today.

How are the CDKs themselves regulated? The levels of these proteins remain pretty constant throughout the cell cycle, yet their levels of *activity* rise and fall cyclically. CDKs need to hydrolize ATP for energy in order to perform phosphorylation. They have an ATP binding cleft whose ability to bind ATP is regulated by two mechanisms. First, CDKs have a 'flexible T loop' which contains a threonine (T) residue which normally blocks the ATP binding cleft, but not when the T is phosphorylated. Second, cyclins bind CDKs and induce a conformational change that also helps to expose the ATP binding cleft. Therefore a fully active CDK is one which is both phosphorylated at the T on the T loop *and* is bound to a cyclin.

The various activities of the cell cycle, then, are determined by the combination of cyclins and CDKs that are active at each stage, as shown in the following table:

cell cycle stage	cyclins	CDKs	comments
G1	Cyclin D	CDK4&6	Can react to outside signals such as growth factors or mitogens.
G1/S	Cyclins E & A	CDK2	Regulate centrosome duplication; important for reaching START
S	Cyclins E & A	CDK2	Targets are helicases and polymerases
М	Cyclins A & B	CDK1	Regulate G2/M checkpoint. The cyclins are synthesized during S but not active until synthesis is complete. Phosphorylate lots of downstream targets.

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 24/5

All cyclins contain a conserved 100 amino acid 'cyclin box.' Cyclin/CDK complexes regulate the cell cycle both by promoting activites for their respective stages, and by inhibiting activites for future cell cycle stages that must not yet be reached. Therefore cyclins must be able to be both generated *and* degraded in order for the cell cycle to proceed.



Antibodies against Cyclin D inhibit entry into the S phase, which can be measured by whether BrdU (a fluorescent nucleotide analog) gets incorporated into DNA (which only happens during synthesis).

DNA replication starts at 'prereplication complexes' that get assembled at *origins of replication* during early G1 phase. The S-phase cyclin/CDK complexes phosphorylate these complexes and thereby trigger replication starting from these 'origins'.

Here is a bit more about how the CDKs are regulated through phosphorylation. A CDKactivating kinase (CAK; a trimeric complex composed of CDK-7, Cyclin H and Mat1) phosphorylates amino acid T160 in CDK, located at the T loop, thereby *activating* CDK. CDKs are also regulated by CDK inhibitors p27 (CDKN1B gene), p21 (CDKN1A gene) and p57 (CDKN1C gene), which bind to and inhibit both of the G1 CDKs (CDK4 & CDK6). p27 does this by physically blocking the cyclin/CDK complex's interaction with its targets. p15 (CDKN2B gene) and p16 (CDKN2A gene) are both Ink4s ("inhibitors of kinase 4", though they also inhibit CDK6) control the mid G1 phase by binding to CDK4 and CDK6 and

KARPAGAM ACADEMY OF HIGHER EDUCATION		
CLASS: I B Sc BT	COURSE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: I (Introduction)	BATCH-2018-2021

blocking their binding to cyclin D. This results in decreased phosphorylation of target proteins. Overexpression of p16 arrests the cell cycle by inhibiting CDK4/Cyclin D during early G1.

Another inhibitor called Rb prevents entry into the S phase by binding to E2F transcription factors. E2Fs are transcriptional activators when they act alone but repressors when bound to Rb. The mid-G1 cyclin/CDK complexes partially phosphorylate Rb, reducing its binding to E2Fs; the late G1 complex of CDK2/Cyclin E completely phosphorylates it, preventing its binding to E2F. E2Fs can then act as transcriptional activators for genes needed in the S phase.

S-phase cyclin/CDK complexes accumulate in late G1, but are still bound to an inhibitor called Sic1. But G1 CDK/cyclin complexes polyphosphorylate Sic1 at six sites. When and only when all six sites are phosphorylated (which takes a while), Sic1 is released, whereupon it gets polyubiquitinated and therefore degraded by the proteasome. The S-phase cyclin/CDK complexes are then free to induce DNA replication. This mechanism allows the cell to accumulate many of these complexes 'ahead of time' (starting in G1) but then post-translationally activate them all at once in order to suddenly start massive-scale DNA replication in the S phase.

Targeting of proteins such as Sic1 to the proteasome is mediated by two complexes SCF (Skp, Cullin & F-Box) and APC/C (anaphase promoting complex / cyclosome), each composed of ubiquitin and a protein ligase. These two control three major transiations in the cell cycle:

1. the onset of S phase by degradation of Sic1

2. beginning of anaphase by degradation of securin

3. exit from mitosis by degradation of Cyclin B.

APC/C is composed of several proteins including but not limited to cullin (APC2), Ring (APC11) and SCF-like protein. APC/C's role is to ubiquitinate (add ubiquitin tags to) proteins, thus flagging them for degradation by the proteasome. Cdc20 casues APC/C to polyubiquitinate the anaphase inhibitor securin in point (2) above. Cdh1 causes APC/C to polyubiquitinate Cyclin B in point (3) above. Cdh1 is itself regulated by phosphorylation – the G1/S cyclin/CDK complexes phosphorylate it, preventing it from acting too soon; Cdc14 then later activates Cdh1.

SCF has no binding partner – rather, its ability to ubiquitinate targets (substrates) is determined by the latter's phosphorylation state. SCF is active throughout the cell cycle; the cell regulates its activity by regulating the phosphorylation of SCF's targets. An example is Sic1 (above).

MPF (Maturation or Mitosis Promoting Factor depending on who you ask) is a fancy word for the cyclin B / CDK1 complex, which is active in the M phase - actually, required in order for entry to the M phase. Here's how MPF itself is regulated. CDK1 has two phosphorylation sites (in vertebrates; only one in yeast) at amino acids Y15 and T161. In animals, phosphorylation of Y15 is inhibitory, while phosphorylation of T161 is activating. The cyclin/CDK2 complex begins its life wholly unphosphorylated (thus inactive). then Wee1 phosphorylates Y15, CAK then phosphorylates T161 too, then finally Cdc25 removes the Y15 phosphorylation, allowing the complex to be active (about 100x more activity than with neither site phosphorylated). Cdc25 overexpression results in premature removal of this Y15 phosphorylation so that the cell enters the M phase before it has had enough time to grow during the G2 phase. This results in 'wee cells' which are literally small because the cell growth was inhibited, so that the daughter cells after mitosis are smaller than the mother cell was. (Hence the name Wee1, which is of course backwards since Wee1 acts against the 'wee cell' phenotype).

Once a cell passes the restriction point (R) in late G1, it is committed to passing through the S phase.

Many cancers involve a loss of p16 leading to cyclin D1 overexpression. Other cancers (esp. late onset cancers – breast, lung, bladder) often have hyperphosphorylation of Rb leading to release of E2F to promote the cell cycle. This hyperphosphorylation can be caused by loss or 'misexpression' of one or more things

### Regulatory steps in the cell cycle

Here is an overview of what are considered to be the 9 fundamental steps of cell cycle regulation. *Cell starts in phase G0 or G1.* 

- 1. DNA replication machinery begins to assemble at origins of replication
- 2. G1 cyclin-CDK complexes inactivate Cdh1
- 3. G1 cyclin-CDK complexes activate the S-phase cyclin-CDK expression

- 4. G1 cyclin-CDK complexes phosphorylate and thereby inactivate S-phase inhibitor(s?)
- 5. SCF polyubiquitinates the phosphorylated S-phase inhibitor, targeting it for proteasome degradation. *Cell enters S phase*.
- 6. S-phase cyclin/CDK activates the preplication complexes that began to assemble in step 1. *S phase and G2 phase happen.*
- 7. Cdc25 phosphatase activates M-phase cyclin/CDKs. Cell enters M phase and gets to metaphase.
- 8. APC/C and Cdc20 target securin for proteasomal degradation. *Cell advances from metaphase to anaphase.*
- 9. CdcA phosphatase activates Cdh1, making it possible for APC/C and Cdh1 to target the M-phase cyclins for proteasomal degradation. *Cell returns from M phase to G0 or G1 phase.*

Step 7 relies on a mechanism to recognize unreplicated DNA and stalled replication forks in order to ensure that mitosis does not proceed before all DNA has been replicated. Two proteins are involved in this mechanism: ATR and Chk1 (pronounced 'check one'). Both are protein kinases that 'sense' the state of DNA. ATR is located at the replication fork and activates other protein kinases, leading to phosphorylation and therefore activation of Chk1 which is also a kinase. Chk1 then phosphorylates and *in*activates Cdc25, which would otherwise activate MPF. ATR thus continues to prevent M phase entry until replication is complete (= when replication forks are gone?).

Step 8 involves an even more complicated mechanism called the mitotic spindle assembly checkpoint which prevents entry into anaphase until each and every kinetochore is attached to a microtubule. Sister chromatids are held together at the centromeres by a multi-protein complex called cohesin (composed of smc1, smc3 and kleisin). Separase is an enzyme that cleaves part of cohesin kleisin, physically releasing the chromatids. Securin inhibits separase to prevent separation of the chromatids. Once all kinetochores are attached, securin is targeted for degradation, thus freeing separase from inhibition and causing entry into anaphase. The mechanism for 'knowing' when to target securin for degradation is the really complicated part. It involves Mad1, Mad2, Cdc20, APC/C. Mad2 can have an 'open' or 'closed' conformation. 'closed' Mad2 inactivates Cdc20, thus preventing its association with APC/C. This pretty Wikimedia commons graphic by Dawn08 gives some sense of how complicated it all is:

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 28/5



Proper segregation of the daughter chromosomes is monitored by the 'mitotic exit network'. At this stage, MPF is inactivated, Cdh1 is dephosphorylated, and the cell can enter telophase.

There is also a 'spindle position checkpoint'. In yeast, Tem1 becomes associated with the spindle pole body at the centrosome.

'DNA damage checkpoints' detect whether DNA has been damaged (e.g. by UV light, chemicals, etc.). The cell cycle arrests in G1 or S, preventing the copying of damaged bases until they can be repaired. Arrest in G2 allows double-stranded breaks to be repaired before mitosis. Tumor suppressor genes are important in preventing replication of damaged DNA. Cells can sense UV or gamma radiation damage through a protein called ATM/R. ATM/R phosphorylates and activates Chk2, which then phosphorylates Cdc25, marking it for degradation. ATM/R also stabilizes the tumor suppressor p53 (gene: TP53), which is a

Prepared by Dr. Anil Kumar PK and Dr. Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 29/5

transcription factor that activates p21, which (inhibits the G1 CDKs? and) arrests the cell cycle in G1. p53 undergoes loss-of-function mutations in perhaps 50% of cancers, and many of the mutations are dominant negative so only one allele need be mutated. Mdm2 also regulates p53, and ATM/R prevents Mdm2's binding to p53. p53 is regulated by multiple means, and also has multiple roles in promoting DNA repair, cell cycle arrest, and apoptosis

#### CLASS: I B Sc BT COURSE CODE: 18BTU201

COURSE NAME: GENETICS UNIT: II BATCH-2018-2021

### UNIT-II

### **SYLLABUS**

**GENETICS:** Mendels experiment design, Mono hybrid, di-hybrid and tri-hybrid crosses, Law of segregation & principle of independent assortment, Verification of segregates by test and back crosses, Chromosomal theory of inheritance, allelic interaction concepts of dominance, recessiveness, Concepts of incomplete dominance , co-dominance, Concepts of semi dominance, pleiotropy, Concepts of multiple allele, pseudo-allelic,

### **MENDELIAN GENETICS**

### Mendel's Laws of Inheritance

Gregor Mendel is a Father of Genetics. His work, entitled "Experiments on Plant Hybrids" was published in 1866. Like many great scientific discoveries, it was ignored for 34 years. In 1900, Mendel's work was rediscovered by three botanists working independently

- Hugo de Vriies of Holand
- Carll Correns off Germany
- Eriich von Tschermak of Austria.

#### **Mendel's Experiments**

Gregor Mendel, the Austrian monk famous for his experiments with pea plant characteristics, was the first to identify discrete units of heredity and thus discredit the blending theory. Mendel used characteristics of pea plants and four o'clock flowers to analyze the hereditary patterns of these traits. His historic experiments led him to the conclusion that inherited characteristics were carried in discrete, independent units (later named genes). In Mendel's interpretation, hereditary characteristics occurred in pairs of factors that had specific relationships. Mendel devised two fundamental principles of inheritance:
- Mendel's Principle of Segregation: The factors of inheritance (genes) normally are paired, but are separated or segregated in the formation of gametes (eggs and sperm).
- Mendel's Principle of Independent Assortment: Each factor's distribution in the gametes is not related to the distribution of any other factor. (This principle is not strictly true due to the organization of genes on chromosomes.)

Mendel also defined and described the relationships between the different factors of inheritance and their effects on the observed characteristics of the organism.



#### **Independent** assortment

### **Mendel's Observations**

Mendel made numerous important observations in his exhaustive study of pea plants' characteristics. He elaborated an important distinction between dominant and recessive traits through his work with pea plants.

By studying the characteristics of pea plants, such as their height, seed shape, seed color, flower position, and other traits this discussion will use height as a primary example. Mendel first crossbred one tall, true-breeding plant with one short, true-breeding plant. Contrary to the blending theory, all the offspring were tall. In terms of genotype, the original tall plant was TT (two dominant alleles; homozygous), the short plant was tt (two recessive alleles; homozygous), and the second-generation plants were Tt (one dominant and one recessive allele; heterozygous).

When Mendel next allowed these plants to self-fertilize, he found that the short trait reappeared in the third generation. The ratio of short to tall plants was almost exactly 3:1. Their genotypes were as follows - 1 short (tt) : 2 tall (Tt) : 1 tall (TT).

### **Punnett Squares: Simulation**

These results can be simulated with a Punnett square, a calculation device used to determine the possible genotypes of offspring given the genotypes of the parents. The parents' genotypes are represented by letters, one allele in each cell of the upper row (traditionally the mother) and rightmost column (traditionally the father). The offspring's genotypes are then calculated by observing the intersection of the mother's and father's individual alleles (mush like a multiplication table). Use the interactive Punnett square below to simulate Mendel's results or experiment with combinations of your own. (If you do not select a genotype, the simulation will assume Mendel's original cross: a true-breeding short plant with a true-breeding tall plant.)

### Keys to Mendel's Experiments

Prepared by Dr. Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 3/5

CLASS: I B Sc		KSE NAME: GENETICS
COURSE CODE: 18BTU2	01 UNIT: II	BATCH-2018-2021
The garden pea was an ideal organ	nism for study because:	
– Vigorous growth		
– Self fertilization		
– Easy to cross fertilize		<u>_</u>
– Produces large number of offspr	ring each generation	
Mendel analyzed traits with discre	rete alternative forms (one	of two options)
– purple vs. white flowers		
– yellow vs. green peas		
- round vs. wrinkled seeds		
– long vs. short stem length		
Mendel established pure/true breed	ding lines to conduct his e	xperiments.
(traits remain constant from generation)	ration to generation)	
Monohybrid Crosses and Mendel'	's Principle of Segregation	
Terminology used in breeding exp	periments:	
- Parental generation is the P gene	eration.	
- True breeding unless stated othe	erwise	
- Progeny of P generation is the fi	irst filial generation,	
" designated F1.		
- When F1 interbreed or are "selfe	ed",	
- the second filial generation, F2,	is produced.	

Prepared by Dr. Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE Pa

## – F2 is always F1 x F1

Subsequent interbreeding produces F3, F4, and F5 generations

## Mendel's First Three Postulates:

Using the consistent pattern of results in the monohybrid crosses, Mendel derived the following three postulates or principles of inheritance. The Monohybrid cross reveals how one trait is transmitted from generation to generation.

**Unit Factors in Pairs:** Genetic characters are controlled by unit factors that exist in pairs in individual organisms.

.**Dominance/Recessiveness:** When two unlike unit factors responsible for a single character are present in a single individual, one unit factor is dominant to the other, which is said to be recessive.

**.Segregation:** During the formation of gametes, the paired unit factors separate or segregate randomly so that each gamete receives one or the other with equal likelihood.

## Monohybrid Crosses and Mendel's Principle of Segregation:

A monohybrid cross involves true-breeding strains that differ in a single trait. To determine whether both parents contribute equally to the phenotype of a particular trait in offspring a set of reciprocal crosses is performed. In Mendelian genetics, offspring of a monohybrid cross will exactly resemble only one of the parents.

Mendel noted that traits that disappear in the F1 reappear in the F2. The F2 has a ratio of about three individuals with the phenotype of the F1 to one individual with the

"reappearing" phenotype

– 3:1 ratio

- 3/4 to 1/4

Mendel reasoned that information to create the trait was present in the F1 in the form of "factors". now called genes.

By convention, letters may be used to designate alleles,

– with the dominant a capital letter (S)

– and the recessive in lowercase (s).

Individuals with identical alleles are said to be homozygous for that gene

- (e.g., genotypes SS and ss)

because all their gametes will have the same allele for this trait.
Individuals with different alleles are heterozygous (e.g., Ss).

- because 1/2 of their gametes will contain one allele, and 1/2 the

Other.When Mendel had conducted experiments for the seven different traits in garden peas he made these conclusions:

1. Results of reciprocal crosses are always the same.

2. The F1 resembled only one of the parents.

3. The trait missing in the F1 reappeared in about ¼ of the F2 individuals.

## **Punnett Squares**

A Punnett square is a grid that enables one to predict the outcome off simple genetic crosses proposed by the English geneticist, Reginald Punnett.

1. Write down the genotypes of both parents

- Malle parentt = *Ttt*
- Femalle parentt = Ttt

Prepared by Dr. Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE Pa

CLA55: 1	I B Sc BT		COUR	SE NAME: GENETICS
COURSE CODE: 18	BTU201	UN	IT: II	BATCH-2018-2021
2. Write down the possible	gametes each	parent c	an make.	
– Malle gamettes: <i>T</i> or <i>tt</i>				
– Femalle gamettes: <i>T</i> or <i>tt</i>				
3. Fill in the Punnett square	e with the pos	ssible gei	notypes of tl	he offspring. Determine the relative
proportions of genotypes an	d phenotypes	of the o	ffspring	
– Genotypic ratio				
"TT : Tt : tt				
" 1 : 2 : 1				
– Phenotypic ratio				
" Tallll :: dwarff				
" 3 :: 1				
	Mo	onohybri	id cross	
	Don (1	ninant and 1 = Tall &	t = short	
		Cross:	Tt × Tt	
	Г	1	· ]	
	т	тт	π	
	. T	Tt	#	

## The Principle of Segregation

The first Mendelian law, the principle of segregation, states:

- "Recessive characters, which are masked in the F1 from a cross between two true-breeding strains, reappear in a specific proportion in the F2." This is because alleles segregate during anaphase I of meiosis and progeny are then produced by random combination of the gametes.

## **Test cross:**

A **test cross** is a cross of the F1 plant to a homozygous recessive plant. Test crosses help establish genotype.

## Mendel's Dihybrid Cross Generated a Unique F2 Ratio:

**Independent assortment**: *During gamete formation, segregating pairs of unit factors assort independently of each other.* 

As a natural extension of the monohybrid cross, Mendel also designed experiments in which he examined two characters simultaneously. Such a cross, involving two pairs of contrasting traits, is a **dihybrid cross** or *two-factor cross*.

For example, if pea plants having yellow seeds that are round are bred with those having green seeds that are wrinkled, the results shown in will occur: the F1 offspring will all be yellow and round. It is therefore apparent that yellow is dominant to green and that round is dominant to wrinkled. When the F1 individuals are selfed, approximately of the F2 plants express yellow and round, express yellow and wrinkled, express green and round, and express green and wrinkled. Instead of crossing one P1 parent with both dominant traits (yellow, round) and one with both recessive traits (green, wrinkled), plants with yellow, wrinkled seeds are crossed with those with green, round seeds. In spite of the change in the P1 phenotypes, both the F1 and F2 results remain unchanged.



## **APPLICATIONS OF MENDEL'S PRINCIPLES:**

The reason for these laws is found in the nature of the cell nucleus. It is made up of several chromosomes carrying the genetic traits. In a normal cell, each of these chromosomes has two parts, the chromatids. A reproductive cell, which is created in a process called meiosis, usually contains only one of those chromatids of each chromosome. By merging two of these cells (usually one male and one female), the full set is restored and the genes are mixed. The resulting cell becomes a new embryo. The fact that this new life has half the genes of each parent (23 from mother, 23 from father for total of 46 in the case of humans) is one reason for the Mendelian laws. The second most important reason is the varying dominance of different genes, causing some traits to appear unevenly instead of averaging out (whereby dominant doesn't mean more likely to reproduce—recessive genes can become the most common, too).

There are several advantages of this method (sexual reproduction) over reproduction without genetic exchange:

Prepared by Dr. Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 9/5

- 1. Instead of nearly identical copies of an organism, a broad range of offspring develops, allowing more different abilities and evolutionary strategies.
- 2. There are usually some errors in every cell nucleus. Copying the genes usually adds more of them. By distributing them randomly over different chromosomes and mixing the genes, such errors will be distributed unevenly over the different children. Some of them will therefore have only very few such problems. This helps reduce problems with copying errors somewhat.
- 3. Genes can spread faster from one part of a population to another. This is for instance useful if there's a temporary isolation of two groups. New genes developing in each of the populations don't get reduced to half when one side replaces the other, they mix and form a population with the advantages of both sides.
- 4. Sometimes, a mutation can have positive side effects. For example, sickle cell anemia is a mutation that can cause the benefit of malaria resistance. The mechanism behind the Mendelian laws can make it possible for some offspring to carry the advantages without the disadvantages until further mutations solve the problems

## NON-MENDELIAN GENETICS

Non-Mendelian inheritance is a general term that refers to any pattern of inheritance in which traits do not segregate in accordance with Mendel's laws. These laws describe the inheritance of traits linked to single genes on chromosomes in the nucleus. In Mendelian inheritance, each parent contributes one of two possible alleles for a trait. If the genotypes of both parents in a genetic cross are known, Mendel's laws can be used to determine the distribution of phenotypes expected for the population of offspring. There are several situations in which the proportions of phenotypes observed in the progeny do not match the predicted values.

Although inheritance of traits in fungi, viruses, and bacteria are all non-Mendelian, the phrase "non-Mendelian inheritance" is usually only used to describe the exceptions which occur in eukaryotic reproduction

### **Incomplete Dominance:**

**Definition:** Incomplete dominance is a form of intermediate inheritance in which one allele for a specific trait is not completely dominant over the other allele. This results in a combined phenotype.

### **Incomplete dominance:**

Incomplete dominance occurs when the phenotype of the heterozygous genotype is an intermediate of the phenotypes of the homozygous genotypes. For example, the snapdragon flower color is either homozygous for red or white. When the red homozygous flower is paired with the white homozygous flower, the result yields a pink snapdragon flower. The pink snapdragon is the result of incomplete dominance. A similar type of incomplete dominance is found in the four o'clock plant where in pink color is produced when true bred parents of white and red flowers are crossed. When plants of  $F_1$  generation is self pollinated the phenotypic and genotypic ratio of the  $F_2$  generation will be same and is 1:2:1



### **Over-dominance:**

It is a condition in genetics where the phenotype of the heterozygote lies outside of the phenotypical range of both homozygote parents. Over dominance can also be described as heterozygote advantage, wherein heterozygous individuals have a higher fitness than homozygous individuals.

An example in humans is sickle cell anemia. This condition is determined by a single polymorphism. Possessors of the deleterious allele have lower life expectancy, with homozygotes rarely reaching 50 years of age. However, this allele also yields some resistance to malaria. A thus in region where malaria exerts or has exerted a strong selective pressure, sickle cell anemia has been selected for its conferred partial resistance to the disease. While homozygotes will have either no protection from malaria or a dramatic propensity to sickle cell anemia, heterozygotes enjoy a partial resistance to both

### **Codominance:**

It is a situation in which two different alleles for a genetic trait are both expressed. Codominance is a relationship between two versions of a gene. Individuals receive one version of a gene, called an allele, from each parent. If the alleles are different, the dominant allele usually will be expressed, while the effect of the other allele, called recessive, is masked. In codominance, however, neither allele is recessive nor are the phenotypes of both alleles expressed.

A slightly more complicated multiple-allele system determines blood type in humans. The three alleles are: A, B, and O, corresponding to A, B, and O type blood respectively. The A and B alleles are dominant over the O allele, which is always recessive. However, there is an additional twist: the A and B alleles exhibit incomplete dominance and, when inherited together, give rise to AB blood type. Yet another catch is waiting, however: humans also have the Rh factor, a special type of protein found in most (but not all) human blood. Rh factor's presence is inherited in the ordinary Mendelian fashion, with Rh-positive dominant over Rh-negative. In total, this means humans have 8 simple blood types.



**Epistasis:** it is an interaction between genotypes at two *different* gene loci, which sometimes resembles a dominance interaction at a single locus. Epistasis modifies the characteristic 9:3:3:1 ratio expected for two non-epistatic genes. Most genetic systems involve complex epistatic interactions among multiple gene loci.

- 1. Recessive epistasis
- 2. Dominant epistasis
- 3. Supplementary epistasis

## **Multiple alleles**

Although any individual has at most two different alleles, most genes exist in a large number of allelic forms in the population as a whole. In some cases, the alleles have different effects on the phenotype, and their dominance interactions with each other can be described as a series.

For example, the best known human blood groups, the <u>ABO system</u>, comprises three sets of alleles at the I locus,  $I^A$ ,  $I^B$ , and  $I^O$ . The first two are dominant to the latter: that is, the **AA** and

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B Sc BT	COUR	<b>SE NAME: GENETICS</b>		
COURSE CODE: 18BTU201	UNIT: II	BATCH-2018-2021		

**AO** genotypes produce indistinguishable blood group phenotypes, called "*Type A*", as do **BB** and **BO**, which produce "*Type B*" blood. In another example, coat color in <u>siamese cats</u> and related breeds is determined by a series of alleles at the <u>albino</u> gene locus (*c*) that produce different levels of pigment and hence different levels of color dilution. Four of these are  $c^+$ ,  $c^b$ ,  $c^s$ , and  $c^a$  (standard, <u>Birman</u>, <u>siamese</u>, and <u>albino</u>, respectively), where the first allele is completely dominant to the last three, and the last is completely recessive to the first three.

## PLEIOTROPY AND LETHAL ALLELES

## **Introduction**

From Mendel's experiments, you might imagine that all genes control a single characteristic and affect some harmless aspect of an organism's appearance (such as color, height, or shape). Those predictions are true for some genes, but definitely not all of them! For example:

- A human genetic disorder called Marfan syndrome is caused by a mutation in one gene, yet it affects many aspects of growth and development, including height, vision, and heart function. This is an example of **pleiotropy**, or one gene affecting multiple characteristics.
- A cross between two heterozygous yellow mice produces yellow and brown mice in a ratio of 2:12:12, colon, 1, not 3:13:13, colon, 1. This is an example of **lethality**, in which a particular genotype makes an organism unable to survive.

In this article, we'll take a closer look at pleiotropic genes and lethal alleles, seeing how these variations on Mendel's rules fit into our modern understanding of inheritance.

## **Pleiotropy**

When we mentioned Mendel's <u>experiments</u> with purple-flowered and white-flowered plants, we didn't discuss any other phenotypes associated with the two flower colors. However, Mendel noticed that the flower colors were always correlated with two other features: the color of the seed coat (covering of the seed) and the color of the axils (junctions where the leaves met the stem) $^{1,2}1,2$ start superscript, 1, comma, 2, end superscript.

In plants with white flowers, the seed coats and axils were colorless. In plants with purple flowers, on the other hand, the seed coats were brown-gray and the axils were reddish. Thus, rather than affecting just one characteristic, the flower color gene actually affected three.

Genes like this, which control multiple, seemingly unrelated features, are said to be **pleiotropic** (*pleio-* = many, *-tropic* = effects)^11start superscript, 1, end superscript. We now know that Mendel's flower color gene specifies a protein that causes colored particles, or pigments, to be made^22start superscript, 2, end superscript. This protein works in several different parts of the pea plant (flowers, seed coat, and leaf axils). In this way, the seemingly unrelated phenotypes can be traced back to a defect in one gene with several jobs.



Simple schematic illustrating pleiotropy.

In pleiotropy, one gene affects multiple features (feature 1, feature 2, feature 3.

Caption: One gene affects multiple characteristics.

Importantly, alleles of pleiotropic genes are transmitted in the same way as alleles of genes that affect single traits. Although the phenotype has multiple elements, these elements are specified as a package, and the dominant and recessive versions of the package would appear in the offspring of two heterozygotes in a ratio of 3:13:13, colon, 1.

### Pleiotropy in human genetic disorders

Genes affected in human genetic disorders are often pleiotropic. For example, people with a hereditary disorder called Marfan syndrome may have a set of seemingly unrelated symptoms, including the following^{1,3}1,3start superscript, 1, comma, 3, end superscript:

- Unusually tall height
- Thin fingers and toes
- Dislocation of the lens of the eye
- Heart problems (in which the aorta, the large blood vessel carrying blood away from the heart, bulges or ruptures).

These symptoms don't seem directly related, but as it turns out, they can all be traced back to the mutation of a single gene. This gene encodes a protein that assembles into chains, making elastic fibrils that give strength and flexibility to the body's connective tissues^44start superscript, 4, end superscript. Mutations that cause Marfan syndrome reduce the amount of functional protein made by the body, resulting in fewer fibrils.

How does the identity of this gene explain the range of symptoms? Our eyes and the aortas normally contain many fibrils that help maintain structure, which is why these two organs are affected in Marfan syndrome^55start superscript, 5, end superscript. In addition, the fibrils serve as "storage shelves" for growth factors. When there are fewer of them in Marfan syndrome, the growth factors cannot be shelved and thus cause excess growth (leading to the characteristic tall, thin Marfan build)^44start superscript, 4, end superscript.

### **Lethality**

For the alleles that Mendel studied, it was equally possible to get homozygous dominant, homozygous recessive, and heterozygous genotypes. That is, none of these genotypes affected the survival of the pea plants. However, this is not the case for all genes and all alleles.

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B Sc BT	COUR	SE NAME: GENETICS		
COURSE CODE: 18BTU201	UNIT: II	BATCH-2018-2021		

Many genes in an organism's genome are needed for survival. If an allele makes one of these genes nonfunctional, or causes it to take on an abnormal, harmful activity, it may be impossible to get a living organism with a homozygous (or, in some cases, even a heterozygous) genotype.

## **Example: The yellow mouse**

A classic example of an allele that affects survival is the lethal yellow allele, a spontaneous mutation in mice that makes their coats yellow. This allele was discovered around the turn of the 20th century by the French geneticist Lucien Cuenót, who noticed that it was inherited in an unusual pattern<sup>{6,7}6,7</sup>start superscript, 6, comma, 7, end superscript.

When yellow mice were crossed with normal agouti (brown) mice, they produced half yellow and half brown offspring. This suggested that the yellow mice were heterozygous, and that the yellow allele, A^YAYA, start superscript, Y, end superscript, was dominant to the agouti allele, AAA. But when two yellow mice were crossed with each other, they produced yellow and brown offspring in a ratio of 2:12:12, colon, 1, and the yellow offspring did not breed true (were heterozygous). Why was this the case?

COURSE CODE: 18BTU2	01 UI	NIT: II	BATCH-2018-2021
•	A <sup>r</sup> A	× Ara	2
	A	A	
A۲	dies as embryo AYAY	AYA	
A	AYA	AA	
	2 yellow among :	: I brown survivors	

Two yellow mice (A<sup>A</sup>YAAYAA, start superscript, Y, end superscript, Agenotype) are crossed to one another. The Punnett square for the cross is:

	A^Y <i>AY</i> <b>A</b> , start superscript, <b>Y</b> , end superscript	AAA
A^Y <i>AY</i> A, start superscript, Y, end superscript	A <sup>Y</sup> <i>AY</i> A, start superscript, Y, end superscriptA <sup>Y</sup> <i>AY</i> A, start superscript, Y, end superscript(dies as embryo)	A <sup>A</sup> Y <i>AY</i> A, start superscript, Y, end superscriptAAA (yellow)
AAA	A <sup>Y</sup> <i>AY</i> A, start superscript, Y, end superscriptAAA(yellow)	AAAAA(agouti/brown)

There is a phenotypic ratio of 2:1 yellow:brown among the mice that survive to birth.

As it turned out, this unusual ratio reflected that some of the mouse embryos (homozygous A<sup>Y</sup>A<sup>Y</sup>AYAYA, start superscript, Y, end superscript, A, start superscript, Y, end superscript genotype) died very early in development, long before birth. In other words, at the level of eggs, sperm, and fertilization, the color gene segregated normally, resulting in embryos with a 1:2:11:2:11, colon, 2, colon, 1 ratio of A<sup>Y</sup>A<sup>Y</sup>AYAYA, start superscript, Y, end superscript, A, start superscript, Y, end superscript, A, start superscript, Y, end superscript, A, and AAAAA, A genotypes. However, the A<sup>Y</sup>A<sup>Y</sup>AYAYA, start superscript, Y, end superscript, A, start superscript, Y, end superscript, Y, end superscript, A, start superscript, Y, end superscript, Y, end superscript, A, start superscript, Y, end superscript, A, start superscript, Y, end superscript, Y, end superscript, A, start superscript, Y, end superscript, Y, end superscript, A, start superscript, Y, end superscript, Y, end superscript, A, start superscript, Y, end superscript mice died as tiny embryos, leaving a 2:12:12, colon, 1 genotype and phenotype ratio among the surviving mice<sup>4</sup>7,8}7,8start superscript, 7, comma, 8, end superscript.

Alleles like A<sup>Y</sup>*AY*A, start superscript, Y, end superscript, which are lethal when they're homozygous but not when they're heterozygous, are called **recessive lethal alleles**.

## PENETRANCE and EXPRESSIVITY

Egg colour of the endangered Hawai'ian <u>Oo'Aa bird</u> is controlled by a single locus, *BLU*, where the B allele is dominant to the ballele. As part of the recovery strategy, a purebreeding blue-egg bird (BB) from one island is crossed with a pure-breeding white-egg bird (bb) from another island. Because all the offspring are Bb heterozygotes, they are all expected to show a uniform, blue phenotype like that of the BB parent. However, if penetrance and / or expressivity vary, three patterns of variation that differ from expectation are possible.

If penetrance varies [left], the Bb genotype sometimes does not *penetrate through to the* phenotype: some Bb eggs are blue just like those of their BB parents, others are white like those of their bb parents, despite presence of the B allele. "Lack of penetrance" is sometimes used to explain individuals whose phenotypes do not reflect their inferred genotypes, e.g., a dominant trait "skip" generation pedigree that appears to a in a If expressivity varies [middle], the Bb genotype will not be *expressed uniformly in the* phenotype: all eggs are blue, but the exact shade of blue varies among individuals with the same genotype. Variable expressivity is frequently attributable to environmental factors or variation at

Prepared by Dr. Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE Page 19/5

	CLASS:	I B Sc BT	COUR	SE NAME: GENET	ICS
COU	RSE CODE: 18	8BTU201	UNIT: II	BATCH-2018-	2021
gene	loci	elsewhere	in	the	genome
If express	sivity and pene	trance both vary	[right], there	may be a continue	ous gradient c
phenotypes 1	between white	and blue eggs. No	ote that lack of	penetrance can be c	onsidered as a
extreme fron	n of expressivit	y, in which the ran	ige of expressio	n includes non-expre	ession.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

### CLASS: I B Sc BT COURSE CODE: 18BTU201

COURSE NAME: GENETICS UNIT: III BATCH-2018-2021

## <u>UNIT-III</u>

## **SYLLABUS**

**GENETICS:** Eukaryotic nuclaear genome, Nucleotide sequence composition-unique & repetitive DNA, Satellite DNA, Centromere and telomere DNA, Respective transposed sequences, Genetic organization of prokaryotic and viral genome, Structure and characteristics of bacterial and eukaryotic chromosomes

## CHROMOSOME STRUCTURE

During nuclear division, the DNA (as chromatin) in a Eukaryotic cell's nucleus is coiled into very tight compact structures called chromosomes. These are rod-shaped structures made of protein and DNA, which are visible (when stained) only during nuclear division.

The DNA in Eukaryotic cells is coiled tightly around proteins called histones, which help in the tight packing of DNA. During interphase, the DNA is not tightly coiled into chromosomes, but exists as chromatin.

When preparing for nuclear division, during the S phase of interphase, the chromosomes copy themselves (i.e. DNA replication occurs). Each half of the chromosome is now called a chromatid. Note that there is still only one chromosome; it consists of two chromatids but has only one centromere. The function of this is to hold the two chromatids together until they separate during anaphase.

Chromosomes are simpler in Prokaryotes. Their DNA is in a single chromosome, and exists as a loop (ccc DNA).

## **Eukaryotic Genomes**

The survival strategy for most bacteria is centered on metabolic adaptability and a short doubling time. Many bacteria can double in less than twenty minutes under ideal conditions. Bacterial genomes, then, must contain enough information to specify a broad array of biosynthetic and catabolic systems, but must also be small enough for rapid duplication. Multicellular eukaryotes are not as dependent on rapid doubling as are bacteria. For most, the

generation time is much longer than the time required to replicate the genome, so that this step is not rate-limiting as it is in bacteria. Consequently, there is less selective pressure for the control of genome size in most eukaryotes. We have already seen that introns represent a part of the genome that does not specify protein sequence. Other **non-coding** genomic elements include **satellite DNA**, **telomeres**, **transposons**, and **pseudogenes**.

The moderately repetitive DNA includes those sequences which have multiple copies in the genome, designed to increase the rate or amount of gene product, and some regulatory sequences found scattered throughout the genome. Sequences such as the ribosomal RNA and transfer RNA, which are required in large amounts for protein synthesis have many copies on the genome. The histone sequences also have large copy numbers in the genome. Another group of moderately repetitive DNA sequences are those scattered throughout the genome; known as SINES (short interspersed elements) and LINES (long interspersed elements). Some famous SINES and LINES: Alu repeats are the major SINE in mammalian genomes. They are ~300 bp long and about a million of such sequences exist scattered throughout the genome. They account for ~10% of the genome. They are transcribed into RNA but have no known function?? They are known as Alu repeats because they contain the recognition sequence for the restriction enzyme Alu. The most common LINE in the human genome is L1, a 6 000 bp sequence which is repeated some 50 000 times in the human genome. L1 sequences are also transcribed and some even encode proteins! Their function in the cell is unknown. Both the Alu and L1 sequences are transposable elements, capable of moving to different sites in the genome. The group of sequences with a small number of copies on the genome include such sequences as the globins. This family of genes contains a number of closely related sequences, varying by only a few bases in the code, will cross hybridise. These are also known as gene clusters. The final group, the single copy sequences make up the vast majority of genes on the genome (gene being a functional unit which codes for a single polypeptide chain). This group is the most complex and takes the longest to re-anneal, hence the log scale on the time. The highly repetitive DNA reanneals in seconds while the most complex single copy group takes hours or days to re-anneal.

## **Chromosome Structure of Prokaryotes: (Bacteria)**

In contrast to the linear chromosomes found in eukaryotic cells, the strains of bacteria initially studied were found to have single, covalently closed, circular chromosomes. The circularity of

Prepared by Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE

the bacterial chromosome was elegantly demonstrated by electron microscopy in both Gram negative bacteria (such as *Escherichia coli*) and Gram positive bacteria (such as *Bacillus subtilis*). Bacterial plasmids were also shown to be circular. In fact, the experiments were so beautiful and the evidence was so convincing that the idea that bacterial chromosomes are circular and eukaryotic chromosomes are linear was quickly accepted as a definitive distinction between prokaryotic and eukaryotic cells. However, like most other distinctions between prokaryotic cells, it is now clear that this dichotomy is incorrect. Not all bacteria have a single circular chromosome: some bacteria have multiple circular chromosomes, and many bacteria have linear chromosomes and linear plasmids.



Experimental evidence for multiple chromosomes and linear chromosomes initially came from studies using pulsed field gel electrophoresis (PFGE), an approach that uses alternating electric fields to separate large DNA molecules on an agarose gel. Subsequently genome sequencing projects have added to the list of bacteria with multiple or linear chromosomes.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B Sc BT COURSE CODE: 18BTU201

### COURSE NAME: GENETICS UNIT: III BATCH-2018-2021

Bacteria	Chromosome(s)	Plasmid(s)
Agrobacterium tumefaciens	one linear (2.1 Mb) + one circular (3.0 Mb)	two circular (450 + 200 Kb)
Bacillus subtilis	one circular (4.2 Mb)	
Bacillus thuringiensis	one circular (5.7 Mb)	six (each >50 Kb)
Borrella	one linear (0.91 Mb)	multiple circular + linear (5-200 Kb)
Bradyrhizobium japonicum	one circular (8.7 Mb)	
Brucella melitensis	two circular (2.1 + 1.2 Mb)	
Brucella suis biovars 1, 2, 4	two circular (1.0 + 2.0 Mb)	
Brucella suis biovar 3	one circular (3.1 Mb)	
Buchnera sp. strain APS	one circular (640 Kb)	two circular (< 7.8 Kb each)
Deinococcus radiodurans	two circular (2.6 + 0.4 Mb)	two circular (177 + 45 Kb)
Escherichia coli K-12	one circular (4.6 Mb )	
Leptospira interrogans	two circular (4.7 + 0.35 Mb)	
Paracoccus denitrificans	three circular (2.0 + 1.1 + 0.64 Mb)	
Pseudomonas aeruginosa	single circular (6.3 Mb)	
Rhizobacterium meliloti	two circular (3.4 + 1.7 Mb	one circular megaplasmid (1,400 Kb)
Rhodobacter sphaeroides	two circular (3.0 + 0.3 Mb)	
Ureaplasma urealyticum	one circular (0.75 Mb)	
Vibrio cholerae	two circular (2.9 + 1.1 Mb)	
Vibrio parahaemolyticus	two circular (3.2 + 1.9 Mb)	
Xylella fastidiosa	one circular (2.7 Mb)	two circular (51 + 1.3 Kb)

#### Some examples of bacterial genome organization

The first convincing evidence that some bacteria have multiple chromosomes came from studies on Rhodobacter sphaeroides. Both molecular and studies clearly demonstrated that R. sphaeroides has two large circular chromosomes. One of the chromosomes is 3.0 Mb and the other is 0.9 Mb. Genes encoding rRNAs and tRNAs required for translation, and metabolic enzymes are distributed between the two chromosomes. Multiple chromosomes have also been found in many other bacteria, including Agrobacterium tumefaciens, Rhizobium, Brucella, Paracoccus denitrificans, Ochrobactrum anthropi, Leptospira interrogans, Burkholderia, Vibrio cholerae, Deinococcus radiodurans, and many others from diverse groups of bacteria.

Furthermore, some bacteria have linear chromosomes. Borrelia have linear chromosomes and most strains contain both linear and circular plasmids; most of the bacteria in the genus Streptomyces have linear chromosomes and plasmids and some have circular plasmids as well. In addition, in some cases there may be a dynamic equilibrium between linear and circular forms

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B Sc BT	COURS	SE NAME: GENETICS		
COURSE CODE: 18BTU201	UNIT: III	BATCH-2018-2021		

of a DNA molecule. There is some evidence that linearization may be due to integration of a linear phage genome into the circular DNA molecule.

Linear chromosomes and plasmids were not discovered in bacteria until relatively recently. The first published evidence for linear chromosomes was in 1979, but because the techniques used at that time were limited and because the dogma that all bacterial chromosomes are circular was so entrenched, few people believed that linear chromosomes and plasmids occured in bacteria until 1989. By that time pulsed field gel electrophoresis had been developed, and this new technique provided convincing evidence that the chromosome of Borrelia burgdoferi was linear.

The ends of linear DNA molecules (called telomeres) pose two problems that do not apply to circular DNA molecules. First, since free double-stranded DNA ends are very sensitive to degradation by intracellular nucleases, there must be a mechanism to protect the ends. Second, the ends of linear DNA molecules must have a special mechanism for DNA replication. These problems are solved by features of the telomeres. Two different types of telomeres have been observed in bacteria: hairpin telomeres and invertron telomeres.

Hairpin telomere

			_	
Invertron telomere		•		~
	_			•

There are examples of linear DNA molecules in bacteria that are protected by both types of telomeres: palindromic hairpin loops are protected by the lack of free double-stranded ends, and invertron telomeres are protected by proteins that bind to the 5'-ends. Both of these mechanisms are also used by some phage, eukaryotic viruses, and eukaryotic plasmids.

The two types of telomeres also solve the problem of DNA replication differently. Invertron telomeres have a protein covalently attached to the 5' ends of the DNA molecule (called the 5'-terminal protein or TP for short). DNA polymerase interacts with the TP at the telomere and catalyzes the formation of a covalent bond between the TP and a dNTP. The dNTP bound to the

TP has a free 3'-OH group which acts as the primer for chain elongation. Replication of hairpin telomeres is less well understood. Apparently multiple hairpin sequences can pair to form concatemers that are replication intermediates.

The important take-home point is that we are just beginning to appreciate the similarity of many processes once thought to be completely different between bacteria and eukaryotes, partly because we now have better tools for studying these processes and partly because most of the earlier studies focused on relatively few types of bacteria. The more we study a wider diversity of bacteria, phages, and plasmids, the more obvious it becomes that E. coli is an excellent model for dissecting broad features of molecular and cell biology, but not all bacteria do everything the same way. Furthermore, we have only recently begun to attack the molecular genetics of the Archae, and what we have learned so far suggests that this diverse group of prokaryotes share even more common features with the eukaryotes.

- 1. The circular genomes of mitochondrial and chloroplast are a notable exception to the rule that eukaryotic chromosomes are linear. However, this nicely fit into the dichotomy that eukaryotic chromosomes are linear and bacterial chromosomes are circular because these organelles seem to have evolved from entrapped bacteria.
- 2. Other examples include the presence of introns, and poly-A tails on mRNA.
- 3. This genus includes B. burgdorferi, the causative agent of Lyme disease.
- 4. Streptomyces make a wide variety of useful antibiotics, including streptomycin.
- 5. For example, linear DNA was precipitated in the most commonly used procedures for purifying bacterial plasmids, and the procedures for purifying chromosomal DNA relied upon the differential binding of ethidium bromide to "sheared DNA fragments" compared to circular DNA.
- 6. It is not intuitively obvious how the ends of a linear DNA molecule could be completely replicated. All known DNA polymerases require a pre-existing primer for initiation of DNA replication. The primer is usually a short RNA molecule with a free 3'-OH group that can be extended by DNA polymerase. If a linear DNA molecule was primed at one end, DNA synthesis could continue to the other end. However, once the primer is removed, the DNA corresponding to the primer could not be replicated.

Prepared by Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE



The telomers at the end of chromosomes of most eukaryotic cells are replicated by a different mechanism: most telomeres are short GC-rich repeats that are added in a 5' to 3' direction by the enzyme telomerase.

## **Eukaryotic Chromosome Structure (Plants and Animals)**

In eukaryotes the chromosomes are found packaged within a nuclear membrane, unlike the case in prokaryotes. This membrane consists of a DNA double helix bound to an octamer of core histones (2 dimers of H2A and H2B, and an H3/H4 tetramer). Together, the DNA bound around this histone core forms what is known as the nucleosome. About 147 base pairs of DNA coil around 1 octamer, and ~20 base pairs are sequestered by the addition of the linker histone (H1), and various length of "linker" DNA (~0-100 bp) separate the nucleosomes.

Packaging of DNA is facilitated by the electrostatic charge distribution: phosphate groups cause DNA to have a negative charge, whilst the histones are positively charged. Most eukaryotic cells contain histones (with a few exceptions) as well as the kingdom Archaea, a protist group. Histones are positively charged molecules as they contain lysine and arginine in larger quantities and DNA is neatively charged. So they make a strong ionic bond in between them to form nucleosome.



### Other variations in numbers of chromosomes—Polyploidy

**Polyploidy** is a term used to describe <u>cells</u> and <u>organisms</u> containing more than two paired (homologous) sets of <u>chromosomes</u> results from having **additional whole haploid sets** of chromosomes. If a normal diploid individual is said to have **2n** chromosomes, and a normal gamete is said to have **n** chromosomes, then polyploid individuals can be represented as: 3n, 4n, 6n, 8n, etc. Polyploidy is rare in many groups of animals, but we do observe it in fish, amphibians, and lizards.

**Autotriploid**. Many domestic plants are autopolyploids. **Usually sterile**. Problems at meiosis if there are odd numbers of chromosome sets (3,5, etc.). Unbalanced gametes. Some "seedless" varieties of fruit are polyploidy.

**Allotetraploid.** Non-homologous sets of chromosomes cannot synapse at meiosis. But if both sets are doubled (**mitotic failure**), all chromosomes now have a homolog to pair with, meiosis can take place normally.

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B Sc BT	COURS	SE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: III	BATCH-2018-2021	

## **Chromosomal Mosaics**

It is when different cells within an individual, who has developed from a single fertilized egg, have a different chromosomal makeup. Most commonly there will be some cells with a typical number of chromosomes (46 chromosomes) and other cells with an altered number or structure of chromosomes. The most common kind of chromosomal mosaicism found at prenatal diagnosis involves trisomy, where the abnormal cells contain 47 chromosomes. Down syndrome mosaicism is an example of trisomy mosaicism. These individuals have some cells with the typical number of chromosomes (46) and some cells with an extra chromosome 21, for a total of 47 chromosomes. Mosaicism may exist for all kinds of chromosome abnormalities (monosomy, triploidy, structural changes, etc). Although more rare, there may even be mosaicism where both different cell types are abnormal in structure or number, and there are no normal cells involved.

Chromosomal mosaicism describes a **group of disorders**. Any chromosomal change, in number or structure, can be present in a mosaic form. However, trisomy mosaicism is the most common kind ascertained during prenatal diagnosis.

All of the cells in our body come from a single cell, the fertilized egg or zygote. In order for the zygote to develop into a baby, this single cell must grow and divide. Before cells can divide each chromosome must make an identical copy of itself. At cell division each chromosome and its identical copy pull apart into two separate cells. Now the resulting cells, also called <u>daughter cells</u>, have the same chromosome make-up as the original cell. The two new cells will repeat this process. In each cell, the chromosomes will duplicate and divide into two new cells. The result, is 4 cells identical to the first original cell. This process of cell division is called <u>mitosis</u>. Sometimes a mistake can occur when the chromosomes are separating into the two daughter cells. An extra chromosome may travel into the wrong cell or a chromosome may get lost in the separation of the cells. The result would be two daughter cells with different chromosomal make-up.



Prepared by Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE

In this illustration, the green cell represents a cell with an abnormal chromosome make-up. All cells that come from the green cell will share the same chromosome change. We say that all cells originating from that cell are in the same <u>cell line</u>. The baby that develops from this embryo will have some cells in his/her body which have the typical number of chromosomes and some that have the chromosome change.



Trisomy mosaicism can occur in one of two ways:

• In an abnormal fertilized egg with 47 chromosomes, one of the cells may lose the extra chromosome at cell division, leaving 46 chromosomes in that cell. All cells that are derived from that cell will have 46 chromosomes. The rest of the cells will have 47 chromosomes.

In a typical zygote with 46 chromosomes, at cell division one of the cells may retain a duplicated copy of one of the chromosomes. This produces a cell with 47 chromosomes. All cells that are

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B Sc BT	COURS	SE NAME: GENETICS		
COURSE CODE: 18BTU201	UNIT: III	BATCH-2018-2021		

derived from that cell also have 47 chromosomes. The rest of the cells will have 46 chromosomes.

## **Polytene Chromosome:**

To increase cell volume, some specialized cells undergo repeated rounds of DNA replication without cell division (endomitosis), forming a giant **polytene chromosome**. Polytene chromosomes form when multiple rounds of replication produce many sister chromatids that remain synapsed together.

In addition to increasing the volume of the cells' nuclei and causing cell expansion, polytene cells may also have a metabolic advantage as multiple copies of genes permits a high level of gene expression. In *Drosophila melanogaster*, for example, the chromosomes of the larval salivary glands undergo many rounds of endoreplication, to produce large amounts of glue before pupation.

Polytene chromosomes have characteristic light and dark banding patterns that can be used to identify chromosomal rearrangements and deletions. Dark banding frequently corresponds to inactive chromatin, whereas light banding is usually found at areas with higher transcriptional activity. The banding patterns of the polytene chromosomes of *Drosophila melanogaster* were sketched in 1935 by Calvin B. Bridges, in such detail that his maps are still widely used today. The banding patterns of the chromosomes are especially helpful in research, as they provide an excellent visualization of transcriptionally active chromatin and general chromatin structure.



**Chromosome puffs** are diffused uncoiled regions of the polytene chromosome that are sites of RNA transcription. A **Balbiani ring** is a large chromosome puff.



Polytene chromosomes were originally observed in the larval salivary glands of *Chironomus* midges by Balbiani in 1881, but the hereditary nature of these structures was not confirmed until they were studied in *Drosophila melanogaster* in the early 1930s by Emil Heitz and Hans Bauer. They are known to occur in secretory tissues of other dipteran insects such as the Malpighian tubules of *Sciara* and also in protists, plants, mammals, or in cells from other insects. Some of the largest polytene chromosomes described thus far (see scale bar in figure below) occur in

larval salivary gland cells of the Chironomid genus *Axarus*. Polytene chromosomes are about 200micron in length. The chromonema of these chromosomes divide but do not separate. Therefore, they remain together to become large in size. Another form of chromosomal enlargement that provides for increased transcription is the lampbrush chromosome.

Polytene chromosomes are also used to identify the species of Chironomid larvae that are notoriously difficult to identify. Each morphologically distinct group of larvae consists of a number of morphologically identical (sibling) species that can only be identified by rearing adult males or by cytogenetic analysis of the polytene chromosomes of the larvae. Karyotypes are used to confirm the presence of specific species and to study genetic diversity in species with a wide range.

## **GENETIC code**

As we learned in Structure of Nucleic Acids, DNA and RNA are made up by sequences of nitrogen bases-pairs: adenine, thymine, guanine, and cytosine. Scientists have long understood that these nitrogen bases somehow contained the information that coded for specific amino acids. However, it took some time before they figured out how the base pairs accomplished this coding. Scientists main problem lay in the fact that while there were only 4 nitrogen bases (nucleotides), there were 20 amino acids for which those nucleotides had to code. If adenine, thymine, guanine, and cytosine each coded for a particular amino acid, then the DNA/mRNA information system would only be able to code for 4 amino acids. If, however, groups of two nucleotides coded for a single amino acid, the story is somewhat different. Given four nucleotides looked at in groups of two, there are sixteen possible combinations (AA, AT, AG, AC, TA, TT, TG, TC, GA, GT, GG, GC, CA, CT, CG, CC); that sixteen is still not enough to code for twenty amino acids. But if the nucleotides code for amino acids in groups of three then there are sixty-four possible combinations. Scientific experiments have verified that nucleotides code for amino acids in successive groups of threes. These groups of threes are called codons.

## **Degeneracy of the Genetic Code**

As we know, since the genetic code is read in triplets and there are four possible bases that can occupy each position, the number of possible codons is 4 X 4 X 4, or 64 codons. However, there are only 20 known amino acids. Experiments have shown that three codons function also function stop codons, acting as termination signals in translation. Yet that brings the count up to only twenty-three necessary codons. The vast difference between possible codon variations and needed codon variations means, as seen in the figure below, that each amino acid is specified by more than one codon. Because the genetic code therefore does not code to its capacity, it is called "degenerate".

		U	С	А	G		
First position (5' end)	U	UUU UUC UUA UUG	UCU UCC UCA UCG	UAU UAC UAA Stop UAG Stop	UGU Cys UGC Cys UGA Stop UGG Trp	U C A G	
	с	CUU CUC CUA CUG	CCU CCC CCA CCG	CAU CACHis CAA CAGGIn	CGU CGC CGA CGG	U C A G	Third position
	А	AUU AUC AUA AUG	ACU ACC ACA ACG	AAU AAC AAA AAA AAG	AGU AGCSer AGA AGGArg	U C A G	(3 end)
	G	GUU GUC GUA GUG	GCU GCC Ala GCA GCG	GAU GAC GAA GAG GIU	GGU GGC GGA GGG_	U C A G	

#### Amino acid names:

Ala = alanine	GIn = glutamine	Leu = leucine	Ser = serine
Arg = arginine	Glu = glutamate	Lys = lysine	Thr = threonine
Asn = asparagine	Gly = glycine	Met = methionine	Trp = tryptophan
Asp = aspartate	His = histidine	Phe = phenylalanine	Tyr = Tyrosine
Cys = cysteine	Ile = Isolevcine	Pro = proline	Val = valine

Prepared by Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE

## KARPAGAM ACADEMY OF HIGHER EDUCATION

## CLASS: I B Sc BT COURSE CODE: 18BTU201

COURSE NAME: GENETICS UNIT: IV BATCH-2018-2021

## UNIT-IV

## **SYLLABUS**

**GENETICS:** Definition and types of mutation, Causes of mutations, Ames test for mutagenic agents, Screening procedure for isolations of mutants and use of mutants, Variations in chromosomes structure deletion, duplication, Inversion and translocation, Sex determination, Sex linkage, sex linked diseases

## **Gene Mutation**

Mutation = a process that produces a gene or chromosome that differs from the wild type 2. Mutation = the gene or chromosome that results from a mutational process 3. a mutant is the organism or cell whose changed phenotype is attributed to a mutation B. General Types 1. Gene mutation = the allele of a gene changes. 2. Chromosome mutation = segments of chromosomes, whole chromosomes, or entire sets of chromosomes change C. What does wild type (wt) mean? Wild type is an arbitrary standard for what "normal" is for an organism. Please remember that what is considered wild type today may have been a mutant in the evolutionary past. D. Direction of the mutation 1. Forward mutations are changes away from the wt 2. Reverse mutations (reversions) are changes from the mutant allele back to the wt allele E. Mechanisms for gene mutation 1. Errors in DNA replication 2. Errors in DNA repair 3. Environmental mutagen causes DNA damage that is not repaired correctly 4. Transposons and insertion sequences (a mobile DNA elements that can move from one location in the chromosome to another; the element may "jump" into a gene thereby mutating it) F. Why study gene mutation? 1. Variants in genes (which are caused by mutations) are needed to study the transmission of traits 2. Mutations can tell the researcher about the function of a gene product in a biological system 3. Mutations are the basis for cancer and other genetic diseases 4. Gene mutations serve as the source for most alleles in a population and is therefore the origin of genetic variation within a population 5. Mutations drive evolution: mutations are the raw material upon which natural selection acts. II. Classification of mutations A. General info 1. Various schemes for classification depending upon which aspect of mutation is being examined 2. Classes are not mutually exclusive B. Point of origin 1. Somatic mutations a) mutations that are in the somatic tissues of the body b) mutations are NOT transmitted to progeny c) the extent of the phenotypic

Prepared by Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE

effect depends upon whether the mutation is dominant or recessive (dominant mutations generally have a greater effect) d) the extent of the phenotypic effect depends upon whether it occurs early or late in development (early arising mutations have a greater effect) (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) e) sectoring phenotypes may be seen when the mutation occurs during embryonic development f) cancer caused by somatic mutations 2. Germinal mutations a) mutations that are in the germ tissues of the body b) mutations MAY BE transmitted to progeny c) dominant mutations are seen in first generation after the mutation occurs d) if a female gamete containing an X-linked mutation is fertilized, the males will show the mutant phenotype e) recessive mutations will only be seen upon the chance mating with an individual carrying the recessive allele too; thus, the recessive mutation may remain hidden for many generations C. Phenotypic effects 1. Morphological mutations are mutations that affect the outwardly visible properties of an organism (i.e. curly ears in cats) 2. Lethal mutations are mutations that affect the viability of the organism (i.e. Manx cat). 3. Conditional mutations are mutations in which the mutant allele causes the mutant phenotype only in certain environments (called the restrictive condition). In the permissive condition, the phenotype is no longer mutant. (i.e. Siamese cat - mutant allele causes albino phenotype at the restrictive temperature of most of the cat body but not at the permissive temperature in the extremities where the body temperatures is lower). 4. Biochemical mutations are mutations that may not be visible or affect a specific morphological characteristic but may have a general affect on the ability to grow or proliferate. a) Most microorganisms are prototrophs which means that they can grow on a simple growth medium including an energy source and inorganic salts. Biochemical mutations include those that affect proteins or enzymes required to grow on various nutrients or to synthesize various components. Thus, these mutations cause the microorganisms to become auxotrophs (they must be supplied with additional nutrients if they are to grow). For example, the bacterium Escherichia coli does NOT require the amino acid tryptophan for growth because they can synthesize tryptophan. However, there are E. coli mutants that have mutations in the trp genes. These mutants are auxotrophic for tryptophan, and tryptophan must be added to the nutrient medium for growth. b) Humans can also have biochemical mutations (also called inborn errors in metabolism). Such examples include hemophilia, phenylketonuria, and galactosemia. D. Loss of function vs. gain of function

mutations 1. Loss of function mutations are those that destroy the function of the gene product. Many times in diploid organisms, these are recessive mutations because the other wild type allele still encodes a functional gene product. However, it is possible to have a dominant loss of function mutation in which the mutant gene product interferes with the activity of the gene product from the wild type allele. a) Null mutation = loss of function mutation where gene product is completely inactive (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) b) Leaky mutation = loss of function mutation where gene product is not completely inactive (partially active still) (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) 2. Gain of function mutations are those that produce a new function for the gene product. Gain of function mutations are dominant. (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) III. The occurrence of mutations A. Frequencies of mutations 1. Mutation frequency = # of times mutation appears in the population / # of individuals in the population where a population can be bacterial cells, people, gametes 2. Mutation rate = # of mutations / unit time where unit time can be per cell division, cell generation (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) 3. Mutations are relatively rare. 4. Different genes have different mutation frequencies (Table 7-1) 5. Different organisms have different overall mutation frequencies (Table 7-2) B. Detection of mutations in humans 1. Detection of germinal dominant mutations by human pedigree analysis (shows up in the pedigree as the sudden appearance of a novel phenotype) 2. Detection of germinal recessive mutations are more difficult because they remain masked by the dominant allele until the union of two heterozygotes 3. Detection of germinal X-linked mutations arising in female gametes appear in some of the males in the generation after the mutation occurred. C. Detection of mutations using the specific-locus test, a system for detecting recessive mutations in diploids. Heterozygote individual for gene(s)A that give phenotype A is crossed with a homozygous recessive individual for gene(s) a that gives phenotype a. The frequency of the mutant phenotype (a) is quantitated. (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) D. Detection of X-linked mutations in Drosophila using the ClB chromosome The ClB chromosome is the X chromosome bearing the C allele which prevents crossover, the l allele which is a recessive lethal, and the Bar allele which is a dominant eve
mutation. (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) E. Detection of mutations in microorganisms 1. General info on microorganism growth and manipulation a) Growth is rapid (some bacterial populations double every 30 minutes) b) Growth can occur in liquid media to cell density of approx 109 bacteria/ml c) Growth can occur on solid media (agar plate). Organisms are spread across a plate of solid media (plating). Single cells are deposited randomly and will grow into a colony of cells that is visible to the naked eye. 2. Microorganisms allow for the use of selective systems for mutation detection vs. the screening systems used for higher organisms. A selective system is one in which the experimenter can DEMAND that the only individuals that grow or survive are the ones that have the mutation of interest. On the other hand, a screening system is one in which the experimenter must examine each individual to see if it has the mutation of interest. 3. Microbial selective systems a) Selection for reversion of an auxotroph to a prototroph: Plate 109 adenine auxotrophs on agar plate with no adenine. The only bacteria that grow are those that have a random mutation in the mutant ad gene that now reverts it back to the wild type allele. (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) b) Selection for resistance to an environmental factor (bacteriophage, antibiotics). Example, Luria-Delbruck experiment: Tested whether mutations arose in response to plating on selective media (in this case bacteriophage resistant mutants were selected for by plating on media containing bacteriophage which normally kill the bacterium). Two hypotheses: (a) physiological change where mutations arose after plating because the bacteria sensed the phage and altered themselves so that they may become resistant to the phage or (b) random mutation where mutations arose randomly before plating and all plating did was select for the resistant bacteria (from An Introduction to Genetic Analysis, 6th ed. By Griffiths et al. W. H. Freeman and Company) Versus Luria and Delbruck tested the above hypothesis by starting 20 very small cultures (0.2 ml) and 1 large culture (10 ml) with low numbers of bacteria. The cultures were allowed to grow for numerous generations and them 0.2 ml aliquots were plated on phage plates (all of the small cultures and aliquots of the large culture) and the # of resistant bacteria were counted. If the physiological hypothesis was correct, then one would expect that each plate would contain the same number of phage resistant bacteria. However, if the random mutation hypothesis was correct, some of the small cultures may have had the random phage resistance mutation occur

early and thus almost the entire population is phage resistant while other small cultures may have had the random phage resistance mutation occur very late early and thus almost the entire population is not phage resistant. They found a large fluctuation in the number of phage resistant bacteria on plates from the small 0.2 ml cultures (Table 7-3), and thus random mutation was the correct hypothesis. In the larger culture, the fluctuations are averaged out and so you do not see the large variation in #s. 4. Microbial screening systems a) Screening for forward mutations from wild type to auxotrophy: Plate bacteria on complete media to form colonies. Replica plate many, many colonies to plates with and without the nutrient you are testing for auxotrophy. Compare the plus and minus nutrient X plates to look for a colony that appears on the plus nutrient plate but not the minus nutrient plate. If you expect the mutation frequency to be 1 in 10,000, then you will need to replica plate at least 10,000 colonies to get the 1 mutant. b) Enrichments are important in screening because they reduce the number of organisms that you have to screen. Enrichment for auxotrophs works by selectively removing or killing growing microorganisms while they are in medium that allows only the prototrophs to grow. (1) Filtration enrichment (Figure 7-18) (2) Penicillin enrichment IV. Mutations and cancer A. Cancer is a group of diseases characterized by rapid, uncontrolled proliferation of cells within a tissue resulting in the formation of a tumor. Cancer has many causes and phenotypes but the fundamental mechanism underlying all cancers is genetic. B. There are two types of genes that are involved in cancer formation. 1. Tumor suppressor genes are genes that encode a product that normally stops cell division. Mutations in these genes result in uncontrolled activation of cell division and therefore tumor formation. Mutations are generally recessive and thus you need mutations in both alleles to have cancer. A mutation in one allele predisposes the carrier to cancer. a) Rb gene retinoblastoma (retinal cancer) b) BRCA1 - hereditary breast cancer gene c) p53 gene mutations are found in a variety of cancers including breast, lung, bladder, and colon cancers. Over 1/2 of all cancers are associated with p53. 2. Proto-oncogenes are genes that encode a product that normally controls cell division (kind of like an on/off switch). Mutations in these genes make the gene product permanently in the on position which results in uncontrolled activation of cell division and therefore tumor formation. a) N-ras – neuroblastoma (tumor formed of embryonic ganglion cells), leukemia b) N-myc – neuroblastoma c) man – mammary carcinoma V. Mutagens in genetic dissection A. Mutagens are agents that cause mutations at a rate higher than the

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B Sc BT	COURS	SE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: IV	BATCH-2018-2021	

spontaneous rate. B. Mutagens can be chemical (i.e. cigarette smoke, mustard gas) or radiation (i.e. UV, X-rays, )

#### CHOMOSOME MUTATION:

#### CHANGES IN CHROMOSOME STRUCTURE)

Chromosomal mutations are processes that result in rearranged chromosome parts, abnormal numbers of individual chromosomes, or abnormal numbers of chromosome sets. The resulting products are also known as chromosomal mutations. B. For our purposes here, we will be talking about alterations in large regions of the chromosome spanning numerous genes C. Abnormalities from chromosomal mutations are frequently due to: 1. change in gene number (balance) 2. change in gene location 3. break internal to a gene D. Can occur in somatic cells, germinal cells, and gametes E. How can you detect a chromosomal mutation? cytogenetics 1. Cytological examination of the actual chromosomes 2. Genetic analysis of inheritance II. Physical features of the chromosomes A. Size B. Centromere position (appears constricted under microscope) 1. Telocentric (at one end of the chromosome) 2. Acrocentric (off center on the chromosome) 3. Metacentric (in the center of the chromosome) B. Arm length: centromere to the end defines an arm (p =short arm and q =long arm) C. Nucleolar organizer position 1. Nucleoli = intranuclear organelles that contain rRNA (appears as dark spot under microscope) 2. Number of nucleoli ranges from 1 to many per chromosome set 3. Position of nucleolus is next to constriction of the chromosomes called nucleolar organizers which themselves have specific chromosome locations D. Chromomere patterns (beadlike, localized thickening along the chromosomes during prophase) E. Heterochromatin patterns 1. Constitutive heterochromatin is highly condensed chromosomal regions that are for the most part genetically inert and are found in particular chromosomal regions 2. Detected by staining with chemicals (such as Feulgen) that react with DNA as very dense regions of staining F. Banding patterns due to special chromosome staining procedures 1. Specific stains reveal specific bands at specific chromosomal sites a) Quianacrine hydrochloride ‡ Q bands b) Giemsa stain ‡ G bands c) Reversed giemsa ‡ R bands 2. Specialized case of polytene chromosomes in Drosophila E. Karotype = the entire metaphase chromosome complement of individual F. Physical characteristics of chromosomes are useful for detecting chromosome mutations such as chromosome rearrangements including deletions, duplications,

#### KARPAGAM ACADEMY OF HIGHER EDUCATION

#### CLASS: I B Sc BT COURSE CODE: 18BTU201

#### COURSE NAME: GENETICS UNIT: IV BATCH-2018-2021

inversions, and translocations. III. Deletions (loss of a chromosomal segment) A B C o D E F ‡ A C o D E F A. Types 1. Interstitial (Not at the end of the chromosome) 2. Terminal (Near the end of the chromosome) B. Deletions require two breaks in the chromosome followed by loss of the chromosomal segment and rejoining of the ends. C. The deleted fragment is acentric (without a centromere) and will be lost upon multiple rounds of cell division. D. In meiosis I cells heterozygous for the deletion, chromosomes line up normally except that region corresponding to deletion forms a "loop formed to get proper alignment of the homologous chromosomes. E. Effect of multigenic deletions 1. Homozygous for multigenic deletion = usually lethal. 2. Heterozygous for mutigenic deletion = frequently lethal because a) Unmasking of lethal allele on homologous chromosome without the deletion b) Upset balance of gene number F. Detection 1. Genetic a) Failure of homozygotes to survive b) Inability of the mutation to revert back to wild type c) Recombination frequency between the genes flanking the deletion is lower than in the wild type d) Unmasking of a recessive allele present on the homologous chromosome without the deletion (pseudodominance of the recessive allele) 2. Cytological a) Deletion loop in meiosis b) Change in banding patterns G. Using deletions to map genes 1. Deletion mapping a) Cross a new recessive mutant with a set of deletion mutants of known map location. The new mutation will show pseudodominance when heterozygous for the deletion that contains the region to which the new mutation maps. See figure 8-8 in text. b) Cross a new mutant suspected of containing a deletion with a set of recessive mutants of known map location. Look for the pseudodominance effect which indicated that your putative deletion is in the same area as the gene that is showing pseudodominance 2. In situ hybridization Add a radioactively labeled gene of interest to a chromosome deletion set and look for the presence of radioactivity for each deletion. The chromosome deletions that do no hybridize to the radioactive gene probe are deleted for the region that contains the gene of interest. IV. Duplications (presence of two copies of a chromosomal region) ABC 0 DEF 1 ABCBC 0 DEF OR A B C O D E F ‡ A B C C B O D E F A. Types 1. tandem (adjacent to each other in same order) 2. reverse tandem (adjacent to each other in reverse order) 3. nontandem (at a different chromosomal location) B. Tandem duplications can occur due to unequal crossing over where homologous chromosomes pair inaccurately during meiosis I

(Deletion) (Duplication) C. Nontandem duplications may result from crossing over during meiosis within segments of the chromosome that contain inversions or translocations (see later). D. In meiosis I cells heterozygous for the duplication, chromosomes line up normally except that region corresponding to duplication forms a "loop" formed to get proper alignment of the homologous chromosomes. Also, asymmetric pairing within the duplicated region may occur during meiosis I, leading to unequal crossover. (from An Introduction to Genetic Analysis, 6th ed. by Griffiths et al. (W. H. Freeman and Company) E. Effect of duplications 1. Duplications are rare. 2. Heterozygous for duplication = can be lethal because of imbalance generated by extra copies of the duplicated region. 3. Homozygous for duplications is unknown in medical genetics. F. Detection is difficult. Cytology is the main tool (change in bands and duplication loop in meiosis I) G. Duplications have been proposed to be important for evolution because they provide an extra copy of genes that can then undergo mutation while still leaving the original gene unaffected. V. Inversions (flipping of chromosomal segment relative to the rest of the chromosome) A B C D E o F G H ‡ A D C B E o F G H OR ABC O DEF ‡ AED O CBF A. Types 1. Paracentric – centromere outside the inversion 2. Pericentric - centromere inside the inversion B. Inversions require two breaks in the chromosome followed by a 180° rotation of the chromosomal segment and rejoining of the ends. C. In meiosis I cells heterozygous for the inversion, inversion loop is formed to get proper alignment of the homologous chromosomes. 1. Paracentric (from An Introduction to Genetic Analysis, 6th ed. by Griffiths et al. (W. H. Freeman and Company) 2. Pericentric (from An Introduction to Genetic Analysis, 6th ed. by Griffiths et al. (W. H. Freeman and Company) D. In meiosis I cells homozygous for the inversion, crossover is normal but linkage map shows inverted gene order. E. Effects of inversions are negligible because there is no change in the genetic material. However, those heterozygous for inversions produce abnormal meiotic crossover products and possibly abnormal progeny (if they survive at all). F. Detection 1. Genetic – recombinant progeny are usually not recovered from heterozygotes 2. Cytological a) If the duplication is pericentric, the arm lengths change. b) Change in the banding pattern c) Dicentric bridge in meiosis for paracentric inversion IV. Translocations (movement of a chromosomal segment from one location to another) \_A\_B\_C\_\_\_o\_D\_E\_\_ and  $\_G_H_I_o_J_K_$   $\ddagger$   $\_A\_B\_C\_o\_J\_K\_$  and  $\_G_H_I\_o\_D\_E\_$  A. Types 1.

Reciprocal (exchange of segments) 2. Nonreciprocal (1 segment moves to a new location without an exchange) B. Reciprocal translocations require two breaks in two different chromosomes followed by rejoining of the ends. C. In meiosis I cells heterozygous for the deletion, cross conformation forms to get proper alignment of the homologous chromosomes. Showing only meiosis I on the next page: A B C D E G H I J K A B C J K G H I D E A B C D E I H G D E A B C J K J K I H G Lining up during Meiosis I Alternate Segregation A B C D E G H I J K A B C J K G H I D E Adjacent-1 Segregation A B C D E G H I J K A B C J K G H I D E wild type translocated Deleted for JK Duplicated for DE Deleted for DE Duplicated for JK OR Heterozygous for translocation (In most cases, these gametes form nonviable progeny even when mated with wild type.) D. Effect of translocations 1. Semisterility due to adjacent segregation in meiosis 2. Position effects: altered expression of a gene when it is moved to a new location E. Detection 1. Genetic a) Semisterility b) Apparent linkage of genes on separate chromosomes c) Position effects 2. Cytological a) Can change the location of the centromere b) Change in the size of the chromosome c) Cross formation in meiosis I

#### VARIATION IN CHROMOSOME NUMBER & STRUCTURE

Changes in chromosome number can occur by the addition of all or part of a chromosome (**aneuploidy**), the loss of an entire set of chromosomes (**monoploidy**) or the gain of one or more complete sets of chromosomes (**euploidy**). Each of these conditions is a variation on the normal diploid number of chromosomes. As you would expect each of these can have drastic effects on phenotypic expression.

**Aneuploidy** - the abnormal condition were one or more chromosomes of a normal set of chromosomes are missing or present in more than their usual number of copies

Monoploidy - the loss of an entire set of chromosomes

Euploidy - an entire set of chromosomes is duplicated once or several times

**Chromosome Number in Different Species** 

In "higher" organisms (**diploids**), members of same species typically have identical numbers of chromosomes in each somatic cell. **Diploid chromosome number** (2n). Nearly all chromosomes will exist in pairs (identical wrt length and centromere placement) except the sex chromosomes. Members of pair are **homologous chromosomes**. **Haploid number** (n) is the number of chromosome pairs.

	<u>2n</u>	<u>n</u>
Human	46	23
Horse 64	32	
Cat	38	19
Geometrid moth	224	112
Tomato	24	12
Pink bread mold	14	7

#### Autosomal monosomy and trisomy

Occasionally, one finds an organism that has an extra copy of a particular chromosome. This is known as **trisomy**-because there are now 3 copies of an autosome. Some trisomies are viable in animals, but the condition usually has severe effects. These effects are presumably related to the fact that there are 3 copies of every gene on the trisomic chromosome, but only 2 copies of all the genes on the other chromosomes. We will see later, that organisms with three or more copies of **all** the chromosomes are often perfectly viable.

Trisomy of human chromosome 21 is the cause of the disorder known as **Down syndrome.** 

(Remember, humans have 23 pairs of chromosomes--the pairs are numbered 1 through 22, plus the X and Y). It is characterized by multiple physical defects, including epicanthal fold, furrowed tongue, characteristic palm and finger print patterns, and lowered IQ. About 1 in 750 live births produces a child with this condition. It results from the **non-disjunction** of chromosome 21

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B Sc BT	COURS	SE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: IV	BATCH-2018-2021	

during meiotic anaphase I or anaphase II, when the paired homologs (or paired chromatids) normally migrate to opposite poles of the cell.

**Nondisjuction:** Nondisjunction ("not coming apart") is the failure of chromosome pairs to separate properly during meiosis stage 1 or stage 2. This could arise from a failure of homologous chromosomes to separate in meiosis I, or the failure of sister chromatids to separate during meiosis II or mitosis. The result of this error is a cell with an imbalance of chromosomes. Such a cell is said to be an uploid. Loss of a single chromosome (2n-1), in which the daughter cell(s) with the defect will have one chromosome missing from one of its pairs, is referred to as a monosomy. Gaining a single chromosome, in which the daughter cell(s) with the defect will have one chromosome in addition to its pairs is referred to as a trisomy. The members of a chromosome pair (homologs) line up at the metaphase plate during meiotic metaphase I, then separate to opposite poles of the cell during anaphase I--review this material in Klug and Cummings or any introductory Genetics text if you are not thoroughly familiar with it!). If the pair fails to separate, and both migrate to the same pole, half of the resulting gametes will have two copies of chromosome 21, rather than one. When this gamete unites with a normal gamete (bearing one copy of chromosome 21) during fertilization, the resulting gamete has 3 copies of chromosome 21, rather than the normal 2. Nondisjunction of chromosome 21 seems to occur more often in the production of eggs than sperm, and the frequency increases with the age of the parent. Older individuals are often encouraged to test for trisomy 21 by amniocentesis at 15 to 16 weeks after conception. Nondisjunction can happen to other chromosomes in addition to chromosome 21. But human embryos that are trisomic for any other chromosome do not survive to birth. It should be obvious that the other half of the gametes resulting from a non-disjunction event at anaphase 1 will have 0 copies of the chromosome. When a gamete with 0 copies of a chromosome unites with a normal gamete, the result is a zygote that has only one copy of that chromosome. This is monosomy. Monosomy is not well tolerated in animals—usually lethal. Some plants can survive (observed in maize, tomato, Oenothera, and Datura) but they have low viability and are usually sterile. Nondisjunction can also occur at anaphase II, when sister chromatids fail to separate and migrate to opposite poles. Nondisjuntion at anaphase one results in half the gametes being normal, and half being abnormal (see diagram in text).

If one surveys karyotypes of embryos that spontaneously abort, trisomies for all the autosomes are seen, and other forms of chromosomal abnormalities as well, but these conditions are apparently fatal early in development. Normal embryonic development requires a precise diploid complement of chromosomes.

2n	2n
/ \	/ \
n+1 n-1	n n
n+1 n+1 n-1 n-1	n n n+1 n-1
Schematic of nondisjunction in meiosis I.	Schematic of nondisjunction in meiosis II.
Duplicated chromosomes in diploid cell (2n).	Duplicated chromosomes in diploid cell (2n).
All gametes are affected by nondisjunction in	Half of the gametes are affected by
meiosis I. Two gametes have a single extra	nondisjunction in meiosis II. One gamete has a
chromosome; two gametes are missing a single	single extra chromosome; one gamete is missing
chromosome.	a single chromosome.

#### **Chromosomal Rearrangements:**

Variations in chromosome structure also exist, and also cause characteristic genetic effects. We will consider deletions and duplications of parts of whole chromosomes, and inversions and translocations of sections of chromosomes.

**Deletion:** Sometimes a chromosome will arise in which a segment is missing. These chromosomes are said to have **deletions.** Deletions are generally harmful, and typically, the larger the deletion, the more harmful it is. Small deletions are often viable if the deletion is heterozygous, because the other chromosome contains copies of the genes missing in the chromosome with the deletion. But even small deletions are usually lethal if they are

homozygous. A deletion on one homologue can "unmask" recessive alleles on the other homologue--this effect is called pseudo-dominace.



#### **Duplications**

Duplication often results in reduction of viability, but in general duplications are less severe in their effects than are deletions.



Heterozygotes and homozygotes for small duplications can be viable, although often exhibit phenotypic effects (Bar eye in *Drosophila*).

Duplication may be important in generating multiple copies of genes that require very high levels of expression (**gene amplification**)--**rDNA** codes for genes producing RNA for construction of ribosomes. Also may be source of evolutionary origin and divergence of genes coding for similar (but not identical) proteins (**alpha and beta globin components of hemoglobin**).

#### Inversions

Chromosomes can sometimes break and rejoin in a different orientation. If this happens in a germ cell, the gametes will contain a **rearrangement or inversion**. The area between the breaks is inverted (turned around), and then reinserted and the breaks then unite to the rest of the chromosome. If the inverted area includes the centromere it is called a **pericentric inversion**. If it does not, it is called a **paracentric inversion**.



#### Translocations

5% of people with Down Syndrome have one parent who is heterozygous for a translocation. Chromosome 14 is translocated onto chromosome 21. Half the time, meiosis produces a normal set of balanced translocated chromosomes. But half the time, unbalanced chromosomes are produced, either a 14 w/out the translocated 21 segment or a translocated 14 with the attached 21 plus a normal 21.



#### Chromosome rearrangements in evolution

Chromosomal rearrangements are not just rare abnormalities. Many kinds of rearrangements have no deleterious effects, and some may be beneficial (e.g., duplications are the source of novel gene functions and allow evolutionary increase in gene number). Comparisons among species show that chromosomal rearrangements have occurred repeatedly throughout evolutionary history.

Humans normally have 46 chromosomes in each cell, divided into 23 pairs. Two copies of chromosome 8, one copy inherited from each parent, form one of the pairs. Chromosome 8 spans more than 146 million DNA building blocks (base pairs) and represents between 4.5 and 5 percent of the total DNA in cells.

Identifying genes on each chromosome is an active area of genetic research. Because researchers use different approaches to predict the number of genes on each chromosome, the estimated number of genes varies. Chromosome 8 likely contains about 700 genes that provide instructions for making proteins. These proteins perform a variety of different roles in the body.

Geneticists use diagrams called ideograms as a standard representation for chromosomes. Ideograms show a chromosome's relative size and its banding pattern. A banding pattern is the characteristic pattern of dark and light bands that appears when a chromosome is stained with a



#### **Definition(s)**

Absence of a segment of DNA; may be as small as a single base or large enough to encompass one or more entire genes. Large deletions involving a whole segment of a chromosome may be detected by routine examination of the chromosomes; intermediate deletions involving a few genes may be detected by using fluorescent in situ hybridization (FISH); smaller deletions involving a portion of a gene may only be detected by analyzing the DNA.

A loss of part of the DNA from a chromosome; can lead to a disease or abnormality.

Deletion is a type of mutation involving the loss of genetic material. It can be small, involving a single missing DNA base pair, or large, involving a piece of a chromosome.

A deletion occurs when a chromosome breaks and some genetic material is lost.

# CLASS: I B Sc BT COURSE NAME: GENETICS COURSE CODE: 18BTU201 UNIT: IV BATCH-2018-2021

#### Duplication

#### **Definition(s)**

The presence of an extra segment of DNA, resulting in redundant copies of a portion of a gene, an entire gene, or a series of genes, usually caused by unequal crossing-over during gene replication when gametes are formed in meiosis

Duplication is a type of mutation that involves the production of one or more copies of a gene or region of a chromosome. Gene and chromosome duplications occur in all organisms, though they are especially prominent among plants. Gene duplication is an important mechanism by which evolution occurs.

A section of DNA is accidentally duplicated when a chromosome is copied.



#### Inversion

#### **Definition**(s)

A chromosomal rearrangement in which a segment of genetic material is broken away from the chromosome, inverted from end to end, and re-inserted into the chromosome at the same breakage site. Balanced inversions (no net loss or gain of genetic material) are usually not associated with phenotypic abnormalities, although in some cases gene disruptions at the breakpoints can cause adverse phenotypic effects, including some known genetic diseases; unbalanced inversions (loss or gain of chromosome material) nearly always yield an abnormal phenotype.

Inversions occur when a chromosome breaks in two places and the resulting piece of DNA is reversed and re-inserted into the chromosome. Inversions that involve the centromere are called pericentric inversions; those that do not involve the centromere are called paracentric inversions.



#### Translocation

#### **Definition(s)**

A chromosome alteration in which a whole chromosome or segment of a chromosome becomes attached to or interchanged with another whole chromosome or segment, the resulting hybrid segregating together at meiosis; balanced translocations (in which there is no net loss or gain of chromosome material) are usually not associated with phenotypic abnormalities, although gene disruptions at the breakpoints of the translocation can, in some cases, cause adverse effects,

including some known genetic disorders; unbalanced translocations (in which there is loss or gain of chromosome material) nearly always yield an abnormal phenotype

#### **Balanced translocation**

In a balanced translocation, pieces of chromosomes are rearranged but no genetic material is gained or lost in the cell.



#### **Unbalanced translocation**

An unbalanced translocation occurs when a child inherits a chromosome with extra or missing genetic material from a parent with a balanced translocation.



#### Can changes in the structure of chromosomes affect health and development?

Changes that affect the structure of chromosomes can cause problems with growth, development, and function of the body's systems. These changes can affect many genes along the chromosome and disrupt the proteins made from those genes.

Structural changes can occur during the formation of egg or sperm cells, in early fetal development, or in any cell after birth. Pieces of DNA can be rearranged within one chromosome or transferred between two or more chromosomes. The effects of structural changes depend on their size and location, and whether any genetic material is gained or lost. Some changes cause medical problems, while others may have no effect on a person's health.

#### Changes in chromosome structure include:

#### Translocations

A translocation occurs when a piece of one chromosome breaks off and attaches to another chromosome. This type of rearrangement is described as balanced if no genetic material is gained or lost in the cell. If there is a gain or loss of genetic material, the translocation is described as unbalanced.

#### Deletions

Deletions occur when a chromosome breaks and some genetic material is lost. Deletions can be large or small, and can occur anywhere along a chromosome.

#### Duplications

Duplications occur when part of a chromosome is copied (duplicated) too many times. This type of chromosomal change results in extra copies of genetic material from the duplicated segment.

#### Inversions

An inversion involves the breakage of a chromosome in two places; the resulting piece of DNA is reversed and re-inserted into the chromosome. Genetic material may or may not be lost as a result of the chromosome breaks. An inversion that involves the chromosome's constriction point (centromere) is called a pericentric inversion. An inversion that occurs in the long (q) arm or short (p) arm and does not involve the centromere is called a paracentric inversion.

#### Isochromosomes

An isochromosome is a chromosome with two identical arms. Instead of one long (q) arm and one short (p) arm, an isochromosome has two long arms or two short arms. As a result, these abnormal chromosomes have an extra copy of some genes and are missing copies of other genes.

#### Dicentric chromosomes

Unlike normal chromosomes, which have a single constriction point (centromere), a dicentric chromosome contains two centromeres. Dicentric chromosomes result from the abnormal fusion of two chromosome pieces, each of which includes a centromere. These structures are unstable and often involve a loss of some genetic material.

#### Ring chromosomes

Ring chromosomes usually occur when a chromosome breaks in two places and the ends of the chromosome arms fuse together to form a circular structure. The ring may or may

not include the chromosome's constriction point (centromere). In many cases, genetic material near the ends of the chromosome is lost.

SEX-Linked Inheritance

#### **X-Linked Inheritance**

Traits that are determined by alleles carried on the **X chromosome** are referred to as **X-linked**. X-linked alleles require a specific notation:  $X^c$  or  $X^+$  where the "+" represents the **dominant** allele and the lowercase letter the **recessive** allele. Females will have two Xlinked alleles (because females are XX), whereas males will only have one X-linked allele (because males are XY). Most X-linked traits in humans are recessive.



An Ishihara plate can be used to test for red-green colorblindness. Affected individuals may not see the number 6. Image courtesy of Wikimedia Commons

One example of an X-linked trait is red-green colorblindness. Let  $(X^c)$  represent the recessive allele that causes colorblindness and  $(X^+)$  represent the normal dominant allele. Females that are  $X^+X^+$  or  $X^+X^c$  have normal color vision, while  $X^cX^c$  females are colorblind. Males that are  $X^+Y$  have normal color vision, while  $X^cY$  males are colorblind.

#### **Punnett Squares**

To determine the inheritance of red-green colorblindness (or any other X-linked trait),

the genotypes of the parents must be considered. For example, if a mother is a **carrier** for colorblindness  $(X^+X^c)$ , and a father has normal vision  $X^+Y$ , then their sons have a 50% chance of

colorblindness because they inherit their X chromosome from their

	Xc	<b>X</b> <sup>+</sup>
<b>X</b> <sup>+</sup>	<u>X</u> <sup>e</sup> X <sup>+</sup>	<b>X</b> <sup>+</sup> <b>X</b> <sup>+</sup>
Y	<u>X°Y</u>	<b>X</b> <sup>+</sup> <b>Y</b>

mother and their Y chromosome from their father. Their daughters will have a 50% chance of being a carrier  $(X^+X^c)$  and a 50% chance of being completely normal  $(X^+X^+)$  (see figure). A Punnett square can be used to determine any possible genotypic combinations in the parents.

#### Pedigree

Here is a pedigree depicting X-linked recessive inheritance. These traits are often passed from a carrier mother to an affected son. X-linked traits are never passed from father to son. Males are more likely to be affected than females. In this pedigree, the carrier (**heterozygous**) females are indicated; however, they do not express the trait being tracked in this pedigree.



#### KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B Sc BT COURSE CODE: 18BTU201 COURSE NAME: GENETICS UNIT: V BATCH-2018-2021

#### <u>UNIT-V</u>

#### **SYLLABUS**

**GENETICS:** Linkage and recombination of genes in a chromosome crossing, Rules of extra nuclear inheritance, Cytoplasmic inheritance, organism heredity, genomic imprinting, Evolution and population genetics, inbreeding and out breeding, Hardy Weinberg law, Alleic and genotype frequency

#### Unit – V

#### Linkage and Crossing over

#### CHROMOSOME THEORY OF LINKAGE

The chromosome theory of linkage of Morgan and Castle states that:

- 1. The genes which show linkage, are situated in the same pair of chromosomes.
- 2. The linked genes remain arranged in a linear fashion on the chromosome. Each linked gene has a definite and constant order in its arrangement.
- 3. The distance between the linked genes determines the degree of strength of linkage. The closely located genes show strong linkage then the widely located genes which show weak linkage.
- 4. The linked genes remain in their original combination during the course of inheritance.

#### Kinds of Linkage:

Linkage is generally classified on the basis of three criteria viz., (i) Crossing over, (ii) Genes involved and (iii) Chromosomes involved.

**Based on crossing over:** Linkage may be classified into (a) complete and (b) incomplete / partial depending up on absence or presence of recombinant phenotypes in test cross progeny.

(a) Complete linkage: It is known in case of males of *Drosophila* and females of silkworms, where there is complete absence of recombinant types due to absence of crossing over.

(b) Incomplete / partial linkage: If some frequency of crossing over also occurs between the linked genes, it is known as incomplete / partial linkage. Recombinant types are also observed besides parental combinations in the test cross progeny. Incomplete linkage has been observed in maize, pea, *Drosophila* female and several other organisms.

#### **Types of linkage**

The types of linkage are two types.

- 1) Complete linkage
- 2) Incomplete linkage

#### **Complete linkage**

Linked genes inherit together for more generation in a continuous and regular fashion on the same chromosome. This phenomena of inheritance is called complete linkage.

Eg: linkage in Drosophila melanogaster.

#### 1. Complete Linkage

In complete linkage, linked genes inherit together for many generations. Here crossing over does not occur. In complete linkage the genes are closely situated.

Eg. 1. Body colour and shape of wings in male Drosophila.

2. Bent wings and shaven bristles in the 4<sup>th</sup> chromosome of *Drosophila*.

Complete linkage is the phenomenon in which two or moregenes or characters are inherited together Jor a number of generation. In this, genes are closely associated and tend to inherit together. Complete linkage is due to the fact that there occurs no break in the

chromosomes. As a result of complete linkage, the young ones inherit only the parental characters. New characters do not appear among the young ones. So complete linkage produces only parental combination; new combinations do not arise. This phenomenon is very rare. It is found only in male *Drosophila*. The F 1 male hybrid is back crossed with recessive female parent. The F 1 male hybrid produces only two types of gametes in which the linked genes (G and L or g and 1) are inherited together. So only two types of offspring are produced in the F2 generation in equal numbers.



**Incomplete linkage** 

The linked genes which are widely located in chromosome and have changes (meiotic prophase) of separation by crossing over. This phenomena are incompletely linked genes and their inheritance is called incomplete linkage.

*The separation of linked genes during inheritance* is called *incomplete linkage*. The linked genes are separated due to crossing over, chromosomal breaks, etc.

Eg. 1. Body colour and wing shape in female Drosophila.

- 2. Seed colour and seed shape in maize.
- 3. Flower colour and pollen grain shape in sweet pea.
  - In incomplete linkage, the linked genes on certain occasions separate. This leads to the formation of new combinations among the young ones.
  - Incomplete linkage is due to the breakage of chromosomes during gametogenesis. Incomplete linkage is found in *female Drosophila*.
  - This breakage of chromosomes leads to the separation of linked genes and new combinations appear. Because of this new genetic combination the offspring produced in the F2 generation are different from their parent in their phenotype and genotype.
  - So the incomplete linkage involves the accidental breakage of chromosomal segments or linked genes, resulting in new combination of genes.
  - In the below experiment, the F 1 female hybrid produces four types of gametes. Among four types, two types of gametes carry new combinations due to the separation of linked genes. The gene G is separated from *L* and joins with *I*. In the same way the *L* joins with *g*.
  - These combinations are different from the original combination. This type of inheritance is different from the *independent assortment*.
  - If the genes are assorted independently the four types of offspring produced in the F2 generation of the above experiment should be in the 1:1:1:1 ratio.



#### **Importance of linkage**

*Morgan* and *Castle* proposed the chromosomal theory of linkage. The main features of the chromosomal theory of linkage are the following:

- 1. The genes are arranged in a linear fashion on the chromosomes.
- 2. The genes on a chromosome are linked
- 3. Genes showing linkage are located on the same chromosome.
- 4. Linked genes will remain together during inheritance.

5. The distance between the genes will determine the strength of linkage. The closely located genes show strong linkage. Distantly located genes show weak linkage.

**Factors effecting linkage** 

Linkage is affected by the following factors:

1. Distance: Closely located genes show strong linkage while genes widely located show weak linkage.

2. Age: With increasing age the strength of linkage decreases.

3. Temperature: Increasing temperature decreases the strength of linkage.

4. X-rays: X-ray treatment reduces the strength of linkage.

**Based on genes involved:** Depending on whether all dominant or some dominant and some recessive alleles are linked together, linkage can be categorized into (a) Coupling phase and (b) Repulsion phase.

(a) **Coupling phase:** All dominant alleles are present on the same chromosome or all recessive alleles are present on same chromosome.

(b) **Repulsion phase:** Dominant alleles of some genes are linked with recessive alleles of other genes on same chromosome.

**Based on chromosomes involved:** Based on the location of genes on the chromosomes, linkage can be categorized into (a) autosomal linkage and (b) X-chromosomal linkage / allosomal linkage / sex linkage.

(a) Autosomal linkage: It refers to linkage of those genes which are located in autosomes (other than sex chromosomes).

(b) X-chromosomal linkage / allosomal linkage / sex linkage: It refers to linkage of genes which are located in sex chromosomes i.e. either 'X' or 'Y' (generally 'X').

#### LINKAGE GROUPS:

Linkage group refers to a group of genes which are present in one chromosome. In other words, all those genes which are located in one chromosome constitute one linkage group. The number of linkage groups is limited in each individual. The maximum number of linka ge groups is equal to the haploid chromosome number of an organism. For example there are ten linkage groups in corn (2n = 20), seven in garden pea (2n = 14), seven in barley (2n = 14), four in *Drosophila melanogaster* (2n = 8) and 23 in man (2n = 46).

#### **CROSSING OVER**

The term crossing over was first used by Morgan and Cattell in 1912. The exchange of precisely homologous segments between non-sister chromatids of homologous chromosomes is called crossing over.

#### **Types of crossing over:**

Depending upon the number of chiasmata involved, crossing over is of three types.

**Single crossing over**: It refers to the formation of single chiasma between non-sister chromatids of homologous chromosomes. It involves two linked genes (Two point test cross).

**Double crossing over**: It refers to the formation of two chiasmata between non-sister chromatids of homologous chromosomes. It involves three linked genes (Three point test cross).

**Multiple crossing over**: Occurrence of more than two crossing overs between non-sister chromatids of homologous chromosomes is known as multiple crossing over. However, the frequency of such type of crossing over is extremely low.

Mechanism of Meiotic crossing over:

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B Sc BT	COUR	SE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: V	BATCH-2018-2021	

It is responsible for recombination between linked genes and takes place during pachytene stage of meiosis i.e. after the homologous chromosomes have undergone pairing and before they begin to separate. It occurs through the process of breakage and reunion of chromatids. During pachytene, each chromosome of a bivalent (chromosome pair) has two chromatids so that each bivalent has four chromatids or strands (four-strand stage). Generally one chromatid from each of the two homologues of a bivalent is involved in crossing over. In this process, a segment of one of the chromatids becomes attached in place of the homologous segment of the nonsister chromatid and vice-versa. It is assumed that breaks occur at precisely homologous points in the two nonsister chromatids involved in crossing over; this is followed by reunion of the acentric segments. This produces a cross (x) like figure at the point of exchange of the chromatid segments. This figure is called chiasma (which is seen in diplotene stage of meiosis) (pluralchiasmata).

Obviously, each event of crossing over produces two recombinant chromatids (involved in the crossing over) called cross over chromatids and two original chromatids (not involved in crossing over) referred to as noncrossover chromatids. The crossover chromatids will have new combinations of the linked genes, i.e. will be recombinant; gametes carrying them will produce the recombinant phenotypes in test-crosses, which are called crossover types. Similarly, the noncrossover chromatids will contain the parental gene combinations and the gametes carrying them will give rise to the parental phenotypes or noncrossover types. Therefore the frequency of crossing over between two genes can be estimated as the frequency of recombinant progeny from a test-cross for these genes. This frequency is usually expressed as percent. Thus, the frequency of crossing over (%) can be calculated using the formula;

No. of recombinant progeny from a test cross

Frequency of crossing over(%) = ------ x 100

Total number of progeny

#### KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B Sc BT COURSE CODE: 18BTU201

#### COURSE NAME: GENETICS UNIT: V BATCH-2018-2021

#### Kinds of crossing over

#### Single cross over

- When only one chiasma occurs at one point of the chromosome pair is called single cross over.
- It produces two known cross over chromatid and two cross over chromatid.

#### **Double cross over**

- When the crossing over occurs at two point between any points in the same chromosome pair is called double crossing over.
- It produces 4 cross overs

#### **Triple cross over**

- When the cross over occur at 3 point between any 3 points in the same chromosome pair is called triple cross over.
- It produces 6 cross over.

#### Multiple cross over

• When crossing over occur at more than 3 or 4 point between any two or more points in the pair of chromosomes is called multiple cross over.

• It produces multiple cross over.

#### Factors effecting crossing over

- 1) High temperature to increase the frequency of crossing over.
- 2) X ray
- 3) Age

- 4) Some genetical mutations decrease the frequency of crossing over.
- 5) Inversion of chromosome segments suppress the crossing over.

#### Theories about the mechanism of Crossing over

Important Theories That Can Explain the Mechanism of Crossing Over are listed below:

- 1. Janssen's partial chiasma type theory.
- 2. Belling's copy choice theory.
- 3. Darlington's breakage and reunion theory.

#### 1. Breakage and reunion theory:

This theory is based on the assumptions that:

- 1. Prior to crossing over each chromosome of each bivalent get duplicated to form tetrad.
- 2. Crossing over occurs only between non-sister chromatids.

3. Crossing over involves the mechanical breaks in non-sister chromatids due to twisting around each other and reunion or recombination of chromatids take place.

According to this theory first of all, chromatids break and then form chiasmata. Crossing over does not produce chiasmata but it is caused by chiasmata.

#### **2.** Copy choice theory:

This theory was proposed by Belling in 1933. According to this theory, the paired chromosomes duplicate their genes before the fibres join them to form strands.

When the chromosomes are twisted around each other, reciprocal exchange of the chromatids take place during pachytene or just before.

There may be some recombination during the period of DNA synthesis affecting short, unpaired segments of the chromosomes.

A small part of new DNA helix being synthesized may copy a non-sister helix rather than a sister helix to produce recombination in a very short period.

There are two main objections:

1. Only two chromatids out of four involved in crossing over thus newly formed chromatids would be altered by recombination.

2. Duplication should occur during late meiotic prophase but now it is clear that DNA duplication occurs even before syn-apsis.

Hence, these theories assume that crossing over occurs in the interphase rather than pachytene.

#### **3.** Partial chiasma type theory:

According to this theory breaks occur only in two chromatids out of the four at the pachytene stage. These four again rejoined and the chiasmata are formed i.e. chiasmata is the result of cross-ing over.

#### Cytological detection of crossing over.

• The genetical detection of crossing over in Drosophila melanogaster we will know using their (f1 off springs) phenotype characters. But we unable to see under microscopic examination whether the crossing over occur or not, because the chromosomes are in homologous strains or condition.

• This crossing over or blocks had changed places cannot observed until visible marker may be incorporated on the chromosome.

Stern's experiment for cytological detection of cross over

1) Crossing over is the interchange of chromosome parts between homologous chromosome, and this crossing over is proved by Stern, in 1931, on Drosophila melanogaster.

2) The female Drosophila carries XX chromosome and the male Drosophila carries one X chromosome and one Y chromosome.

3) In a type of female Drosophila the two X chromosomes are different from each other.

4) An X chromosome has a piece of Y chromosome attach to it, the other X chromosome has been broken into two unequal segments and it is shorter than the unbroken X chromosome. Thus the two X chromosome are structurally different from the normal X chromosome.

5) In Drosophila red eye (C) is dominant and carnation eye (c) is recessive. Similarly bar eye (B) is dominant and round eye (b) is recessive.

6) The broken X chromosome contains a recessive gene (c) for carnation eye colour and a dominant eye (B) for bar eye, while it is homologous contains C & b.

7) This female having red bar eyes is crossed with a double recessive male, having carnation round eyes.

8) In the absence of crossing over only two types of female gamates are produced, one type having broken X chromosome containing c & B genes, the other type X chromosome having with a piece of Y chromosome attached and contain C & b genes.

9) If crossing over occurs two more type of gamates are produced. One type having C & B on a broken X chromosome with a piece of Y chromosome. So these 4 types of gamates after fertilization will produce 4 types of off spring that are,

• Carnation colour & bar shape eyes

- Red colour & round shape eyes
- Carnation colour & round shape eyes
- Red colour & bar shape eyes

10) The X chromosome of above said 4 types we can identified under the microscopic examination. This experiment proves that inter change of chromosomal material takes place between the homologous chromosomes.

#### Creighton & Mc Clintock's experiment

1) Creighton & Mc Clintock proved experimentally the exchange of chromatids during crossing over in maize.

2) They used to strains of maize which showed difference in the 9th chromosome. Because the 9th chromosome has knob at one end and a cell maker at the other end.

3) The other strain has no knob and no cell maker.

4) In additional to genetical characters are selected they are, colour of Kernal and nature of endosperm. Coloured kernel is dominant (C) & the colourless kernel is recessive (c).

5) Starchy endosperm (WX) is dominant and waxy endosperm (wx) is recessive.

6) A maize with knobbed chromosome, coloured kernel and waxy endosperm is crossed with another maize having knobless chromosome colourless kernel and starchy endosperm.

7) Hybrid maize having heterozygous chromosomes and heterozygous genotype are produced in the F1 generation.

8) The F1 hybrid is test crossed with a double recessive knobless chromosome.

9) The outcome F2 generations have examine genetically and cytologically.

10) The outcome F2 off springs have the following characterestics features.

• Knobbed coloured waxy

Knobbed coloured starchy

- Knobless coloured waxy
- Knobless coloured starchy

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B Sc BT	COUR	SE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: V	BATCH-2018-2021	

11) This result shows the knobbed chromosomes were transformed in meosis process through the crossing over.

#### Significance of crossing over

a. It produces new individuals having new combinations of traits.

b. Crossing over has helped in establishing the concept of linear arrangement of genes.

c. The frequency of crossing over helps in the mapping of chromosomes. i.e., determining the location of the genes in the chromosomes.

d. Selection of useful recombination by geneticists has brought about green revolution in our country.

#### CYTOPLASMIC INHERITANCE

Inheritance due to genes located in cytoplasm (plasmagenes) is called cytoplasmic inheritance. Since genes govering traits showing cytoplasmic inheritance are loc ated outside the nucleus and in the cytoplasm, they are refered to as plasmagenes. The sum total of genes present in the cytoplasm of a cell or an individual is known as plasmon. The plasmagenes are located in DNA present in mitochondria (mt DNA) and in chloroplasts (cp DNA). Together both the DNAs are called organelle DNA. Therefore, this type of inheritance is often referred to as organellar inheritance, plastid inheritance or mitochondrial inheritance. In this, generally, the character of only one of the two parents (usually female) is transmitted to the progeny. Hence such inheritance is usually referred to as extra - nuclear or extra-chromosomal or maternal or uniparental inheritance. The cytoplasmic inheritance is of two types: 1) Plastid inheritance and 2) mitochondrial inheritance.

**Plastidial or Chloroplast Inheritance:** Plastids self duplicated and have some amount of DNA and plays an important role in cytoplasmic inheritance. Plastids have green pigments called chloroplasts. Chloroplasts contain a unique circular DNA (cp DNA) in the stroma that is completely different from the nuclear genome. Some examples of plastid inheritance are given below.

a) Leaf variegation in *Mirabilis jalapa* : The conclusive evidence for cytoplasmic inheritance was first presented by C. Correns in Mirabilis jalapa (Four 'O' clock plant) in 1909. He studied inheritance of leaf variegation in *M. jalapa*. Variegation refers to the presence of white or yellow spots of variable size on the green background of leaves. In *M. jalapa*, leaves may be green, white or variegated. Some branches may have only green, only white or only variegated leaves. Correns made crosses in all possible combinations among the flowers produced on these three types of branches. When flowers from green branch were used as female parent, all the progeny were green irrespective of the phenotype (green, white or variegated) of male parent. Similarly, progeny from crosses involving flowers bloomed on white branches as female parent were all white irrespective of the phenotype of male parent. But in progeny from all crosses involving flowers born on variegated branches as female parent, all the three types i.e. green, white and variegated individuals were recovered in variable proportions.

Female Parent	Х	Male Parent			
Green	Х	Green			
	Х	White	Green		
	Х	Variegated			
White	x	Green			
	х	White	pale Green		
	X	Variegated			
Variegated	Х	Green			
Branarad by Anil Kumar DK and Nichy Sakar, Acct Braf, Danartmant of Diotach, KAHE					
KARPAGAI	KARPAGAM ACADEMY OF HIGHER EDUCATION				
----------------	--------------------------------------	-----------------------	--------------------------	--	--
CLASS:	I B Sc BT	COURSE NAME: GENETICS			
COURSE CODE: 1	18BTU201	UNIT: V	BATCH-2018-2021		
Х	White	Green, w	white and variegated		
Х	Variegated	in variat	ble ratio in each of the		

cases.

The green leaf branches have normal chloroplasts, white branches have mutant chloroplasts and variegated have a mixture of both normal and mutant chloroplasts. The above results indicated that the inheritance is governed by chloroplasts. Since the cytoplasm is contributed to the zygote mainly by female parent, the plastids are transmitted to the zygote from the female parent. Thus the plastids are responsible for variation in the crosses of green, white and variegated leaves.

**b**) **Mitochondrial inheritance:** The inheritance of some characters, such as cytoplasmic male sterility in plants, pokyness in *Neurospora* etc., is governed by mitochondrial DNA (mtDNA).

- a) Cytoplasmic Male Sterility (CMS) in maize : In several crops, cytoplasmic control of male sterility is known. In maize, cytoplasmic male sterility (CMS) is governed by mitochondrial DNA. In such cases, if female parent is male sterile, F1 progeny also will be male sterile, because cytoplasm is mainly derived from female parent.
- **b**) **Pokyness in** *Neurospora: Neurospora*, which is a breadmold has two strains i.e. wild and poky. The wild strain has normal growth. While the poky which is a mutant has very slow growth. A cross betw een a poky female and a wild male produce only poky progeny. In reciprocal cross (a cross between wild female and poky male) all the progeny would be wild. This suggests the presence of cytoplasmic inheritance because only difference between the reciprocal crosses is in the main contributor of cytoplasm.

#### **Characteristic Features of Cytoplasmic Inheritance**

**a. Reciprocal difference:** Reciprocal crosses show marked differences for characters governed by plasmagenes. In most cases, plasmagenes from only female parent are transmitted and hence this phenomenon is also called uniparental inheritance.

**b.** Lack of segregation: In general, F1, F2, F3 and subsequent generations do not show segregation for a cytoplasmically inherited trait, as F1 individuals receive plasmagenes from female parent only.

**c. Somatic segregation:** Plasmagenes generally show the features in somatic tissues such as leaf variegation features which is of rare occurrence in case of nuclear genes.

**d.** Association with organelle DNA: Several plasmagenes have been shown to be associated either with chloroplast or mitochondrial DNA. For example: Cytoplasmic Male Sterility (CMS) in sorghum and maize is associated with mitochondrial DNA.

**e. Nuclear transplantation:** Nuclear transplantation means nucleus of a cell is removed and replaced by nucleus of another genotype from a different cell. If nuclear transplantation reveals a trait to be governed by genotype of cytoplasm and not by that of nucleus, it clearly indicates that the trait or character is governed by cytoplasmic inheritance.

**f. Mutagenesis:** Some mutagens are highly specific mutagens which act only on the plasmagenes and do not affect nuclear genes Eg; ethidium bromide, Induction of mutations by such agents or chemicals in a gene clearly indicates that it is a plasmagene.

**g. Lack of chromosomal location:** In many organisms extensive linkage maps of nuclear genes are available. If a gene is shown to be located in one of these linkage groups, obviously it cannot be a plasmagene.

**h. Transfer of nuclear genome through back crosses:** Nucleus of a variety or species may be transferred into cytoplasm of another variety or species through repeated back crossing with former, which is used as recurrent male parent. Lines produced in this way are called alloplasmic lines, since they have cytoplasm and nucleus from different species.

**i. Lack of association with a parasite or symbiont or virus:** Only those cytoplasmically inherited traits which are not associated with parasites, symbionts or viruses can be regarded to be governed by plasmagenes.

#### **Extranuclear Inheritance**

It is the transmission of genes that occur outside the <u>nucleus</u>. It is found in most <u>eukaryotes</u> and is commonly known to occur in cytoplasmic <u>organelles</u> such as mitochondria and <u>chloroplasts</u> or from cellular parasites like viruses or bacteria.

#### **Extranuclear Inheritance of Organelles**

Mitochondria are organelles which function to produce energy as a result of <u>cellular respiration</u>. Chloroplasts are organelles which function to produce sugars via <u>photosynthesis</u> in plants and algae. The <u>genes</u> located in <u>mitochondria</u> and chloroplasts are very important for proper cellular function, yet the genomes replicate independently of the DNA located in the nucleus, which is typically arranged in chromosomes that only replicate one time preceding cellular division. The extranuclear genomes of mitochondria and chloroplasts however replicate independently of cell division. They replicate in response to a cells increasing energy needs which adjust during that cells lifespan. Since they replicate independently, genomic recombination of these genomes is rarely found in offspring contrary to nuclear genomes, in which recombination is common. Mitochondrial disease are received from the mother, sperm does not contribute for it.

#### **Extranuclear Inheritance of Parasites**

Extranuclear transmission of viral genomes and symbiotic bacteria is also possible. An example of viral genome transmission is <u>perinatal</u> transmission. This occurs from mother to fetus during the perinatal period, which begins before birth and ends about 1 month after birth. During this time viral material may be passed from mother to child in the bloodstream or breastmilk. This is of particular concern with mothers carrying <u>HIV</u> or <u>Hepatitis C</u> viruses. Examples of cytoplasmic <u>symbiotic</u> bacteria have also been found to be inherited in organisms such as insects and protists.

#### **Types of Extranuclear Inheritance**

Three general types of extranuclear inheritance exist. These are <u>vegetative</u> segregation, uniparental inheritance and biparental inheritance.

• **Vegetative segregation** results from random replication and partitioning of cytoplasmic organelles. It occurs with chloroplasts and mitochondria during mitotic cell divisions and results in daughter cells that contain a random sample of the parent cell's organelles. An example of vegetative segregation is with mitochondria of asexually replicating yeast cells.

#### Maternal inheritance

- The determination of the phenotype of offspring by the genotype of female parent is called *maternal inheritance* or *uniparental inheritance or maternal effect*.
- Example: Pattern of shell coiling in snail.
- oHere the shell coiling is determined by the genotype of the mother and not by the individual's own genotype.

#### **Examples for cytoplasmic inheritance**

Plastid inheritance in Mirabilis Shell-coiling in snail Kappa particles in Paramecium Cytoplasmic male sterility in maize Sigma virus in *Drosophila melanogaster* Milk factor in mice

#### LEAF VARIEGATION IN PLANTS

- The first example of cytoplasmic inheritance was reported by Correns (1909) in a variegated variety of the four-o'clock plant *Mirabilis jalapa*.
- Variegated plants have some branches which carry normal green leaves, some branches with variegated leaves (mosaic of green and white patches) and some branches which have all white leaves.
- Flowers on wholly green branches produce seeds that grow into normal plants.
- Flowers on variegated branches yield offspring of three kinds- green, white and variegated in variable proportions.

- Flowers from branches wholly white produce seeds that grow into white plants that is without chlorophyll.
- But in every case the source of pollen has no influence on the offspring.
- In other words, the phenotype of the progeny always resembled the female parent and the male made no contribution at all to the character. So cytoplasm of the egg influences the type of leaf in Mirabilis.
- The explanation for this unusual pattern of inheritance is that the genes concerned are located in the *plastids* within the cytoplasm, not in the nucleus and are therefore transmitted only through the female parent.
- Plastids are of two types, namely green *chloroplasts* and colourless *leucoplasts*.
- *Green* branches contain Green plastids in their leaves, Variegated branches contain Green plastids and Colourless plastids and Colourless branches are due to the presence of Colourless plastids.



#### SNAIL SHELL COILING

- The classic phenotype which exhibits maternal effects is coiling direction of snail shells.
- Shell coiling in *Limnaea peregra*, a fresh water snail, is of two types, Dextral (clockwise) and Sinistral (anticlockwise).
- The dextral shell is dominant and is controlled by dominant gene D.
- The sinistral shell is recessive and is controlled by recessive gene d.
- The following crosses were made between pure line snails.
- When dextral female (DD) was crossed with sinistral male (dd), all the offsprings of F1 generation (Dd) have dextral coiling.
- If sinistral female (dd) is crossed with dextral male (DD), the offspring have Dd genotype but coiling is sinistral.

- In the above two crosses, the F1 snails have the same genotypes.
- The F1 phenotype is not the same for both crosses.



- The coiling phenotype that is seen in the offspring is controlled by the genotype of the mother.
- In the first cross, the offspring has dextral shell because the mother's genotype is DD.
- In the second cross, the offspring has sinistral shell because the mother's genotype is dd.
- Since, zygote receives whole of its cytoplasm from the egg, the direction of shell coiling in the offspring is governed by cytoplasm of the mother.

#### **KAPPA PARTICLES IN PARAMECIUM**

- T. M. Sonneborn described the inheritance of some cytoplasmic particles known as kappa and their relation to nuclear gene in the common cillate protozoan, *Paramecium aurelia*.
- There are two strains of Paramecium. They are killer and sensitive.
- Killer strain produces a toxic substance called paramecin that is lethal to other individuals called "sensitives".
- The production of paramecin in killer type is controlled by certain cytoplasmic particles known as kappa particles. The sensitive strains lack these particles.



**Binary Fissio** 

Killers

Prepared by Anil Kumar PK and Nishu Sekar, Asst Prof, Department of Biotech, KAHE

**Binary Fission** 

Killers

Sensitives

- The development of a particular type depends upon the duration of cytoplasmic exchange
- oIf conjugation is normal, i.e., lasts only for a short time, and no exchange of cytoplasm takes place between the two, both killers and non-killers
- (sensitive) are produced.
- oHowever in rare or prolonged conjugation (i.e., lasting for long time) the cytoplasmic bridge between the two conjugants is larger. In such cases, in addition to the nuclear material, the cytoplasmic materials are also exchanged.
- oDuring this cytoplasmic exchange, the kappa particles present in the cytoplasm of the killer type enter the non-killer type and convert it into a killer type. So all the offspring produced by the exconjugants are killer type.
- This shows that a Paramecium becomes a killer when it receives kappa particles and it becomes a sensitive when it does not receive kappa particles.

• Uniparental inheritance occurs in extranuclear genes when only one parent contributes organellar DNA to the offspring. A classic example of uniparental gene transmission is the <u>maternal inheritance</u> of human mitochondria. The mother's mitochondria are transmitted to the offspring at <u>fertilization</u> via the egg. The father's mitochondrial genes are not transmitted to the offspring via the sperm. Very rare cases which require further investigation have been reported of paternal mitochondrial inheritance in humans, in which the father's mitochondrial genome is found in offspring Chloroplast genes can also inherit uniparentally during <u>sexual reproduction</u>. They are historically thought to inherit maternally, but paternal inheritance in many species is increasingly being identified. The mechanisms of uniparental inheritance from species to species differ greatly and are quite complicated. For instance, chloroplasts have been found to exhibit maternal, paternal and biparental modes even within the same species.

• **Biparental inheritance** occurs in extranuclear genes when both parents contribute organellar DNA to the offspring. It may be less common than uniparental extranuclear inheritance, and usually occurs in a permissible species only a fraction of the time. An example of biparental mitochondrial inheritance is in the <u>yeast</u>, <u>Saccharomyces cerevisiae</u>.

#### Hardy-Weinberg Principle

The Hardy-Weinberg Principle (also Hardy Weinberg Equilibrium) states that the <u>allele</u> <u>frequency</u> (and genotype frequency) of a <u>population</u> remains constant over generations, unless a specific factor or combination of factors disrupts this equilibrium. Such factors might include <u>non-random mating</u>, <u>mutation</u>, <u>natural selection</u>, <u>genetic bottlenecks</u> leading to increased <u>genetic drift</u>, the immigration or emigration of individuals (<u>gene flow</u>) or <u>meiotic</u> <u>drive</u>. The Hardy-Weinberg Equilibrium does not actually exist in nature because one or more of these factors is *always* in play.

The concept of an equilibrium exists instead as a baseline against which to measure genetic change between generations. According to the Hardy-Weinberg principle, then, changes in allele frequency (and thus evolution) would be theoretically impossible if the following conditions were met: 1.There mutation was no There 2. were no selective pressures 3. The population size was infinite (this would bring the rate of genetic drift infinitely close to zero)

4. All members of the population breeding were 5. All breeding random was 6. All individuals produced the of offspring same amount 7. There was no migration of individuals into, or out of, the population (i.e. the rate of gene flow was zero)

Represent	epresenting		the		equilibrium			mathema	tically:		
When	considering	a	lo	cus	th	nat	has	two	О	alleles, A	and a:
The	frequency	of	A (the		domi	nant	al	lele)	is	de	enoted <i>p</i>
The	frequency	of	a (the		recess	sive	al	lele)	is	de	noted q
And $p + q$	= 1, since	every	locus	in	the	popul	lation	must	carry	either	allele.

KARPAGAM ACADEMY C	OF HIGHER EDU	JCATION	
CLASS: I B Sc BT	COUR	SE NAME: GENETICS	
COURSE CODE: 18BTU201	UNIT: V	BATCH-2018-2021	

If the population is in Ha	rdy-Weinberg equili	brium, then the alleles	are distributed e	evenly among
heterozygotes		and	1	nomozygotes:
Dominant homozygotes	(genotype AA) are	denoted <i>p</i> ^2 because	the probability	of inheriting
two	dominant	alleles		is <b>p*p</b>
Recessive homozygotes	(genotype <b>aa</b> ) are de	noted $q^2$ because the	probability of i	nheriting two
recessive				allelesis <b>q*q</b>

Heterozygotes (genotype Aa) are denoted 2pq because the probability of inheriting both alleles is (p\*q)+(q\*p) And  $p^2 + 2pq + q^2 = 1$ , since every individual in the population must be one of these genotypes This equation can be used, for example, to predict the frequency of carriers of a disease in a population. Consider the autosomal recessive disease, phenylketonuria. If the frequency of the recessive (in this case, harmful) allele is 1% (q = 0.01), then the number of people who suffer (i.e. who are homozygous recessive) is  $q^2 = 0.0001$  or 0.01% of the population.

The number of carriers - or heterozygotes - is  $2pq = 2 \times 0.99 \times 0.01 = 0.198$ , or 1.98% of the population. That means that carriers of the disease exist in the population at a frequency of almost 200 times more than actual sufferers. This can help us to identify the likelihood of one carrier mating with another, and potentially producing an offspring who suffers from the condition. One exception to the Hardy-Weinberg principle is a phenomenon called recurrent *mutation*. When looking at the rate of mutation in a population, forward mutations (i.e. those that cause functional genes to become non-functional) are more common than reverse mutations (i.e. those that cause non-functional genes to restore functionality). This is simply because a forward mutation can involve any number of base changes, while a reverse mutation requires the specific reversal of the change that initially made the gene non-functional. Hence, forward mutations occur at far greater rate than reverse mutations, and will proliferate in a population when they pose no selective disadvantage. This is evident in the proliferation of the O allele (which does not code for a glycosyltrasnferase enzyme) in the ABO blood-grouping system, at the expense of В the Α alleles. and Thus if q to determines the frequency of the non-functional allele, and  $\mu$  determines forward mutation rate while v determines reverse mutation rate, where  $\mu$  is taken to be 10 times the value of *v*:*q*(equilibrium) =  $\mu / (\mu + v)$ 

CLASS: I B Sc BT

#### COURSE CODE: 18BTU201

#### COURSE NAME: GENETICS UNIT: V BATCH-2018-2021

#### **NATURAL SELECTION**

History of Evolution:

- Plato & Aristotle believed species were fixed & could be arranged according to their complexity
- In the mid eighteenth century, Carolus Linnaeus developed a system of classification that called binomial nomenclature
- George Cuvier, in the eighteenth century, explained changes in the fossil record by proposing that a whole series of catastrophes (extinctions) and re-populations from other regions had occurred giving the appearance of change over time
- Prior to Darwin, it was thought that the world was young & species did not change
- Lamarck (1744-1829) was first to state that descent with modification occurs and that organisms become adapted to their environments
- Inheritance of acquired characteristics was the Lamarckian belief that organisms become adapted to their environment during their lifetime and pass on these adaptations to their offspring
- Lamarck believed that the long necks of giraffes evolved as generations of giraffes reached for ever higher leaves; known as the Law of Use & Disuse



- Because it is supported by so many lines of evidence, evolution is no longer considered a hypothesis
- Evolution is one of the great unifying theories of biology

Darwin's Background & Voyage:

- His nature was too sensitive to become a doctor like his father so he studied divinity
- He attended biology and geology lectures and was tutored by the Reverend John Henslow who arranged his trip on the HMS Beagle
- In 1831, at the age of 22, Charles Darwin accepted a naturalist position aboard the ship HMS Beagle & began a five-year voyage around the world



- He read Principles of Geology by Charles Lyell that stated that the observed massive geological changes were caused by slow, continuous processes (erosion, uplifting...)
- Darwin carried this book with him on his voyage as he witnessed Argentina coast earthquakes raising the earth several feet, & marine shells occurring far inland and at great heights in the Andes
- Darwin's many observations led him to the idea that species slowly change over time
  - Darwin's comparison of the animals of South America and the Galapagos Islands caused him to conclude that adaptation to the environment can cause diversification, including origin of new species

Examples: Patagonian hares replaced rabbits in the South American grasslands

The Galapagos Islands: Volcanic islands off the South American coast

- Island species varied from the mainland species, and from island-to-island
- Each island had either long or short necked tortoises depending on the island's vegetation

CLASS: I B Sc BT COURSE CODE: 18BTU201 COURSE NAME: GENETICS UNIT: V BATCH-2018-2021



- Finches on the Galapagos Islands resembled a mainland finch, but there were more types
  - Bill shapes are adaptations to different means of gathering food.



• Galapagos finch species varied by nesting site, beak size, and eating habits

#### CLASS: I B Sc BT COURSE CODE: 18BTU201

#### COURSE NAME: GENETICS UNIT: V BATCH-2018-2021



Darwin's Theory of Evolution:

- An adaptation is a trait that helps an organism be more suited to its environment
- Darwin decided adaptations develop over time
- Natural selection was proposed by both Alfred Russell Wallace and Darwin as a driving mechanism of evolution
- Darwin and Wallace both read an essay by Thomas Malthus that proposed that human populations outgrow resources so there is a constant struggle for existence



- Fitness is a measure of an organism's reproductive success
- Organisms most fit to reproduce are selected by environment which results in adaptation of the population
- Natural selection is also called "survival of the fittest"
- Conditions for natural selection include:
  - a. Variations exist among members of a population
  - b. Many more individuals are produced each generation than will survive
  - c. Some individuals are better adapted so they survive & reproduce
  - d. Members of a population compete for food, space, mates...
- Variations that make adaptation possible are those that are passed on generation to generation
- Extinction occurs when previous adaptations are no longer suitable to a changed environment

On the Origin of Species by Darwin:

- After the HMS Beagle returned to England in 1836, Darwin waited over 20 years to publish
- Darwin was forced to publish <u>Origin of Species</u> after reading a similar hypothesis by Alfred Russell Wallace
- Both men concluded that life forms arose by descent from a common ancestor, and that natural selection is the mechanism by which species change and new species arise

Fossil Evidence:

- Fossils are relics or impressions of ancient organisms
- Most fossils are found in layers (strata) of sedimentary rock



- The fossil record traces history of life and allows us to study history of particular organisms
- Through radioactive dating, geologists estimate the age of the earth at about 4.6 billion years
- Fossils are at least 10,000 years old and include skeletons, shells, seeds, insects trapped in amber, imprints of organisms, organisms frozen in ice (wooly mammoth), or trapped in tar pits (saber-toothed tiger)
- Transitional forms reveal links between groups (Example: Therapsids were mammal-like reptiles and Pterosaurs were bird like reptiles)

CLASS: I B Sc BT COURSE CODE: 18BTU201 COURSE NAME: GENETICS UNIT: V BATCH-2018-2021



PTEROSAURS

Biogeographical Evidence:

- Biogeography is the study of the geographic distribution of life forms on earth
- Physical factors, such as the location of continents, determine where a population can spread
- Example: Placental mammals arose after Australia separated from the other continents, so only marsupials diversified in Australia

Anatomical Evidence:

- Organisms have anatomical similarities when they are closely related because of common descent
- Homologous structures in different organisms are inherited from a common ancestor have have similar structures
- Example : Vertebrate forelimbs contain the same sets of bones organized in similar ways, despite their dissimilar functions

CLASS: I B Sc BT COURSE CODE: 18BTU201 COURSE NAME: GENETICS UNIT: V BATCH-2018-2021



- Analogous structures are inherited from different ancestors and have come to resemble each other because they serve a similar function
- Example: Bird wing & bat wing are both for flight but they are structurally different
- Vestigial Structures are remains of a structure that is no longer functional but show common ancestry
- Example: Humans have a tailbone but no tail

Embryological Evidence:

- During development, all vertebrates have a post-anal tail and paired pharyngeal pouches
- Organisms that show similarities in their embryonic development may have a common ancestry

CLASS: I B Sc BT COURSE CODE: 18BTU201 COURSE NAME: GENETICS UNIT: V BATCH-2018-2021



Biochemical Evidence:

- Almost all living organisms use the same basic biochemical molecules, e.g., DNA, ATP, enzymes ...
- Similarities in amino acid sequences, DNA codes, etc. can be explained by descent from a common ancestor

Examples of Evolution in Modern Times:

- Peppered moth light colored vs. dark colored (industrialization influence) Manchester, England
- Insect resistance to insecticides
- Bacterial resistance to antibiotics

CLA55.1D 5C D1	COUR	SE NAME: GENETICS
COURSE CODE: 18BTU201	UNIT: V	BATCH-2018-2021

During which stage of majorie crossing over takes place?	Lantotana	Zurrotano	Diakinasis	Bachatana
Meiosis – II is :	Cell division	Mitotic division	Commonly cell elongation	Reduction division
Significance of meiosis is associated with : Who is the father of appendix 2	DNA duplication Gregorichennmendel	Asexual reproduction	Sexual reproduction	Growth of the body Tschermak
Which plant Mendel chose for his experiment	Maize	Garden pea	Bean	Wheat
Mendel chose how many pair of characters for his studies	8	3	7	9
Progeny anses with the fusion of is the ratio of law of segregation	03:01	Pemale gametes 01:03	9:3:3:1	03:03:01
Mendel selected varietes for his studies	23	22	21	20
Laws of inneritance are Complete set of genes are known as	allele	law of independent assortment genome	law of seggregation and independent chromosome	nucleotide
Study of genes are called as	genome study	gene	genetics	molecular study
Embryo contain pairs of chromosomes, half from each parent Each unique form of a single gene is called as	21 genome	23 gene	24 chromosome	43 allele
means the information in a given genes is not always exactly the in every individual in that species	heritability	inheritance	assortment	polymorphism
Allele dominates the instruction from another, it is called the allele	recessive	dominant	embryo	zygote
Mendel laws projosed at Mendel laws re-discovered at	1865 and 1866 1900	1885	1882	1863
Dominant allele hide the phenotypic effects of the recessive allele known as	law of inheritance	law of dominance	law of segregation	law of heritance
During gamete formation, the alleles for each segregate from each other and carries only one allele of each gene is called as Genes for different traits independently during the formation of gametes are known as law of	law of dominance	law of inheritance	law of segregation	law of heritance
Allele that is overridden is called the allele	Recessive	dominant	embryo	zygote
is the ratio of dihybitd cross	03:01	01:03	9:3:3:1	09:01
Alleles are also known as	1.5 hereditarymetrial	ri	12 hereditary determinants	14 chromosome
An individual contains only one allele at the allelic pair are	Homozygote	heterozygote	embryo	genome
The specific allelic combination for a certain gene or set of genes are known as The cruss of nur individual to a homeworking the result of the certain of the cruss of the certain of the	phenotype dihybrid gross	allelic pair grossing over	gamete monohubrid groce	genotype test gross
A cross between parents that differ at a single gene pair are called as cross	Monohybrid	dihybrid	test	crossing over
The offspring of two parents that are homozygous for alternative alleles of a gene pair are known as Mendal observed traits that are easily recognized	Monohybrid	dihybrid 4	test cross	crossing over
Flower colour is apparently only occur in one of two forms	red or purple	purple or white	white or red	yellow or purple
Seed colour is apparently only occur in one of two forms	red or yellow	yellow or green	red or white	green or red
Fod colour is apparently only occur in one or two forms The traits physically shows up in the organism are called	yenow or green hereditary	inheritance	green or white Phenotype	genotype
disease caused by a single allele of a gene and are inherited in families	mental disorders	genetic disorders	infectious disease	sex linked disorders
In law of segregation pair of chromosomes separate during the formation of gametes Mendel chose pea plant for his experiment because	homologous easy to breed	heterozygous short breeding period	allelic it had some measurable character	two all the three options
In F1 generation of law of segregation pod characteristic had disappeared	green	yellow	red	white
Law of dominance states that a dominant allele express itself in a cross	monohybrid	dihybrid	trihybrid X-II	penta hybrid
in me r-2 generation of raw of segregation pod variant appeared in a quarter Mendel's law of segregation states that allele pairs separate or segregate during formation	green egg	embryo	i enow zygote	gamete
In law of dominance offspring that are hybrid for a trait will have only the trait in the phenotype	recessive	dihybrid	dominant	monohybrid
Dominant atteles are expressed exclusively in Recessive traits are expressed only if the organism is for the recessive allelo	homozygous	heterozygote homologous	homozygote heterozygote	homologous heterozygous
Mendal's approach was transfer of pollen from the one pea plant to the pistil of the other	stamen	pollen grains	gamete	ovule
The law of independent assortment refers that any plant contain genes	determines flower colour	detetmines length	determines shape	All the three options
is the interaction of alleles belongings to the same gene pair Alleles segregates independently of the other pairs of alleles during gamete formation. This is known as law of	extra allelic interaction Dominance	intra allelic interaction inheritance	inter allelic interaction independent assortment	anenc interaction segregation
Most familiar animals and plants have paired chromosomes that are described as	diploid	polyploidy	haploid	monoploid
The gametes have only one copy of each chromosome that are described as	diploid Round	polyploidy	haploid round and wrinklad	monoploid
In F2 generation of law of dominance all seeds were found to be	Round	spherical	wrinkled	round and wrinkled
In the law of dominance the ratio of the F2 generation is	03:01	01:03	9:3:3:1	03:03:01
Which cross would best illustrate mendel's law of segregation In the cross Yv×Yv, what percent of offspring would have the same phenotype as the parents	TT×tt 25%	Hh×hh 50%	Bb×Bb 100%	rr ×rr 75%
Crossing two hybrid organisms results in which phenotypic ratio	01:02:01	9:3:3:1	03:01	01:01
Mendal formulated his law of segregation after he had	studied F2 offspring	studied F1 offspring	produced mutations	produced hybrids
Which cross would produce phenotypic ratios that would inustrate the law of dominance The gene expression of a trait in an organism is reffered to as	genotype	an allele	phenotype	tt×tt independent assortment
How many alleles for one trait are normally found in the genotype of an organism	1	2	3	4
The genes present in an organism represent the organism's	genotype looking at it	phenotype analysing it genes	physical traits crossing it with a recessive plant	gene expression eating it
Unit 2	looking it h	unity sing it genes	crossing a what a recessive plant	
In both the dominant and recessive characters occur side by side in F1 hybirds to give a mixed colour	dominance	co dominance	incomplete dominance	complete dominance
Individual has an intermediate phenotype between the two parents, as there is no dominant allele. This pattern of inheritance is called Incomplete dominance occurs in the	ncomplete dominance polygenic inheritance	co dominance inheritance	complete dominance monogenic inheritance	dominance tetragenic inheritance
is a condition where the phenotype of the heterozygote lies outside the phenotypical range of both homozygous parents	complete dominance	incomplete dominance	co dominance	over dominance
inheritance is the transmission of genes that occur in cytoplasmic organelles	chromosomal inheritance	non chromosomal inheritance	cytoplasmic inheritance	mitochondrial
L Moniacimic unharmanca alco Enoura ac				
Cytopiasmic inneritance also known as Genes located in cell organelle in the cytoplasm are	cytoplasmic genes	extra chromosomal inheritance chromosomal genes	non chromosomal inheritance mitochondrial genes	non chromosomal genes
Cytoplasmic internance also known as Genes located in cell organelle in the cytoplasm are Gene present in cytoplasm and mitochondria are known as	cytoplasmic genes chromosomal inheritance	extra chromosomal inheritance chromosomal genes non chromosomal inheritance	non chromosomal inheritance mitochondrial genes extra chromosomal	non chromosomal genes mitochondrial inheritance
Cytoparam: mnertance also known as Genes located in cell organelle in the cytoplasm are Gene present in cytoplasm and mitochondria are known as Genes sociated in cell organelle in the cytoplasm are Gene present in cytoplasm and mitochondria are known as	cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal genes inter chromosomal	non chromosomal inheritance mitochondrial genes extra chromosomal mitochondrial genes extra chromosomal	non chromosomal genes mitochondrial inheritance non chromosomal genes chromosomal
Cytoplasme innertiance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are and Genes located in cell organelle in the cytoplasm are and Genes located in cell organelle in the cytoplasm are and Genes located in cell organelle in the cytoplasm are dimensioned and Genes located in cell organelle in the cytoplasm are dimensioned and Holoroplasm are exercically and have their own system of heredity	cytoplasmic genes cytoplasmic genes cytoplasmic genes inter chromosomal normal	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal genes inter chromosomal abnormal	non chromosomal inheritance mitochondrial genes extra chromosomal mitochondrial genes extra chromosomal stable	mitochondrial inheritance non chromosomal genes mitochondrial inheritance non chromosomal genes chromosomal autonomous
Cytopiame innertiance also known as Genes located in cell organelle in the cytopiasm are Genes located in cell organelle in the cytopiasm are Gene present in cytopiasm and mitochondria are known as Didroopiasm are centerically and have their own system of herodity During excual reproduction chloropiast chromosomes are only transmitted through the of the egg cell is the net where of home michowheredite ansees are historical to the context of the contex	circumsonial intertance cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal normal nucleus in chromidi l'abasitance	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal genes inter chromosomal abnormal cytoplasm careluesic inheritance	non chromosomai inheritance mitochondrial genes extra chromosomal mitochondrial genes extra chromosomal stable mitochondria	mitochondria inieritance non chromosomal genes mitochondrial inheritance non chromosomal genes chromosomal autonomous stamen
Cytopiamic innertiance also known as Cytopiamic innertiance also known as Genes located in cell organile in ithe cytopiasm are Genes located in cell organile in ithe cytopiasm are Genes located in cell organile in the cytopiasm are Genes located in cell organile in the cytopiasm are Genes located in cell organile in the cytopiasm are Genes located in cell organile in the cytopiasm are Genes located in cell organile in the cytopiasm are Gene present in cytopiasm and mitochondria are known as Chioropiasts are genetically and have their own system of heredity During secual reproduction chicropiast chromosomes are only transmitted through the of the egg cell is the study of how mitochondrial genes are inherited Mitochondrial interiance also known as	chromosonal inheritance cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal normal nucleus mitochondrial inheritance cytoplasmic inheritance	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal genes inter chromosomal abnormal cytoplasmi cytoplasmic inheritance extranuclear inheritance	non chromosomal inheritance mitochondrial genes extra chromosomal mitochondrial genes extra chromosomal stable mitochondria extra nuclear inheritance chromosomal inheritance	nniceronarian internance non chromosomal genes mitochondrial inheritance non chromosomal genes chromosomal autonomous stamen chromosomal inheritance both cytoplasmic and extranucle
Cytoplasmic intertaince also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are more direction organicate by a cell organicate in the cytoplasmic direction of the cell organicate in the cytoplasmic di	chromosona internance cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal normal nucleus mitochondrial inheritance cytoplasmic inheritance cytoplasmic inheritance	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal genes inter chromosomal abnormal cytoplasmic inheritance extranuclear inheritance inheritance	non chromosomal inheritance mitochondrial genes extra chromosomal stable mitochondrial genes extra chromosomal extra nuclear inheritance chromosomal inheritance chromosomal inheritance	nnochonaria ninerianee non chromosomal genes ehromosomal genes ehromosomal autonomous stamen ehromosomal inheritanee both evtoplasmic and extranucle non chromosomal inheritanee
Cytoplasmic miteriance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Gene present in cytoplasm and mitochondria are known as Chloroplasm are gene cilcully — and have their own system of herodity During sexual reproduction chloroplast chromosomes are only transmitted through theof the egg cell is the study of how mitochondrial genes are inherited Mitochondrial inheritance task oknown as Trait controlled by a gene of the modern but expressed in the progeny is called Extramelear inheritance fast formor Jinterinate Non meredical inheritance fast formored the second to the seco	caronisonan inner inner cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal nucleus mitochondrial inheritance maternal effect cytoplasmic inheritance maternal effect	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal genes inter chromosomal abbornal cytoplasm cytoplasm cytoplasm cytoplasm chromic inheritance extranuclear inheritance inheritance mitochondria Catl correns	non chromosomal inheritance mitochondrial genes extra chromosomal mitochondrial genes extra chromosomal stable mitochondria extra nuclear inheritance chromosomal inheritance chromosomal inheritance mendelian Buth saver	Innoctionata internance non chromosomali genes mitochondrial inheritance non chromosomal genes chromosomal autonomous stamen chromosomal inheritance both evtoplasmic and extramucle non chromosomal inheritance non medelian Herschel Mitchell
Cytopamic minertance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Gene present in cytoplasm and mitochondria are known as Chloroplasts are centically and have their own system of hereditiv During sexual reproduction chloroplast chromosomes are only transmitted through the of the egg cell is the study of how mitochondrial genes are inherited Mitochondrial inheritance that oknown as Trait controlled by a gene of the mother but expressed in the progeny is called Extransclear inheritance is a form of inheritance Non medicalin inheritance first discovered by	circumonian internative cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal nucleus mitochondrial inheritance cytoplamic inheritance maternal effect chromosomal Mendel 1904	extra chromosomal inheritance chromosomal genes non chromosomal inheritance chromosomal abnormal abnormal cytoplasmi inheritance extramuclear inheritance extramuclear inheritance mitochondria Carl correns 1907	non chromosomal inheritance mitchoudrial genes extra chromosomal mitchoudrial genes extra chromosomal stable mitochoudrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance mendelian Ruth sager 1908	mitochonican internance non chromosomali genes mitochondrial inheritance non chromosomal genes chromosomal genes stamen chromosomal inheritance both cytoplasmic and estramacle non chromosomal inheritance non mendelian Hershel Mitchell 1902
Cytoplasmic interfance also known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Gene present in cytoplasm and mitochondria are Known as Choroplasm are second and have their ono system of heredity During secual reproduction chloroplast chromosomes are only transmitted through the of the egg cell is the study of how mitochondria genes are inherited Mitochondrial inheritance that form of inherited Extramelear inheritance first Genovered by Mon meddelin inheritance first Genovered by Low meddelin inheritance first Genovered by Unignerated inheritance is ferm of inheritance Unignerated inheritance is ferm of inheritance Low meddelin inheritance first Genovered by Low meddelin inheritance first Genovered by Monoveredin inheritance first Geno	consolution interaction compared and a second and a second chromosomal inheritance cytoplasmic genes inter chromosomal necleus necrosal necleus cytoplasmic inheritance mitocheantic inheritance cytoplasmic inheritance cytop	extra chromosonia inheritance chromosonia genes non chromosonia genes inter chromosonia abhormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cheritance extramoclear inheritance inheritance inheritance Carl corrers 1907 Mendelian Mendelian	non chromosomal inheritance mitchondrali genes extra chromosomal mitchondrali genes extra chromosomal i stabile mitochondral genes extra nackear inheritance extra nackear inheritance chromosomal inheritance chromosomal inheritance mendelian Rouh sager 1908 Ruha sager	nuccentrating speeds non-dromosouring speeds mitochondrial inheritance non-dromosourial genes chromosourial inheritance chromosouri inheritance chromosouri inheritance both evoplasmic and extramacle non-dromosouri linheritance men methian inheritance men methian heritance Heritah Mitchell Heritah Mitchell
Cytoplasme innertiance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are is the study of how mitchchordinal genes are inherited Genes located in cell organelle in the cytoplasm are genes of the other study of the cell organized in the progeny is called Extranuclear inheritance frant discovered by Non mendelian inheritance frant discovered by Changement Interinates is an from of inheritance Transmission of genes that are located in mitchchordin as termed	circumoniar internance cytoplasmi genes chromosomal inheritance cytoplasmi genes inter chromosomal nucleus mitochondrial inheritance mitochondrial inheritance extoplasmic inheritance matemal effect chromosomal Mendel 1904 nonmendelian Matemal matemalificatione	extra chromosonia inheritance chromosonia genes non chromosonia genes inter chromosonia abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cheritance extramclear inheritance inheritance Mendelian chromosonial cytoplasmi inheritance	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance nendelian Ruth sager 1908 Ruth sager mendelian chromesomal inheritance chromosomal inheritance	minicronouran internance non chromosomali genes mitochondrial inheritance non chromosomali genes chromosomal autonomous stamen chromosomal inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1902 Hershel Mitchell uniparental inheritance
Cytoplasmic miteriance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are	circulonian internation cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal nucleus mitochondrial inheritance cytoplasmic inheritance extra diffect chromosomal Mendel 1904 nonmendelian Maternal maternal/inheritance structural/variation	extra chromosomal inheritance chromosomal genes non chromosomal genes inter chromosomal abnormal cytoplasm cytoplasm cytoplasm cytoplasm cytoplasm cheritance extramcelar inheritance inheritance fantorens 1907 Mendelian chromosomal cytoplasmic inheritance annéer variation	non chromosomal inheritance mitchoudrial genes extra chromosomal mitchoudrial genes extra chromosomal stable mitochoudrial genes extra nackear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance Ruth sager mendelian Ruth sager mendelian chromosomal inheritance chromosomal inheritance chromosomal inheritance tamaslation of gene	minicromata internance non chromosomali genes mitochondrial inheritance non chromosomali genes chromosomali autonomous stamen chromosomal inheritance both evtoplasmic and extramucle non endromosomal inheritance non endromosomal inheritance Herschel Mitchell 1902 Herschel Mitchell unigarental inheritance expression of gene
Cytoplasmic interfance also known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are is the study of how mitochondria are kentron on system of heredity Mitochondral inheritance also locom as Tati controlled by a gene of form of the progenys is called Evon metadelim inheritance find for organication in the progeny is called Evon metadelim inheritance find for organication in the structure of a reganication in the structure of an organication is termado Unignerated inheritance is a	cerestoriant nan-tance cytophamic genes chronocomal inheritance cytophamic genes inter chronocomal notice-homosonial notice-homosonial notice-homosonial notice-homosonial chronocomal Honda Hon	extra chromosonia inheritance chromosonia genes non chromosonia genes inter chromosonia genes abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cheritance extramuclear inheritance inheritance chromosonia mitochondria C9007 Mendeliana chromosonial cytoplasmic inheritance muber variation chromosonial inheritance	non chromosomal inheritance mitchondral genes extra chromosomal mitchondral genes extra chromosomal stable mitochondral genes extra nackear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance mendelian B908 Ruth suger mendelian chromosomal inheritance etyroplasmic cytoplasmic inheritance	nmixtromaria internance non chromosomi genes mitochondrial inheritance non chromosomali genes chromosomali antoennoos stamen bodi veroalsmic and extramacle novaissmic and extramacle nova mendelli i laheritance en mendelli i laheritance extra nuclear inheritance extra nuclear
Cytoplasme innertiance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are is the study of how mitchchoding ares known as interided matching are of the mother but expressed in the progeny is called Extranuclear inheritance is an Genor of inheritance Non medical in inheritance is an Genor of inheritance Transmission of genotypes from one parental type to all progeny is known as inheritance Transmission of genotypes from one parental type to all progeny is known as inheritance is an from one inheritance Transmission of genotypes from one parental type to all progeny is known as inheritance Transmission of genotypes from one parental type to all progeny is known as inheritance is an	circumonia interime cytoplasmi genes chromosomal inheritance cytoplasmi genes inter chromosomal mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance matemal effect chromosomal Mendel 1904 matemal inheritance structural variation matemal inheritance matemal inheritance	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia abnormal abnormal cytoplasmi chieritance extrancelar inheritance mitochondria Carl correns 1907 Mendelina Mendelina chromosonial chromosonial uniparenta inheritance number variation chromosonial uniparenta inheritance mental	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal sinheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance cytoplasmic inheritance cytoplasmic inheritance	minicronical internance mitochonical internance non chromosomali genes chromosomali autonomous stamen chromosomal inheritance both evroplasmic and extranucle non mendelian Hershel Mitchell 1002 Hershel Mitchell uniparental inheritance expression of gene extra nuclear intermediate inheritance phenotypic disease
Cytopame innertiance also known as Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are Genes located in cell organelle in the cytoplasm are is the study of how mitchcofthat are known as interiated study of how mitchcofthat are shown as some of heredity Larging escual reproduction childroplast chromosomes are only transmitted through the of the egg cell is the study of how mitchcofthat are har ender one system of heredity Mitchondrial inheritance fast discovered by	circumoniar internance cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal nucleus mitochodrali iheritance mitochodrali iheritance cytoplasmic inheritance matemal iffect chromosomal Mendel 1904 nonmendelian Matemal matemal iheritance structural variation matemal iheritance matemal iheritance genetic multiple mechanism	extra chromosomal inheritance chromosomal genes non chromosomal genes inter chromosomal abnormal cytoplasmi cytoplasmi inheritance extranuclear inheritance inheritance carl correns 1907 Mendelian chromosomal chromosomal chromosomal uniparentai inheritance number variation chromosomal uniparentai inheritance number variation chromosomal uniparentai inheritance number variation corrent of homosoma	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitochondria extra naclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance etranslation of gene cytoplasmic inheritance infactivus disease single mechanism	minicronomata intercement on chromosomal genes mitochondrial inheritance non chromosomal genes chromosomal autonomous stamen chromosomal inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1902 Hershel Mitchell 1902 extra nucleon display and the state inheritance expression of gene extra nucleon phenotypic disease vidi mechanism
Cytoplasme interfance laso known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochondria are known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochondria are known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochondria are known as Genes present in cytoplasm and mitochondria are known as some of heredity Danima bial rempt of thore mitochondria are stored with the cytoplasm are genes are compared with the cytoplasm are genes are compared with the cytoplasm are discovered are Genes present in cytoplasm are genes are compared with the cytoplasme are compared with the cytoplasme are Train controlled by a gene of the mother but cepressed in the progeny is called Extramulecin inheritance first discovered at Uniparential inheritance first discovered at Uniparential inheritance first discovered at Uniparential inheritance for discovered at Uniparential inheritance for discovered at Uniparential inheritance for discovered at 	Chromosome Indexianisme interventione elytopharmic press inter chromosomal normal normal encleus mitochromofai inheritance matemal effect chromosomal effect chromosomal Mendel 1904 nonmendelain Matemal matemalinheritance structural variation matemal inheritance structural variation matemal inheritance structural variation matemal inheritance genetic multiple mechanism Chromosome	extra chromosonia inheritance chromosonia genes non chromosonia genes inter chromosonia genes inter chromosonia genes adnormal cytophamo inheritance cytophamo inheritance inheritance mitochondra Carl correns Carl	non chromosomal inheritance mitchondral genes extra chromosomal mitchondral genes extra chromosomal sahle mitochondral genes extra chromosomal inheritance chromosomal inheritance chromosomal inheritance mendelian Ruth sager mendelan et al. Sager mendelan infectious disease single mechanism infra chromosome	nunctionata internance non chronosomia genes mitochondrial inheritance non chronosomial genes chronosomal antonomos antonomos both cvtoplasmi and extranucle non chronosomal inheritance both cvtoplasmi and extranucle non chronosomal inheritance non mendelian both cvtoplasmi and extranucle non chronosomal inheritance pom schollen Usopper school (statistication) Hershel Mitchell 1902 Hershel Mitchell 1902 Hershel Mitchell uniparental inheritance extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abnormal chronosomes
Cytoptame interfance also Arown as Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are Genes located in cell organelie in the cytoptam are is the study of how mitochondria ark betromoses are only transmitted through theof the egg cellis the study of how mitochondria genes are inherited Mitochondrial inheritance field socreted by Non medical inheritance field socreted by Non medical inheritance is a form of inheritance Transmission of genotypes from one paternal type to all progeny is known asinheritance Transmission of genotypes from one paternal type to all progeny is known asinheritance Transmission of genotypes form one homeropytes is known as The transmission of genotypes form one homeropytes is known as The transmission of genotypes form one paternal type to all progeny is known as	cerestonian international complanting upon compared and a second and a compared and a second a neckets compared and a second a compared and a second compared and a second compa	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi chromosomia dheritance extrameclear inheritance mitochondria Carl correnss 1907 Mendelian Mendelian dherotance cytoplasmi cinheritance manker variation chromosomial uniparental inheritance mental various mechanism extra diromosomes extra diromosomes variantalian tanget directiones and series directiones and various mechanism	non chromosomal inheritance mitchondrali genes extra chromosomal mitchondrali genes extra chromosomal stable mitchondrali genes extra nuckear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethoromosoma inheritance ethoromosoma inheritance ethoromosoma inheritance strafactoria disease single mechanismi intra chromosome geneticante in the strafactoria printi a chromosome geneticante in the strafactoria printi schoromosome geneticante in the strafactoria printi schoromosome in the sc	minicronouran internance mitochonical internance mitochonical internance chromosomal autonomous stamen ochromosomal interitance chromosomal interitance both evoplasmic and extranucle non chromosomal interitance mon chromosomal interitance mon chromosomal interitance mon chromosomal interitance prost chromosomal interitance thershel Mitchell 1902 Hershel Mitchell uniparental interitance expression of gene extra nuclear interitance photolyce disease volume deromosomes many dipteran files
Cytoptame interfance into Aroon as Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are Gene located in cell organelle in the cytoptam are is the study of how mitchchording ares known as intri controlled by agene of the mother but expressed in the progeny is called Extranuclear inheritance is an Genor of inheritance Non medicalin inheritance is an Genor of inheritance Transmission of genotypes from one parental type to all progeny is known as inheritance Transmission of genotypes from one parental type to all progeny is known as inheritance Transmission of genotypes from one parental type to all progeny is known as The transmission of genes that are located in mitchchordina is termed The transmission of genes that are located in mitchchordina is termed The transmission of genes that are located in mitchchordina is known as Photype of the heterozygne falls between that of the two homozygnes is known as Photype of the heterozygne falls between that of the two homozygnes is known as Photype of the heterozygne falls between that of the two homozygnes is known as	circumonian internance cytoplasmic genes chromosomal inheritance cytoplasmic genes inter chromosomal nucleus mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance matemal diffet chromosomal Mendel 1904 Mendel 1904 Matemal 1904 Matemal matemal inheritance structural variation matemal inheritance structural variation matemal inheritance penetic multiple mechanism Chromosome normal chromosome microbes salivary glands	extra chromosomal inheritance chromosomal genes non chromosomal genes inter chromosomal abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cheritance extranuclear inheritance mitochondria Carl correns 1907 Mendelian chromosomal cytoplasmi inheritance number variation chromosomal uniparental inheritance mental various mechanism extra chromosoms variant extra chromosoms variant throwid gland hundreds of dexoyribonucleic acid	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethoromosomal inheritance ethoromosomal inheritance single mechanism intra chromosomes gaint chromosome gaint chromosome plants wings	minicronouran internance mitochondrial inheritance non chromosomali genes mitochondrial inheritance chromosomali autonomous stamen chromosomal inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1902 Hershel Mitchell 1902 Hershel Mitchell uniparental inheritance expression of gene extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abnormal chromosomes abnormal chromosomes many dipteran flies thvroid gland bundred of ribonucleic acid
Cytoptame interfance also known as Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are Gene socated in cell organelle in the cytoptam are Genes located in cell organelle in the cytoptam are is the study of how mitchchorthar are known as initia controlled by a gene of the mother but expressed in the progeny is called Extranuclear inheritance frant discovered by	Chromosome informance development informance development information development information normal normal normal encleux matcenal effect ectoronosomal effect development information Mendel 1904 nonmendelain Maternal maternalinteritance structural variation maternal inferitance maternal inferitance solution solution of devorphilocular information more based of thousance maternal inferitance maternal inferitance maternal inferitance maternal inferitance maternal inferitance solution of devorphilocular information more based of thousance maternal information maternal information maternal information maternal inferitance solution of devorphilocular information maternal information maternal information maternal inferitance solution maternal inferitance sol	extra chromosomia inheritance chromosomia genes non chromosomia genes inter chromosomia abnormal cytoplasmi cytoplasmi cytoplasmi inheritance extranuclear inheritance extranuclear inheritance inheritance carl correns 1907 Mendelian chromosomial chromosomial chromosomial unigarental inheritance namber variation chromosomial various mechanism extra chromosomes variant thirotal eland handeles di doxyribonaucleic acid Rah sage	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitochondria extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance etytoplasmic inheritance etytoplasmic inheritance etytoplasmic ethoritance etytoplasmic inheritance gaint chromosomes gaint chromosome	nmix.tomatan internance pino chromosomi genes mito.chondrial inheritance pino chromosomi genes chromosomal anamen observer and server and server chromosomal inheritance both cvropaseni and extramucle non chromosomal inheritance pon mendelian and extramucle non chromosomal inheritance pon mendelian Hershel Mitchell 1902 Hershel Mitchell 1902 Hershel Mitchell 1902 Hershel Mitchell uniparental inheritance extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes andorral chromosomes andorral chromosomes many diperan files thyvoid glan handred of ribonucleic acid Carl ocross
Cytoptame interfance also known as Gene located in cell organelie in the cytoptam are Gene present in cytoptam and mitochordia are known as Chronoptass are seenetically and have their on system of heredity. During sexual reproduction chtoroptast chronosness are only transmitted through the of the egg cell is the study of how mitochordinal genes are inherited Mitochordial inheritance that down and as Tatic controlled by agene of the mother test epseed in the progeny is called Extramulear inheritance is an Non medician inheritance first discovered by [Inheritance is an Non medician inheritance first discovered by [Inheritance is an Inheritance is an [Inheritance is an [Inhe	cristionian international company and an end of the second chromosomal inheritance chromosomal inheritance inter chromosomal necleus company and the second company and the second company and the second company and the second company and the second maternal inheritance maternal since internation common company and the second company and the s	extra chromosonia inheritance chromosonia grees non chromosonia grees inter chromosonia grees inter chromosonia abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cheritance extrameclear inheritance inheritance anticologi 1907 Mendelian Mendelian Mendelian Mendelian Chromosonial chromosonial chromosonial chromosonial inheritance marker variation chromosonial market schemes extrameter inheritance marker variation chromosonial market schemes warks dromosomes extrant market schemes extrameter schemes market schemes throwich schemes market sc	non chromosomal inheritance mitchondral genes extra chromosomal mitchondral genes extra chromosomal stable mitchondral genes extra nuckear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance etranslation of gene cytophasmic cytophasmic stigler mechanismi inter deromosomes ingen etromosomes ingen etromosomes ingen etromosomes participationes partici	ministrometra internance mitochonical internance mitochonical internance chromosomal autonomous stamen on chromosomal internance both evoplasmic and extramacle non chromosomal internance both evoplasmic and extramacle non chromosomal internance mon chromosomal internance pron chromosomal internance mon chromosomal internance promosoma internance international internance promosoma internance expression of gene expression of gene
Cytoplasme interfance iso Arown as Genes located in expoplasm are	circumonia internance cytoplasmi genes chromosomal inheritance ecytoplasmi genes inter chromosomal nacleus mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance matemal effect chromosomal Mendel 1904 Mendel 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal Chromosomal Matemal Inheritance structural inheritance structural inheritance salivary danda Mendel 1881 Matemal	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi chromosomia cutorerss 1907 Mendelina Mendelina chromosomia cytoplasmic inheritance number variation chromosomia chromosomia uniparentai inheritance number variation chromosomia uniparentai inheritance number variation chromosomis extra chromosoms variants hvroid gland hundreds of dexoyrihomucleic acid Rath sager 1883	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal sabile mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethomosomal inheritance ethomosomal inheritance single mechanism intra chromosoms gaint chromosoms gaint chromosoms gaint chromosoms gaints of riboucleic acid Balbani 1880 nullisomy aneuploidy	minicrometar internance pino chromosomali genes mitochondrial inheritance non chromosomali genes chromosomali autonomous stamen ochromosomal inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1002 Hershel Mitchell 1002 Hershel Mitchell uniparental inheritance expression of gene extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abnormal chromosomes abnormal chromosomes many dipteran files thyrotid sland hundred of riboucleic acid Carl correns 1882 aneuploidy
Cytoplasme interfance laso known as Gene located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochoadria are known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochoadria are known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochoadria are known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm and mitochoadria are known as Genes located in cell organelie in the cytoplasm are Genes located in the cytoplasm are defined and located are Mitochoadrial interfaces losa known on as Trait controlled by a gene of the mother bet cepressed in the progeny is called Extramuleer interfaces losa known of in inferitance Mitochoadrial interfaces losa known of inferitance Tramsinsion of genes that are located in mitochoadria is termed	Chromosome induced and a solution of the solution of the solution of the solution of the solution neural and the solution of the solution neural ended and the solution of the solution of the solution neural effect chromosomal effect chromosomal effect chromosomal effect chromosomal and the solution matemal inheritance matemal inheritance solitance matemal inheritance solitance matemal inheritance solitance matemal inheritance solitance matemal inheritance solitance matemal inheritance solitance matemal inheritance solitance matemal inheritance solita	extra chromosonia inheritance chromosonia grees non chromosonia grees inter chromosonia grees inter chromosonia grees inter chromosonia grees cytoplane inheritance inheritance inheritance inheritance inheritance inheritance inheritance inheritance inheritance inheritance inheritance inheritance inheritance mittochondia Carl correas Carl coreas Carl correas Carl corre	non chromosomal inheritance mitchondral genes extra chromosomal mitchondral genes extra chromosomal stable mitochondral genes extra chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal inheritance ethoronosomal stable and ethoronosoma infactoronosomes gaint chromosomes plants thousands of nboucleic acid Babbani 1880 naneiploidy anceploidy	nmixtonian interinance pino chronosomia genes mitochondrial inheritance en ochronosomial genes chronosomal antonomos antonomos both cvtoplasmi and extramucle non chronosomal inheritance both cvtoplasmi and extramucle non chronosomal inheritance pino mendelian definitatione extra nuclear inheritance extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abnormal chronosomes abnormal chronosomes handred of ribonucleic acid Carl correns 1882 aneuploidy multisony multisony multisony multisony
Cytoplasme interfance into Aroon as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are is the study of how mitochondria are therrown system of heredity During sexual reproduction chloroplast chromosomes are only transmitted through the of the egg cell is the study of how mitochondria genes are inherited Mitochondrial inheritance fish Genoverod by Mitochondrial inheritance fish Genoverod by Mitochondrial inheritance fish Genoverod by Transmission of genes that are located in introchondria is termed is the study initia in the structure of an organity is chromosome Fiberacytor of the hereory operated list view to that of the two homozygets is is known as Genesurge common in Mitochondria as termed Mosaichin as	cristolomia nariance cytoplamic genes chronosomal inheritance chronosomal inheritance inter chronosomal naticular naticular naticular naticular naticular naticular composition of the second maternal effect chronosomal for the maternal inheritance maternal inheritance sinturular variation maternal inheritance sinturular variation maternal inheritance sinturular variation maternal inheritance sinturular variation maternal inheritance sinturular variation maternal inheritance sinturular sinturular sinturular maternal inheritance sinturular sinturu	extra chromosonia inheritance chromosonia grees non chromosonia grees inter chromosonia grees inter chromosonia grees inter chromosonia grees cytoplasmi cytoplasmi choritance extramoclear inheritance extramoclear inheritance chromosonia 1907 Carl correns 1907 Carl correns 1907 Chromosonia Chro	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitochondrial genes extra nackear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance extra nackear inheritance extra nackear inheritance extra nackear inheritance cytoplasmic cytoplasmic inheritance extra nackear inheritance single mechanismi intra chromosomes plants thousands of ribonucleic acid Babbiani 1880 nancuploidy aneuploidy aneuploidy	nunctionata internance non chromosomali genes mitochondrial inheritance chromosomali autonomos stamen ochromosomali inheritance bosh evoplasmic and extramacle non chromosomali linheritance bosh evoplasmic and extramacle non chromosomali linheritance bosh evoplasmic and extramacle non chromosomali linheritance bosh evoplasmic and extramacle and chromosomali linheritance terramacle and the state state and the state state state and the state s
Cytoplasme innertiance also known as Genes located in exploylasm are	circusonia internance cytoplasmi genesic chromosomal inheritance expoplasmi genes inter chromosomal nacleus mitochondrial inheritance mitochondrial inheritance matemal effect chromosomal Mendel 1904 Mendel 1904 Matemal 1904 Matemal 1904 matemalinheritance structural inheritance structural inheritance matemalinheritance structural inheritance matemalinheritance structural inheritance structural inheritance statistica (document) matemal inheritance structural inheritance structural inheritance statistica (document) matemal inheritance structural i	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess extrancelear inheritance extrancelear inheritance mitochondria Carl correnss 1907 Mendelian Mendelian chromosomia extra chromosomia extra chromosomis entral inheritance mental various mechanism entral various mechanism entral strat chromosomis entral for decoyribonucleic acid Ruth sager 1883 eaploidy eapl	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal sable mitochondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance single mechanism intra chromosoms gaint chromosoms gaint chromosoms gaint chromosoms gaint chromosoms gaint spanson housands of niboucleic acid Balbani 1880 nullisomy aneuploidy aneuploidy aneuploidy trisomy thisomy	minicromotaria interfance pino chromosomali genes mitochondrial inheritance non chromosomali genes chromosomali autonomous stamen ochromosomal inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1002 Hershel Mitchell 1002 Hershel Mitchell uniparental inheritance expression of gene extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abnormal chromosomes abnordal chromosomes abnordal chromosomes abnordal chromosomes many dipteran files thyvid sland hundred of riboucleic acid Carl correns 1882 aneuploidy nullisomy nullisomy tetrasomy
Cytoptame interfance into Aroon as Gene located in edigrandle in the cytoptam are	Chromosome informance dyrupaming genes dyrupaming genes inter chromosomal informance cytoplasmic informance matchess mitochorodial infortiance vytoplasmic informance matemal effect chromosomal effect chromosomal effect dyrupamic informance matemal informance matematical informance	extra chromosomia inheritance chromosomia genes non chromosomia genes inter chromosomia abnormal cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi choritance extranuclear inheritance inheritance carl correns 1907 Mendelian chromosomial cytoplasmi inheritance number variation chromosomial cytoplasmi inheritance mental various mechanism ettra. chromosoms variant mental thvorid gland hundreds of deoxythomucleic acid Ruth suger cuplokdy cuplokd	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance cytoplasmic inheritance single mechanism intra chromosomes gaint chromosomes gaint chromosomes gaint chromosomes gaint chromosomes gaint chromosomes gaint space plants wings husans ingle mechanism ittra chromosomes gaint chromosomes gaint chromosomes gaint space gaint genes source and space gaint genes gaint genes ga	Innocementari interviewe mitochonical interviewe on chromosomial genes mitochonical interviewe chromosomal autoen on chromosomal genes chromosomal interviewe beth cvroplasmi and extramucle non chromosomal interviewe beth cvroplasmi and extramucle non chromosomal interviewe beth cvroplasmi and extramucle non mendelian der der der der der der Herstel Mitchell 1902 Herstel Mitchell 1902 Herstel Mitchell 1902 Herstel Mitchell unigarental interritate interviewe extra nuclear intermodiate interviewe phenotypic disease void mechanism mutated chromosomes abnormal chromosomes abnormal chromosomes many diperan files thyvoid gelan handted of ritonucleic acid Carl corress actification lastice strasomy tetrasomy mullisomy mullisomy mullisomy mullisomy mullisomy mullisomy mullisomy mullisomy mullisomy mullisomy
Cytoplasme interfance laso known as Genes located in cell organelie in the cytoplasm are Genes located in cell organelie in the cytoplasm are is the study of how mitochondria genes are inherited Mini controlled by the cell of the cytoplasm are in the study of how mitochondria genes are inherited Mini controlled by the cell of the cytoplasm are inheritance is a form of inheritance Non mendelian inheritance first discovered by inheritance is a form of inheritance The transmission of genes that are located in nitochondria is termed is the variation in the structure of an organis's dormonous The pattern of mtDNA inheritance is a form of inheritance The pattern of mtDNA inheritance is a form of the two homozygets is known as Beneatype of the herotrozyget falls between that of the two homozygets is known as disease are suppected to be caused by structural variation Mossicitum as by the chromosomes are Polytene chromosomes are Polytene chromosomes are Polytene chromosomes are commonin [ Polytene chromosomes are commonin [ Polytene chromosomes are commonin [ Polytene chromosomes are to chromosomes are known as	cristonian internet cytophonic grant chronobarnal inheritance chronobarnal inheritance inter chronosomal inter chronosomal matchess monophololy monopholo	extra chromosonia inheritance chromosonia grees non chromosonia grees inter chromosonia grees inter chromosonia grees inter chromosonia grees cytoplasmi cytoplasmi cytoplasmi chromosonia cytoplasmi chromosonia Carl correns Carl correns Car	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitochondria genes extra anckear inheritance chromosomal inheritance chromosomal inheritance extra anckear inheritance extra anckear inheritance extra anckear inheritance erosomal inheritance infectious disease single mechanism intra chromosome plants thousands of ribonucleic acid Babbiani 1880 naneuploidy aneuploidy aneuploidy aneuploidy erosom trisomy trisomy trisomy trisomy 2N + 2 2N + 1	nmix.romoutum internance in ordromosomali genes mitochondrial inheritance inon dromosomali genes chromosomali autonomous stamen obche votoalauria met eta hort ordromosomali inheritance bron chromosomali inheritance pron mendelian Hershel Mitchell 1902 Hershel Mitchell 1902 Hershel Mitchell 1902 Hershel Mitchell uniparental inheritance extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abnormal chromosomes many dipteran files thvoid genes lass2 aneuploidy nullisomy nullisomy tetrasomy tetrasomy tetrasomy andisony 2N + 1 2N + 2
Cytoptame interfance also known as Genes located in cell organelie in the cytoptam are	cristolomia interative cytoplamic genes chromosomal inheritance cytoplamic genes inter chromosomal mechanics cytoplamic inheritance mitochondrial inheritance mitochondrial inheritance matemal effect chromosomal Matemal 1904 Ma	extra chromosonia inheritance chromosonia genes non chromosonia genes inter chromosonia genes inter chromosonia genes cytoplasmi cytoplasmi cytoplasmi cytoplasmi cytoplasmi chromosonia low mitochondria Carl corres low chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia mental markets of deoxyribonucleic acid Ruh suge lissz euplokity ceplokity	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuckar inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance infactions disease single metadaian disctions disease single metadaians intra chromosome plants wittes Babbuni Bab	mukcronuctan interinance inno chromosomali genes mitochondrial inheritance inno chromosomali genes chromosomali autonomous stamen no chromosomali inheritance bosh evtoplasmic and extranucle non echromosomali inheritance non echromosomali inheritance inno chromosomali inheritance inno chromosomali inheritance expression of gene extra nuclear linetradulate inheritance phenotypic disease void mechanism mattade chromosomes ahnormal chromosomes ahnormal chromosomes ahnortan files thvroid gland landred of ribourcleic acid Carl correns 1882 aneuploidy nullisomy nullisomy nullisomy tetrasmy intimestican chromosomes andisomy nullisomy nullisomy nullisomy nullisomy zN = 1 ZN = 1
Cytoptame interfance info Aroon as Gene Societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are Gene societ of cell cignalle in the cytoptam are is the study of how mitchondria are known as Chicoptastar are cell cignalle in the cytoptam are is the study of how mitchondria genes are inherited Mitchondrial inheritance fill as Genover dot the cytoptam are inheritance is an Geno of inheritance Non meddelin inheritance is an Genor of inheritance Transmission of genotypes from one parental type to all progeny is called Transmission of genotypes from one parental type to all progeny is known as The transmission of genotypes from one parental type to all progeny is known as The transmission of genotypes from one parental type to all progeny is known as Photype of the heterozygne fills between that of the two homozygots is known as Photype of the heterozygne fills between that of the two homozygots is known as Photypee of the heterozygne fills between that of the two homozygots is known as Photypee fromsomes are common in Photypee chromsomes are consend by the presence of a Photypee chromsomes are consend by the presence of a Photypee chromsomes are consend by the commonsomes of a normal set of chromsome are missing or present and changes in chromsome number is the absormal combination of transomes. The Loss of single chromsomes is known as	circumonia internance cytoplasmic genes chromosomal inheritance merital nacleus mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance matemal effect chromosomal Mendel 1904 Mendel 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal Chromosomal Matemal Inheritance surcutural viriation matemal inheritance surcutural viriation matemal inheritance surcutural viriation matemal inheritance surcutural viriation matemal inheritance surcutural viriation matemal inheritance surcutural viriation matemal inheritance surcutural decostration matemal inheritance surcutural viriation matemal viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriation viriat	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess extrancelar inheritance extrancelar inheritance inheritance extrancelar inheritance inheritance carl correnss 1907 Mendelan chromosonial chromosonial chromosonial extra chromosonias uniparentai inheritance mentai various mechanism extra chromosonis extra chromo	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance cytoplasmic inheritance infactious disease single mechanism intra chromosomes gaint chromosomes gaint chromosomes gaint chromosome plants wings husas Mongalianti Babbini Babbini Babbini Babbini Babbini Babbini Composition Babbini Composition Compos	mukchonical interlance inno chromosomaligenes mitochondrial inheritance inno chromosomaligenes chromosomaligenes chromosomaligenes chromosomaligenes chromosomaligenes both evtoplasmic and extranucle non chromosomali inheritance both evtoplasmic and extranucle non chromosomali inheritance inno chromosomali inheritance inno chromosomali inheritance inno chromosomali inheritance area chromosoma inheritance phenotypic disease vid mechanism phenotypic disease vid mechanism mark disease vid mechanism mark disease intermediate inheritance phenotypic disease vid mechanism mark disease vid mechanism mark disease vid mechanism mark disease vid mechanism multatel chromosomes ahnormal chromosomes ahnormal chromosomes mary dipteran files thyroid gland hundred of ribotucleic acid Carl correns 1882 aneuploidy nullisomy nullisomy nullisomy zN = 1 ZN = 1 ZN = 1 ZN = 1 ZN = 1
Cytoplasme interfance info Aroon as Genes located in cell cognadie in the cytoplasm are	cristonian internet cytophoneii generate cytophoneii generate cytophoneii generate inter chronosomal inter chronosomal matcleus matcraul effect cytophasnic interinance matcraul effect chronosomal effect chronosomal effect chronosomal effect matcraul effect chronosomal interinance matcraul interinance m	extra chromosonia inheritance chromosonia grees non chromosonia grees inter chromosonia grees inter chromosonia grees inter chromosonia grees cytoplasmi cytoplasmi cytoplasmi chromosonia cytoplasmi chromosonia Carl correns Carl correns Car	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitochondria genes extra anckear inheritance echromosomal inheritance chromosomal inheritance echromosomal inheritance echromosomal inheritance ethonosomal inheritance infectious disease single mechanism intra chromosome plants thousands of ribonucleic acid Babbiani 1880 naneuploidy aneuploidy aneuploidy trisomy	nnucronouran internance in ordronosomia genes mitochonolaria inheritance ino chronosomali genes chromosomali autonomous stame obcho civoplaumia mad extranacce chromosomal inheritance choli civoplaumia mad extranacle non chromosomal inheritance non mendelian deritance extra nuclear inheritance extra nuclear intermediate inheritance phenotypic disease vidi mechanism mutated chromosomes abormal chromosomes many diperan files thivoid genes lass2 aneuploidy nullisomy nullisomy nullisomy nullisomy tetrasomy tetrasomy nullisomy zN - 1 ZN - 1 ZN - 1 monosomic mouse obcomic mouse obcomis mouse obcomis mous
Cytoptame interfance info Aroon as Gene Societ on cell organelie in the cytoptam are	cristionian international company of the second sec	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi chieritance extrameclear inheritance inheritance and concess 1907 Mendelian chromosonia inheritance munker variation chromosonia inheritance munker variation curve and there and the second wariant animala thurveid ella chromosones variant sequeloidy euploidy euploidy euploidy euploidy euploidy euploidy euploidy monosony tetrasonic 2 N + 1 2 N + 1 double tetrasonic 2 N + 2 + 2 2 N + 2 2	non chromosomal inheritance mitichondrial genes extra chromosomal mitichondrial genes extra chromosomal stable mitochondrial genes extra nuckear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance etranalconi gene cytophamic invertiance chromosomal inheritance etranalconi gene cytophamic invertiance chromosoma inheritance chromosoma inheritance chromosoma inheritance etranalconi gene cytophamic invertiance etranalconi gene cytophamic invertiance chromosoms gaint chromosomes gaint chromosomes platts howsmash of fibonucleic acid Babhani 1880 nancupicidy anceptoidy trisomy triso	minicronous internance in dromosomi genes mitochondrial inheritance inon dromosomali genes chromosomal inheritance hornosomal inheritance both evoplasmic and extramcle non chromosomal inheritance men methal in the strands of the strands of the strands of the strands of the strands of the strands methanisme even such as a strands of the strands of the the strands of the strands of the strands of the strands of the strands of the strands of the strands of the strands of th
Cytoplasme interfance iso Aroon as Genes located in expoplasm are	circumonia international cytoplasmi genes chromosomal inheritance cytoplasmi genes inter chromosomal mackus mitochondrial inheritance mitochondrial inheritance mitochondrial inheritance matemal effect chromosomal Mendel 1904 Mendel 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal 1904 Matemal Chromosoma matemal inheritance structural variation matemal inheritance structural variation monsploidy monosploidy	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess extrancelar inheritance extrancelar inheritance inheritance extrancelar inheritance inheritance carl correnss 1907 Mendelan chromosomia cytoplasmic inheritance number variation chromosomia extra chromosoms variant setta chromosoms variants thvoid gland hundreds of doxyribonucleic acid Ruth sager 1883 euploidy eupl	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance infractious disease single mechanism intra chromosoms gaint chromosoms gaint chromosoms gaint chromosoms gaint chromosoms gaint achromosoms gaint achromosoms gaint achromosoms gaint achromosoms gaint achromosoms gaint achromosoms gaint chromosoms gaint achromosoms gaint achro	minicronous internance mitochonical internance in chromosomali genes chromosomali genes chromosomali autonomos stamen ochromosomali inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1002 Hershel Mitchell 1002 Hershel Mitchell 1002 extra nuclear intermediate inheritance phenotytic disease void mechanism mattade chromosomes abnormal chromosomes abnormal chromosomes abnorder filos thytoid sland hundred of riboucleic acid Carl correns 1882 aneuploidy nullisomy nullisomy nullisomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy mullisomy tetrasomy nullisomy nullisomy tetrasomy
Cytoplane interfance info Aroon as Genes located in cell copation are	terrshormin internations chypopune insertance chypopune insertance chypopune insertance chypopune insertance markets matches matches matches matches matches terrchestance matchest mat	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi cytoplasmi chromosonia cytoplasmi miticchrondia miticchrondia miticchrondia miticchrondia miticchrondia cytoplasmic inheritance momber variation chromosomal cytoplasmic inheritance momber variation chromosomal chromosomal multication chromosomal multication chromosomal multication chromosomal momber variation chromosomal multication chromosomal momber variation momber variation chromosomal motification momber variation chromosomal momber variation capitoli variant momber variation capitoli variant variant	non chromosonial inheritance mitichondrial genes extra chromosonial mitichondrial genes extra chromosonial stable mitochondria genes extra nackear linkeritance extra nackear linkeritance echromosonial inheritance chromosonial inheritance echromosonial inheritance erromosonial inheritance etromosonial inheritance etromosonia inheritance etromosonia inheritance etromosonia inheritance etromosonia inheritance etromosonia single mechanism intra chromosonie giant chromosonie giant chromosonie giant chromosonie plants thousands of ribonucleic acid Babbiani 1880 naneuploidy aneuploidy aneuploidy aneuploidy aneuploidy 2N + 2 2N + 2N	nmix.rom/an internance in or dromosomali genes mito.chondrial inheritance in or dromosomali genes chromosomali autonomous stamen on dromosomali martinance to obstationance obstationance obstationance obstationance obstationance non mendelian distributionanti distributionanti distributionanti meritance extra nuclear inheritance extra nuclear inheritance extra nuclear inheritance extra nuclear inheritance extra nuclear inheritance extra nuclear inheritance extra nuclear inheritance extra nuclear inheritance distribution biomale diromosomes abornal chromosomes abornal chromosomes anary diperan files thyvid genes lasse aneuploidy nullisomy nullisomy ettrasomy attrasomy ettrasomy attrasomy ettrasomy at
Cytoplane interfance info Aroon is Cytoplane inter-Change and mitochodria are known as Gene shocad in call cignaphic in the cytoplasm are	ciristonia nancana cirphannii genes chronosomal inheritance chronosomal inheritance scytoplannii genes inter chronosomal natechronosomal natechronosomal effect versonosomal for the state scheme scheme scheme scheme maternal effect chronosomal effect maternal affect chronosomal maternal inheritance sinturula variation maternal inheritance silvaru glands thousands of deoxyrihomucleic acid Mendel 1881 monoploidy mono	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi chieritance extramuclear inheritance inheritance cari correnss 1907 dendelian chromosonia chromosonial cuplicity cup	non chromosomal inheritance mitichondrial genes extra chromosomal mitichondrial genes extra chromosomal stable mitochondrial genes extra nuckear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance trobalistical of genes mendelian factoria inheritance trobalistical genes infectiona disease single mechanismi infra chromosomes gaint chromosomes plants infra chromosomes p	minicronous internance indicational internance in chronosomal ignes chromosomal ignes chromosomal inheritance bosh evoplasmic and extramcle non chromosomal inheritance bosh evoplasmic and extramcle non chromosomal inheritance merchankania inheritance informational inheritance information information inheritance extra nuclear intermediate inheritance phenotypic disease vorta nuclear intermediate inheritance intermediate inheritance inte
Cytoptame interfance also Atown as Genes hourse of corporation are	circumonia international company of the second seco	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess extra clear inheritance extra nuclear inheritance inheritance carl correnss 1907 Carl correns 1907 Mendelina chromosomia chromosomia chromosomia chromosomia chromosomia ental various mechanism ental various mechanism ental varias mechanism ental start chromosomis ental inheritance mental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism ental varias mechanism exploidy euploidy euploidy euploidy etrasomic 2N + 1 2N	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance infractious disease single mechanism intra chromosoms gaint chrom	munctionata interlance pin chromosomali genes mitochondrial inheritance inon chromosomali genes chromosomali autonomos stamen ochromosomali inheritance both evtoplasmic and extranucle non mendelian Hershel Mitchell 1002 Hershel Mitchell 1002 Hershel Mitchell 1002 extra nuclear intermödiate inheritance phenotypic disease void mechanism matade chromosomes abnormal chromosomes abnormal chromosomes abnormal chromosomes abnormal chromosomes many dipteran files thyroid gland handred of riboaucleic acid Carl correns 1882 aneuploidy mullisomy mullisomy mullisomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy mulli
Cytoplane interfance info Gene located in cell equalie in the cytoplasm are	terrsponseries chyropaenia genes chyropaenia genes chyropaenia genes chyropaenia genes matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus matcelsus Mendel 1904 nonmendelainn Matemal Matemal matemal inheritance sutcurul variation matemal inheritance sutcurul variation matematicul sutcurul variation sutcurul variation sutcur	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplanni chineritance cytoplanni chineritance ensistetti chineritance ensistetti chineritance minicchondia minicchondia Gari correas 1907 Mendelian chromosonia cytoplanni chineritance number variaton chromosonia chromosonia chromosonia ettra chromosoness variaus entra chromosoness variaus ettra chromosoness variaus thevioti gland hundreds of doxyribonucleic acid Ruch sager euploidy	non chromosomal inheritance mitichondrial genes extra chromosomal mitichondrial genes extra chromosomal stable mitochondria genes extra nackari labetimace extra nackari labetimace chromosomal inheritance extra nackari labetimace extra nackari labetimace extra nackari labetimace extra nackari labetimace extra nackari labetimace extra nackari labetimace extra nackari diperimace extra nackari diperimace infectiona diperimace infectiona diperimace infectiona diperimace extra nackari diperimace infectiona diperimace extra nackari diperimace plants mitra diperimace plants infra diperimace extra nackari mitra diperimace plants infra diperimace extra nackari mitra diperimace plants infra diperimace extra nackari mitra diperimace extra nackari mitra diperimace extra nackari monaticism obsti dipat dark chromosomal narrangenents	munctionara internance innorthomous internance in chromosomali genes mitochonolial inheritance innorthomosomali genes chromosomali inheritance both cvtoplaama and extranucle non chromosomali anheritance both cvtoplaama and extranucle non mendelian distributed inheritance extra nuclear inheritance extra nuclear intermediate inheritance phenotypic disease void mechanism mutated chromosomes abornal chromosomes abornal chromosomes honoral defineosomes abornal chromosomes abornal chromosomes intermediate inheritance phenotypic disease void mechanism mutated chromosomes abornal chromosomes abornal chromosomes anary dipteran files thyvid genes lass2 aneuploidy nullisomy tetrasomy aulisomy tetras
Cytoplane interfance info Aroon is Gene located in cell cognable in the cytoplasm are	ciristonia nanchanic cyriphanic genes chronosomal inheritance chronosomal inheritance sinter chronosomal inter chronosomal matechanical inheritance stracture maternal effect chronosomal ford Mendel 1904 nonneeldalian Maternal maternal inheritance structural variation maternal inheritance salivary alans thousanso of deoxyribonucleic acid Mendel 1881 monoploidy mono	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi inheritance extranuclear inheritance inheritance and construction of the construction inheritance and construction of the construction of the construction of the construction of the construction of the construction construction of the construction of the construction of the construction of the construction of the construction of the construction of the construction of the construction of the construction of the construction of the construction of the constru	non chromosomal inheritance mitichondrial genes extra chromosomal mitichondrial genes extra chromosomal stable mitochondria genes extra nackear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance thromosomal inheritance thromosomal inheritance thromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomal inheritance tromosomes gaint chromosomes gaint chromosomes plants thousands of ribonucleic acid Babbiani 1880 nancuploidy aneuploidy trisomy t	minicronous internance indications and genes mitochonois and inferitance inon chronosomal genes chromosomal inheritance bosh evoplasmic and extramacle non-thronois and inheritance bosh evoplasmic and extramacle non-thronois inheritance bosh evoplasmic and extramacle non-thronois inheritance bosh evoplasmic and extramacle are chromosomal inheritance percession of gene extra nuclear intermediate inheritance phenotypic disease void mechanism mututed chromosomes anary dipteran files throvid selan bandred of ribonucleic acid Carl correns 1882 aneuploidy nullisomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy tetrasomy andilisomy 2N - 1 2N - 1 2N - 1 monosomic 2N + 1 2N - 1 monosomic 2N + 1 2N - 1 monosomic 2N + 1 2N - 1 monosomic and the three options 1940 all the three options 1940
Cytoptame interfance iso Jown is Gene located in cell cognale in the cytoptam are	circulonian international circulation of the second of the	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi ciheritance extrancelor inheritance mitochondria Carl correns 1907 Carl correns 1907 Mendelian denotonia chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonic chr	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuckar inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance infactions disease single metadation of gene cytoplasmic inheritance infactions disease single metadation functions disease single metadation functions disease single metadation functions disease single metadation intra chromosomes plants wittes mata chromosomes plants wittes anceptoidy anceptoidy anceptoidy anceptoidy anceptoidy 2N + 1 2N + 2 2N + 2 2N + 1 2N + 2 2N + 1 2N + 1 2N + 1 2N + 2 2N + 1 2N + 1 2N + 2 2N + 1 2N + 2 2N + 1 2N + 2 2N + 1 2N + 2 2N + 1 2N + 3 2N + 1 2N + 1 2N + 1 2N + 3 2N + 1 2N + 1 2N + 1 2N + 3 2N + 1 2N + 1 2N + 3 2N + 1 2N + 1 2N + 2 2N + 1 2N + 1 2N + 1 2N + 3 2N + 1 2N + 1 2N + 1 2N + 3 2N + 1 2N + 1 2N + 3 2N + 1 2N + 3 2N + 1 2N + 3 2N + 1 2N + 3 2N + 1 2N + 3 2N + 3 2N + 1 2N + 3 2N + 3 2	minicronous intercemponent genes mitochonical interfiance ino chromosomal genes chromosomal genes chromosomal interfiance both evolasmic and extranucle non chromosomal interfiance tersite Mitchell 1902 Hershel Mitchell uniparental intermédiate interfiance phenotypic disease void mechanism mutated chromosomes ahnormal chromosomes ahnormal chromosomes mutated chromosomes anormal chromosomes mutated chromosomes anormal chromosomes mutated chromosomes mutated chromosomes mutated chromosomes anormal chromosomes mutated chromosomes mutated chromosomes mutated chromosomes mutated chromosomes mutated chromosomes mutated chromosomes anormal chromosomes mutated chromosomes
Cytoplane interfance iso Aroon as Genes located or calculated in the cytoplasm are	terrsponse in international in	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess cytoplanni chineritance extra chromosomia cytoplanni chineritance enhoritance mitticchendria Gari correas unitochendria Gari correas indicchendria Gari correas indicchendria etta chromosoma indicchendria etta chromosoma etta chromosoma etta chromosomes variant etta chromosomes variant etta chromosomes variant animals thvovid eland hundreds of deoxyrbonucleic acid Ruth sager euploidy e	non chromosomal inheritance mitchondria [genes extra chromosomal mitchondria] genes extra chromosomal stable mitochondria [genes extra ackear inheritance extra ackear inheritance chromosomal inheritance chromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal stable and erromosomes gaint chromosomes gaint chromosomes gaint chromosome plants thousands of ribonucleic acid Babbiani 1880 naneuploidy aneuploidy aneuploidy aneuploidy aneuploidy aneuploidy aneuploidy aneuploidy errosomic erosomic errosomic errosomic errosomic errosomic erosomic errosomic	minicipantia interview in chromosomia genes mitochondria i inheritance in chromosomali genes chromosomali inheritance istore both cvtoplasmi and extranucle both cvtoplasmi and extranucle pron chromosomal inheritance pron meddelian distribution and extranucle pron meddelian distribution on meddelian distribution dist
Cytoplane interfance info Gene located in cell cognated in the cytoplasm are Gene present in cytoplasm and mitochodria are known as Gene present in cytoplasm and mitochodria are known as Gene present in cytoplasm and mitochodria are known as Choroplass are sentedialV and have their on system of heredity During sexual reproduction chloroplast chromosomes are only transmitted through the of the egg cell is the study of how mitochodrial genes are inherited Mitochodrial inheritance taks form of here in the progeny is called Extranuclear inheritance is a form of here inherited Mitochodrial inheritance taks form of here inherited Mitochodrial inheritance taks form of here inherited Mitochodrial inheritance fragments in the strongeny is called Extranuclear inheritance fragments in the strongeny is known as inheritance The strongeny form one presental type to all progeny is known as is the variation in the structure of an organism's dormonome is the variation in the structure of an organism's dormonome disease are supported from one present of a two homozygets is known as Structure in the structure of an organism's dormonome structure in all molecohorities is known as 	circumoname international internation of the second	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi chromatomic cytoplasmi chromatomic hiberitance inheritance extra notice inheritance inheritance chromosonia diverso somal cytoplasmi inheritance chromosonial cytoplasmi inheritance chromosonial motellistic cuploidy cup	non chromosomal inheritance mitichondrial genes extra chromosomal mitichondrial genes extra chromosomal stable mitochondria genes extra nackear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance extra nackear inheritance infections disease single mechanism ittra extra nackear inheritance plasts ittra nackear inheritance extra nackear inheritance infections disease single mechanism ittra nackear inheritance extra nackear inheritance infections disease inheritance infections disease infections disease	minicipandia internance in chronosomali genes mitochonosiani internance chromosomali internance stamen inchronosomali internance bodi vironismic and extramale no chronosomali internance bodi vironismic and extramale no mendellin linterinance extra nuclear internanciate internance operations extra nuclear internediate internance operations phenotypic disease viron ancelear internediate internance operations phenotypic disease vironi ancelear internediate internance operations phenotypic disease viron ancelear internation nutison nutison vironi second 2N + 1 2N + 1 monosomic color intuition single cell interosis color al the three options 1940 So
Cytoplane interfance iso Jown is Gene bocaed in cell cognable in the cytoplana are	circumona nanca nance cycipalamic genes chrononomal inheritance chrononomal inheritance sermal nacleus cycipalamic genes cycipalamic genes cyclipalamic genes c	extra chromosonia inheritance chromosonia gress non chromosonia gress inter chromosonia gress inter chromosonia gress inter chromosonia gress cytoplasmi citeritance extrancelor inheritance inheritance anticological inheritance mitochondria Carl correns 1907 Mendelian chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonia chromosonic chromoson	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuckar inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethormosomal inheritance ethormosomal inheritance translation of gene cytoplasmic inheritance intra chromosoms gaine chromosoma intra chromosoms gaine thromosoms gaine thromos	minicipandia internance inno chromosomi genes mitochondria inheritance inno chromosomi genes chromosomi inheritance inno chromosomi inheritance inno chromosomi inheritance bosh evoplasmic and extrancile non endenin inheritance bosh evoplasmic and extrancile non endenin inheritance internedian Hershel Mitchell uniparental inheritance phenotypic diease void mechanism phenotypic diease phenotypic diease void mechanism phenotypic diease phenoty
Cytoplane interfance iso Aroon as Genes koard in cell organite in the cytoplasm are	terrspinner inkninger terrspinner chytiphamie green chytiphamie green inkelses mitochomosinal nermal nermal nermal nermal nermal nermal nermal nermal nermal nermal nermal ffer terrspinner maternal inferitance maternal inferitance maternal inferitance structural variation maternal inferitance structural variation maternal inferitance genetic maternal inferitance structural variation maternal inferitance genetic maternal inferitance silvary almós thousands of deoxyribonucleic acid thousands of deoxyribonucleic acid denblemonosomic doublemonosomic d	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess cytoplanni chineritance extra chromosomia cytoplanni chineritance enhoritance enhoritance enhoritance indicchendia Gari correas 1907 Mendelian deromosomal eritochendia moder variant enhoritance momber variant ectro alternation etro antomosome etro antomosome etro antomosome variant momber variant etro antomosome variant etro antomosome etro antomosome variant etro antomosome etro antomosome variant etro antomosome etro antomosome variant moto etro antomosome variant numeter de antomosome etro antomosome etro antomosome variant moto etro antomosome variant moto etro antomosome variant enploidy eup	non chromosomal inheritance mitchondria genes extra chromosomal mitchondria genes extra chromosomal stable mitochondria genes extra chromosomal extra nackari inheritance extra nackari inheritance erbremosomal inheritance chromosomal inheritance erbremosomal inheritance erbremosomal inheritance erbremosomal inheritance erronalistic inheritance erotopismic cytopismic eytopismic inheritance infectious disease single mechanism intra chromosome gants single mechanism intra chromosome plants thousands of ribonucleic acid Babbani 1880 naneupiody aneupiody trisomy tri	minicipandia interfance incichanosania genes mitochondria inheritance incichanosania genes chromosomal inheritance ben chronosomal inheritance ben chronosomal inheritance ben chronosoma inheritance ben chronosoma inheritance in methelian and extranucle non chronosoma inheritance pon medelian distribution and extranucle non methelian distribution and extranucle non methelian distribution distr
Cyptemic interfance iso Aroon iso Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the cyptemic and Gene located in cell organedic in the progeny is called Extramedicate influentance fragments Gene located in cell on mitochondria is termed Gene medication in the structure of an organity is chromosome The pattern of mDNA indextance is the two bornoxygets is known as Gene structure of an organity is chromosome and Gene located in an located in mitochondria is termed Gene chromosome serve and Gene structure of an organity is chromosome and Gene structure and the composition of a gene of an organity is chromosome and Gene structure of an organity is chromosome and Gene structure and and the composition of a gene of an organity is chromosome and Gene structure and the composition of an organity is chromosome and Gene structure and and the composition of an organity is chromosome and Gene structure and and the composition of an organity is chromosome and Gene structure and and the compositient and the compos	circusonam numerance cycipalamic genes chronopounal inhoritance chronopounal inhoritance interchemican interchemican conservation interchemican interchemican maternal effect chromosomal effect chromosomal effect chromosomal effect chromosomal effect maternal inheritance structural variation maternal inheritance structural variation maternal inheritance structural variation maternal inheritance structural variation maternal inheritance structural variation maternal inheritance structural variation maternal inheritance maternal inheritance structural variation maternal inheritance structural variation monopolisity mono	extra chromosonia inheritance chromosonia greess non chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess inter chromosonia greess cytoplasmi chrotitance extramoclear inheritance inheritance inheritance mitochondria Cytoplarmic inheritance mitochondria Cytoplarmic inheritance mitochondria Cytoplarmic inheritance momber variation chromosomal deromosomal ettra chromosomas unigarental inheritance mental ettra chromosomes variant ettra chromosomes variant hundreds of deoxyribonucleic acid Ruth sager euploidy e	non chromosomal inheritance mitichondria genes extra chromosomal mitichondria genes extra chromosomal stable mitochondria genes extra naclear inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance extra naclear inheritance infections disease single mechanism intra chromosome plants intra naclear inheritance extra naclear inheritance extra naclear inheritance infections plants intra naclear inheritance extra naclear inheritance infections intra naclear inheritance extra naclear inheritance infections infections infections infections infections infections infections infections inheritance inheritance inheritance inheritance inheritance inheritance infections infections inheritance inheritance infections infections inheritance inheritance infections inf	minicipantial genes mitochonical inferitance ino chromosomial genes chromosomial genes chromosomial genes chromosomial inheritance chromosomial inheritance chromosomial inheritance bon chromosomial inheritance or no medellin Hershel Mitchell 1902 Hershel Mitchell 1902 Hershell Mitchell 1902 Hershell Mitchell 1902 Carl corres 1882 aneuploidy nullisomy tetrasomy nullisomy tetrasomy nullisomy tetrasomy nullisomy tetrasomy 1904 N - 1 monosomic monosomic 2N - 1 monosomic monosomic color al the three options 1940 Hershell Mitchell 1940 Hershell Mitchell 1940 Hershell Mitchell 1940 N - 1 monosomic monosomic color al the three options 1940 Chromosomes color al the three options 1940 Chromosomes color and the three color and the t
Cypipame interfance iso known is Genes hocased in cargonale in the cypipam are	circumoname numerations of experiments of expendences of expension of the experiment of experiments of experime	extra chromosonia inheritance chromosonia gress non chromosonia gress inter chromosonia gress inter chromosonia gress inter chromosonia gress inter chromosonia gress occupatami inheritance extrametoer inheritance mitochondria Carl correns 1907 Mendelian chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial inheritance market variation chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonial chromosonia chromosonia chromosonia chromosonia composition chromosonia composition capbioly chromosome paffs capbioly chromosome paffs capbioly	non chromosomal inheritance mitchondrial genes extra chromosomal mitchondrial genes extra chromosomal stable mitchondrial genes extra nuckar inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance chromosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosomal inheritance ethornosoma inheritance ethornosoma inheritance ethornosome galatichornosomes exaultichornosomes fishornosomes galatichornosomes gala	minicronous internance indications and genes mitochonial inheritance inon chromosomal igness chromosomal inheritance chromosomal inheritance both evoplasmic and extrancle non nervolution inheritance non metallin Hold Mitchell 1902 Mitchell 1902 Mitchell 1902 Mitchell 1903 Mitchell 1904 Mitchell 1904 Mitchell 1904 Mitchell 1905 Mitchell 1905 Mitchell 1904 Mitchell 1904 Mitchell 1905 Mitchell 1905 Mitchell 1905 Mitchell 1907 Mitchell 1908 Mitchell 1908 Mitchell 1908 Mitchell 1908 Mitchell 1908 Mitchell 1909 Mitchell 1909 Mitchell 1908 Mitchell 1907 Mitchell 1908 Mitchell 1908 Mitchell 1908 Mitchell 1908 Mitchell 1907 Mitchell 1908 Mitchell 1907 N - 1 20 N - 1 20
Cycipamic interfance iso Anoma is Greek beach in cargonale in the cycipamic and michonalia are known as  Gree present in cycipamic and michonalia are known as  Gree present in cycipamic and michonalia are known as  Gree present in cyclpamic and michonalia are known as  Gree present in cyclpamic and michonalia are known as  Trait controlled by a gree of the mode bet corposed in the progeny is called  Extrandear inferiance is fast on  Gree inferiance are known as  Trait controlled by a gree of the mode bet corposed in the progeny is called  Extrandear inferiance is fast on  Gree inferiance is a fast on  Gree inferiance is  Gree inferiance is  Gree information in the structure of an organian's  Gree information in the structure of an  Gree information in  Gree information  Gree inform	terrspinner inkninger terrspinner terrspinner terrspinner terrspinner terrspinner mitochondrial inheritance matemal effect terromosonal matemal effect terromosonal matemal affect terromosonal matemal inheritance matemal inheri	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess cytoplanne: inheritance ethoritance ethoritance ethoritance inheritance ethoritance mitticchardia of correass 1907 Mendelian chromosomia etro chromosomia etro chromosomia etro chromosomia etro chromosomia etro chromosomia etro chromosomis etro chromosomis etr	non chromosomal inheritance mitchondria genes extra chromosomal mitchondria genes extra chromosomal stable mitochondria genes extra chromosomal extra nackeni inheritance extra nackeni inheritance chromosomal inheritance chromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance erromosomal inheritance infectious disease single mechanism intra chromosomes gaint chromosomes gaint chromosomes gaint chromosome plants thousands of ribonucleic acid Babbani 1880 naneuploidy aneuploidy aneuploidy aneuploidy aneuploidy aneuploidy aneuploidy trisomy	minicronous interfance indictorous metalance indictorous metalance indictorous metalance indictorous and interfance behavioral interfance behavioral interfance behavioral interfance behavioral interfance behavioral interfance behavioral interfance behavioral interfance intermediate interfance accession of gene extra nuclear intermediate interfance phenotypic disease void mechanism mutated chromosomes abnormal chromosomes abnormal chromosomes abnormal chromosomes abnormal chromosomes intermediate interfance phenotypic disease void mechanism mutated chromosomes abnormal chromosomes anary diperan files thvoid gelan bandred of ribonucleic acid Carl correns anapoloity mullisomy mullisomy mullisomy mullisomy interfance tertassomy ter
Cyopama interfance also chooses as Gene located in congrandic in the cyopama rac	crussionar nar. nar.e cytophone: graves cytophone: graves cytophone: graves cytophone: graves inter chromosomal encleus maternal effect cytophasen: internation maternal effect cytophasen: internation Mendel 1904 nonmendedian Mendel 1904 nonmendedian Maternal maternal internation maternal internation maternation maternation maternation monopolos	extra chromosomia inheritance chromosomia greess non chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess inter chromosomia greess cytoplasmi inheritance intertance inheritance metal control of the control of the control of the control of the control of the control of the control of the control of the control of the control of the control	non chromosomal inheritance mitchondria genes extra chromosomal mitchondria genes extra chromosomal stable mitochondria genes extra anckear inheritance extra anckear inheritance chromosomal inheritance extra anckear inheritance infectious disease single mechanism intra chromosome plants intra chromosomes gaint chromosome plants intra chromosomes plants intra chromosome plants intra chromosome plants intra chromosome plants intra chromosome plants intra chromosome plants intra chromosome plants intra chromosome plants intra chromosome reduplication ancuploidy trisomy	minicronouran internance innorkronosomia genes mitochonoiral inheritance inno chronosomali genes chromosomali autonomos stamen bora civopalamia disertiance bora civopalamia mal estimance bora civopalamia mal estimance promosediem inheritance expression of gene extra nuclear intermediate inheritance phenotypic disease vidi mechanism mutated chromosomes anary diperan files thivoid gena borand chromosomes many diperan files thivoid gene babornal chromosomes many diperan files thivoid gene solution pathetica solution terasomy mullisomy tetrasomy tetrasomy mullisomy tetrasomy tetrasomy mullisomy tetrasomy

The sex chromosomes of a human female are	xx	XY	YY	XXY
Who determines the sex of the child? Which of the following supports the chromosomal theory of inheritance	the male Both chromosomesand alleles are pair	the temale homologous chromosomes and alleles	ovum Fertilization restores both the diploi	gamates all the three options
Male and female drosophila flies have	haploid chromosomes	one pair of sex chromosomes	the same number of sex chromosom	homologous chromosomes
I ne allele for drosnopnila eye colour An individual with the genotype XrY will produce which of the following gametes?	is on the Y chromosome XR	is on the A chromosome Xr	Y Y	Xr and Y
Carriers can also be described as	homozygous recessive	homozygous dominant	heterozygous	heterozygous recessive
There is a direct relationship between the frequency of crossing over and the % of recombinant Drosophila chromosome 11 carries the gene for	Gametes antennae type	Phenotype wing type	Genotypes leg length	A.B and C
Triploids have of each kind of chromosomes	1	2	3	4
When an individual has only one of a particular type of chromosome it's described as In humans, turner syndrome	monosomy is a monosomy	disomy involves the sex chromosomes	trisomy results in a female	all the three options
Non disjunction can occur during?	anaphase 1	interphase	metaphase I	prometaphase
polyplody generally results from? which of the following may result in abnormal gametes and offsoring ?	mutation	hybridization translocation	deletion	all the three options
Down syndrom is a?	monosomy	disomy	trisomy	tetrasomy
Down syndrome that tends to run in the family of either the mother or father is the result of? Cri du chat syndrome is the result of a(n) ?	deletion	inversion translocation	duplication deletion	translocation duplication
Chromosomes may be broken by?	viruses	certain organic chemicals	radiation	all the three options
Somatic cells in humans havechromosomes? When a karvotyne is done chromosomes are arranged in pairs based on	23 size	46 shane	92 centromere location	48 all the three options
A display of the pairs of chromosomes is called a?	pedigree chart	karyotype	caricature	chromosome map
In humans there arepairs of autosomes? the most common autosomal trisomy among humans is ?	22 trisomy 21	23 trisomy 13	44 trisomy 18	88 trisomy 17
Individuals with down syndrome have three copies of?	the X chromosome	chromosome 13	chromosome 21	the Y chromosome
Which is an X- linked recessive disorder? The alleles of most on linked genetic diorder are on the	color blindness V	hemohpilia V	muscular dystrophy	all the three options
The anexes of most sex-mixed generic disorder are on the chromosome? Who coined the term linkage?	Correns	1 Mendel	Morgan	de vries
Mendel did not obsorve linkage due to?	mutation	Synapasis	Crossing over	Independent assortment
I ne prenomenon of unkage was first observed in the plant? Crossing over occur during?	Latnyrus odoratus Pachytene	Diplotene	Datura Diakinesis	Mirabilus Jalapa Pachytene
Crossing over more frequent in ?	Males	Females	embryos	gamates
Crossing over in diploid organism is responsible for? Complete linkage has been reported in?	Dominanceot genes Maize	Segregation of alleles Human female	Recombination of linked genes Male Drosophila	Linkage between genes Female Drosophila
Coupling and repulsion phenomenon was concerned with?	Crossing over	Mutation	Linkage	inversion
Linkage prevents? Which of the following is not unique feature of meiosis?	Homozygous condition Synapsis	Segregation of alleles Homologus recombination	Hybrid formation Reduction Division	Heterozygous condition Diakinesis
Terminal chiasmata are characteristic of?	anaphase 1	Prophase II	metaphase I	metaphase II
Most multicellular eukaryotes form gamets by?	Meiosis Syngamy	Mitosis Binary fission	Oncogenes Mitosis	Binary fission Meiosis
Sexual reproduction involves the alternation of?	Mitosis and oogamy	Isogamy and meiosis	Meiosis and fertilization	Meiosis and oogamy
The crossing over process is believed to have originated from?	Recombinant DNA Somatic	Mutation	Symbiosis Semantic	Gene pair Synergic
In anaphase I,theare pulled apart?	Chiasmata	Centromeres	Kinetochores	Centrioles
Which of the following is not a form of asexual reproduction?	Binary fission	Fragmentation	Budding	Syngamy
which of the following are mismatched? Chiasmata formation and crossing over occur during?	riapioid -n Prophase of mitosis	somatic cens -2n Prophase I of meiosis	∠ygote-n prophase II of meiosis	AB and C
Meiosis results in	2 haploid daughter cells	4 haploid daughter cells	2 diploid daughter cells	4 diploid daughter cells
Which of the following cells undergo meiosis? father of genetics?	Sperm cells mendel	Liver cells Darwin	Unicellular organisms Muller	gamates Bateson
Phenotypic monohybrid ratio is?	9:3:3:1	01:02:01	03:01	02:01
Sex chromosomes also called? Pisum satisum is	Allosome chick pea	Heterosome	Autosome	Euchromosome
Pisum sativum is Unit 4	cnick pea	nce	rat	mice
Cytolodists can use which of the following to describe location of a gene at a specific place on the chromosome	DNA sequence of a chromosomalregie	Protein expression from a chromosom	Banding pattern of stained chromos	Comparison to markers located
cytogenetic mapping? Which of these is a leav characteristic of molecular marker?	Requires that the gene have been clone It is a known gene	Is a high resolutionmethod of gene loc It is located at a known site on the chr.	partial resolution known as loci	poor resolution Known as a gamate
A molecular marker which is amplified by PCR and is polymorphic by length is a (n)?	RFLP	VNTR	AFLP	SNP
A polymorphism is?	Any changes in the NNA sequence	The most common variation of a gene	AFLP	SNP
Molecular markers include? Molecular markers are used to construct?	Chromosome maps	cytogenetic maps	Alkaline phosphatase Physical maps	Geographic maps
RFLP involves	used to identify a specific protein	used to identify a specific DNA	used to a identify a specific RNA	used to identify a both DNA and
Location of quantitative genes on chromosomes are called RAPD is a	quantitative trait loci DNA sequencing based method	qualitative trait loci restrictioon direction based method	maps PCR based method	construct or position of gene Enzyme based method
Set of DNAs generated by using random primers in a PCR reaction is called	RAPD	RFLP	AFLP	insitu hybridization
A unit of measurement on physical maps is	kilobases	centomorgans Mutation	cytological bands	centimeters Companyin
A monomorphic DNA segment is	segment of DNA that exists in many fe	segment of DNA that controls a single	segment of DNA inherited in a dom	a segment of shared by over 999
linkage mappin can determined the distance between which of the following pairs of DNA sequences	AFLP and RFLP	two AFLPs	two known gene	all the three options
				- · · · · ·
Linkage analysis? Radiation hybrids ?	Is used in physical mapping Are human cell lines	Is primarialy used with prokaryotes Are produced by irradiation of UV	Requires linked DNA markers Can hold large piece of chromosom	Is based on recombination Have been used in mapping the
Linkage analysis? Radiation bybrids? Physical mapping?	Is used in physical mapping Are human cell lines Requires large number of organisms	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries	Requires linked DNA markers Can hold large piece of chromosom Detects polymorphic DNA sequence	Is based on recombination Have been used in mapping the Uses RFLPs
Linkage analysis ? Rediation bybrids ? Physical mapping ? A genome marker ?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles dcDNA	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries must be atleast 1000 bp long crDNA	Requires linked DNA markers Can hold large piece of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular	Is based on recombination Have been used in mapping the Uses RFLPs must be a unique DNA sequence Greater RNA
Linkage analysis? Linkage analysis? Physical mapping? Ageome marker? Adeno virus geome is in type? Adenoasciated virus geome is in type?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles dsDNA dsDNA	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries must be atleast 1000 bp long ssDNA ssDNA	Requires linked DNA markers Can hold large piece of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular	Is based on recombination Have been used in mapping the : Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA
Linkage analysis? Linkage analysis? Physical mapping? Ademon virus genome is in type? Ademonassociated virus genome is in type? Ademonassociated virus genome is in type? Ademonassociated virus genome is in type?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles dsDNA dsDNA	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizzs genomic libraries must be atleast 1000 bp long ssDNA ssDNA ssRNA	Requires linked DNA markers Can hold large piece of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular	Is based on recombination Have been used in mapping the : Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA
Linkage analysis?  Kaldianio hybrids?  Physical mapping?  A genome marker?  Adenovirus genome is in type?  Adenovirus genome type is?  Hotps virus genome type is?  Hotps virus genome type is?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles dsDNA dsDNA dsDNA dsDNA 30-40kb	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizzs genomic libraries must be atleast 1000 bp long ssDNA ssRNA ssRNA ssDNA Ukb	Requires linked DNA markers Can hold large piece of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular Stb	Is based on recombination Have been used in mapping the ' Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA D00bb
Linkage analysis?  Thysical mapying?  Adenon virus is  Adenonsociated 'virus genome is in type?  Adenonsociated 'virus genome is in type?  Apha virus genome type is?  Genome integration nature of adeno virus is?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA 39-40kb Non- integration	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries must be atleast 1000 bp long ssDNA ssDNA ssDNA ssDNA lokb intergrating	Requires linked DNA markers Can hold large piece of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular Skb transient	Is based on recombination Have been used in mapping the ; Uses RFLPs Circular RNA Circular RNA Circular RNA Circular RNA Oloubar RNA Oloubar RNA Oloubar RNA
Linkage analysis?  Physical mapping?  Adeno virus genome isin type?  Adenoascelated Virus genome isin type?  Apha virus genome type is?  Genome integration nature of adeno virus is?  Genome integration nature of adeno virus is?  Genome integration nature of adeno virus is?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA 04DNA 39-40kb Non-integration Transient	Is primarialy used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries must be atleast 1000 bp long ssDNA ssRNA ssRNA ssRNA libko intergrating Long lasting certain organic chemicals	Requires linked DNA markers Can hold large pices of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular Circular Circular Skb transient potential long lasting rotistion	Is based on recombination Have been used in mapping the ; Uses RPLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA IOU8b Non transient virulant all the three sections
Linkage multiplication hybrids?  Physical mapping?  Adenon wing genome is in type?  Adenonsectined wins genome is in type?  Alpha virus genome type is?  Hopps virus genome type is?  Genome integration nature of adeno virus  Expression pattern of adeno virus  Chromosomes may be broken by?  Chromosomes may	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA daDNA So-40kb Non- integration Transient Virtuses	Is primarily used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries uses be atleast 1000 bp long usDNA skDNA skDNA JOAb Long lasting certain organic chemicals Segregation of alleles	Requires linked DNA markers Can hold large rules of chromsons Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular Circular Circular Detential long lasting radiation Hybrid formation	Is based on recombination Have been used in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA I Diokb Non transient virulant all the three options Heterozygous condition
Linkage multiplication hybrid	Is used in physical mapping Are human cell limes Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA Mon-integration Non-integration Transient Homosygous condition Somatic	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA ssDNA ssDNA ssDNA ssDNA Long tasting Cartian organic chemicals Segregation of alleles Haploid	Requires linked DNA markers Can hold large face of chromosom Detects polymorphic DNA sequence must be atleased SOBP long Circular Circular Circular Circular Circular Circular Skb Transient Insteint potential formation Hybrid formation Hybrid formation	Is based on recombination Have been seld in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Torchar RNA 100kb Non transient virulant all the three options all the three options
Linkage manysis? Physical mapying? Adeno virus is genome isin type? Adenosociated virus genome isin type? Apha virus genome type is? Genome integration nature of adeno virus is? Genome integration nature of adeno virus is? Genome integration nature of adeno virus is? Chromosomes may be blocken by? Linkage prevents? Linkage prevents? Compote linkage has been reported in? Complete linkage has been reported in?	Is used in physical mapping Are human cell limes Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA ddDNA bon-integration Transient viruses Homozygous condition Somatic Ladwyus doortuus	Is primarily used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries stDNA ssDNA ssDNA ssDNA ssDNA Long lasting certain organic chemicals Segregation of alleles Hapfoid Human female pisum sativum	Requires linked DNA markers Can hold large picce of chromsomo Detects polymorphic DNA sequence must be alteased Stop long Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Skb transient John Stap Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop Stop	Is based on recombination Have been used in mapping the ; Uses RPLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Olokb Non transient virulant all the three options Heterozygous condition Symetgic Female Drosophila Mirabius jalapa
Linkage and represents? Linkage and represents? Linkage and represents? Linkage and represents? Linkage re	Is used in physical mapping Are human cell limos Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA daDNA daDNA habba	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries ums be atleast 1000 bp long sdDNA sdDNA sdDNA sdDNA sdDNA lokb intergrating intergrating certain organic chemicals Segregation of alleles Haploid Human female pisum sativum Low of segregation	Requires linked DNA markers Can hold large pices of chromsomo Detects polymorphic DNA sequence must be atleased Circular	Is based on recombination Have been used in mapping the ; Uses RPLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA OlObb Non transient virulant all the three options Heterozygous condition Synergie Franske Drosophila Mirabilos jalapa Lav of hermodynamics
Linkage multiplication by the form of the plant?  Linkage multiplication to mends for price plant for the plant?  Linkage multiplication to mends for plant for the plant?  Linkage multiplication to mends for the plant?  Linkage proverse:  P  Linkage proverse:  P  Linkage proverse:	Is used in physical mapping Are human cell limes Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA hatter segurition National and the second second second second National Second Second Second Second Second Maize Lathyrus odoratus Lathyrus odoratus Dadpendent assortment Bacterium	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries saDNA saDNA saNNA saNNA saNNA saNNA biother santa saNNA saNNNA saNNA saNNNA saNNNA saNNNA s	Requires linked DNA markers Can hold large fuce of chromosom Detects polymorphic DNA sequence must be atleased Circular Circular Circular Circular Circular Circular Circular Sub Transint fung Instanti Polymonia Polymonia Male Dresophila Datura Law of dominance Human mitochondriion 1990	Is based on recombination Have been used in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Diokb Non transient virulant all the three options Heteroxygous condition Synergic Female Drosophila Mirabilus jalapa Law of thermodyamics Fruit fly
Linkage multiplication by bricken by the set of the set	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA Non-integration Non	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA ssDNA ssDNA ssDNA ssDNA Long lasting Long lasting certain organic chemicals Segregation of alleles Haptoid Human female pisum sativum Law of segregation Yeast Yeast 1970	Requires linked DNA markers Can hold large fuces of chromsons Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Skb Mak Drosophila Datura Law of domiance Human mitochondriion 1990	Is based on recombination Have been used in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Olokb Non transient virulant all the three options Heteroxygous condition Symergic Emaile Drosophila Mirablus jalapa Law of thermodynamics Fruit By 2000
Linkage markyis? Physical markyis? Physical markyis? Physical markyis genome isin type? Adeno virus is genome isin type? Adeno virus is genome isin type? Alpha virus genome type is? Genome integration nature of adeno virus is? Composed in large to broken type? Linkage prevents? Linkage is an exception to metal's principle of? The hormon genome type is queeded was that of a? The hormon genome top cost price of was that of a? The hormon genome to contain approximated was that of a? The hormon genome to contain approximated was that of a? The hormon genome to contain approximated was that of a? The hormon genome top contain structure data was that of a deno virus is the plant? Linkage is an exception to metal's principle of? The hormon genome top contain data virus in? The hormon genome top contain data virus in market data with yeard? Linkage for enones the containt data virus in the hormon genome contains approximated virus that data with yeard its matching that data was that data with yeard its matching that data was that data was that yeard its the exercit may as the enones its ? The satisfield of genomics that deals with yeard its matching that data was that yeard its the period that data was that yeard its the pe	Is used in physical mapping Are human cell limes Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA ddDNA bon-integration Transient viruses Homozygous condition Somatic Ladvyus doratus Independent assortment Bacterium J980 J00,000 functional genomics	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA ssDNA ssDNA ssDNA ssDNA lokb intergrating Long lasting certain or ganice chemicals Segregation of alleles Haphoid Human female pisum sativum Law of segregation Yeast 1970 Yeast 1970 Shouthart genomics back crosses	Requires linked DNA markers Can hold large fuces of chromsoms Detects polymorphic DNA sequence must be altess 50p long Circular C	Is based on recombination Have been used in mapping the ; Uses RPLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Diokb Non transient virulant all the three options Heterozygous condition Symergic Heterozygous condition Symergic Hermodynamics Female Drosophila Mirabilus jalapa Law of thermodynamics Fruit fly 2000 30,000,000,000 computational genomics wene malvasis
Linkage analysis?  Adeno virus isin type?  Adeno virus is genome isin type?  Adeno virus is genome isin type?  Apha virus genome type is?  Genome itegration nature of adeno virus is?  Genome itegration nature of adeno virus is?  Genome itegration nature of adeno virus is?  Genome integration nature of adeno virus is?  Genome integration nature of adeno virus is?  Complete linkage has been reported in?  Complete linkage has been reported in her plant?  Linkage year exception to mendels principle of?  The planements of the sequenced was flat of a?  The beam genome contains approximately mucleotds?  The subfield of genomes had east wire gation wire yard?  The genomes model many formation genomes is ?  The genetic maps of genomes are constructed using ?  The genetic maps of genomes are constructed using ?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA daDNA daDNA hatta and the alleles daDNA daDNA hatta and the alleles daDNA hatta and the alleles homozygous condition Somatic Maize Labyrus odoratus Indipendent assortment Bacterium Bacterium Bactorium Jis00 300,000	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries safDNA safDNA safDNA safDNA safDNA safDNA lokb intergrating Long lasting certain organic chemicals Segregation of alleles Haphoid Human female pisam sativum Law of segregation Yeast Jayo Ogo SafDNA SAFDNA SAF	Requires linked DNA markers Can hold large pices of chromsomo Detects polymorphic DNA sequence must be atlassa 50bp long Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Shb Transient Joatura Make Drosophila Datura Law of dominance Law of dominance Law of dominance Human mitochondrion 1990 3.000,0000 c comaparative genomics phylogenetic analysis 24	Is based on recombination Have been used in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Olobb Non transient virulant all the three options Heterozygous condition Synergie all the three options Heterozygous condition Evande Drosophila Mirabilos jalapa Law of thermodynamics Fruit By 2000 300,000,000 computational genomics gene analysis 46
Linkage multiplication hybrids ?  Addition hybrids ?  Physical mapping ?  Addition hybrids ?  Addition hybrids genome is in type?  Addition kybrids genome is ?  Compared and the type is?  Compared and the type is t	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Mon-integration Non-integ	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries subNA subNA subNA subNA subNA loab intergrating Long lasting certain organic chemicals Segregation of alleles Haploid Human female pisum sativum Law of segregation Vesa Vesa UPS of the substantiation Discontrol generation Segregation Law of segregation Vesa UPS of the substantiation Discontrol generation Set Substantiation Set Substan	Requires linked DNA markers Can hold large fuces of chromosom Detects polymorphic DNA sequence must be atleast 50bp long Circular Sab Sab Sab Sab Sab Sab Sab Sab Sab Sab	Is based on recombination Have been seld in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA A Circular RNA A DOA Non transient Vinilant all the three options all the three options All the three options Heteroxygous condition Symetgic Female Drosophila Mirabilus jalage Hermodynamics Fruit By 2000 000,000,000 computational genomics general years Symetgic
Linkage markysis? Physical mapping? Physical mapping? Physical mapping? Adeno virus is genome is in type? Adeno virus is genome is in type? Adeno virus is genome is in type? Apha virus genome type is? Genome integration nature of adeno virus is? Genome integration nature of adeno virus is? Genome integration nature of adeno virus is? Composed pathods broken by? Composed intage to broken by? Linkage prevents?	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA have all the second second ddDNA have all the second homozygous condition Somatic Homozygous condition Somatic Lathynus odoratus Independent assortment Basterium 1980 J00,000 functional genomics genetic crosses 13 1.20%	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA ssDNA ssDNA ssDNA Uokb intergrating Long lasting certain organic chemicals Segregation of alleles Hapdoid Human female pisum sativum Law of segregation Yeast Yeast 2300,000 structural genomics back crosses 23 8,80% centimeter student ttes	Requires linked DNA markers Can hold large fuces of chromsoms Detects polymorphic DNA sequence must be alteased Stop long Circular Skb transient Byber Somantic Semantic Mak Drosophila Datura Law of dominance Human mitschondrion 1990 Circular Shol, Outon Shol, Outon Shol, Outon Shol, Outon Circular Shol, Outon Circular Shol, Outon Circular Shol, Outon Circular Shol, Outon Circular Circula	Is based on recombination Have been used in mapping the ; Uses RFLPA must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA D00kb Montransient virulant all the three options Heterozygous condition Symergic all the three options Heterozygous condition Symergic Hermodynamics Fremale Drosophila Mirabilus jalapa Law of thermodynamics Fruit By 2000 000,000,000 computational genomics gene analysis 46 88%
Linkage markysis?  Addinon hybrids?  Physical mapping?  Addeno virus is in type?  Addeno virus is genome is in type?  Addeno virus is in type?  Addeno virus is in type?  Addeno virus is?  Genome integration nature of addeno virus is?  Chomosomes many be broken by?  Linkage prevents?  Linkage prevents?  Linkage prevents?  Linkage integration of addeno virus is?  Complete linkage via fire observed in the plant?  Linkage integration of addeno virus is?  The homma genome project formally begin in what year?  The homma genome project formally begin in what year?  The human genome contains approximately mucleotido?  The subfield of genomics for addens wire are in genomes is ?  The genetic map of human genome contains of provimated using?  The genetic map of human genome contains of provimated using?  The genetic map of human genome contains of provimated using?  The genetic map of human genome contains of provimated using?  The genetic map of human genome contains of provimated using?  The genetic map of human genome contains of provimated using?  The genetic map of human genome contains of the other of provide linkage here two to genitic loci ?  For human genome 1, map unit corresponded approximately	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA ddDNA Maize Ladyuns odden alles Homozygous condition Somatic Ladyuns oddentus Independent assortment Basterium 1980 Sonodic 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic Basterium 1980 Sonodic 198	Is primarily used with prokaryotes Are produced by irradiation of UV Utilizes genomic libraries subNA subNA subNA subNA subNA subNA lokb intergrating Long lasting certain organic chemicals Segregation of alleles Haphoid Human female pisum sativum Law of segregation Yeast 1970 Yeast 1970 Subok crosses 23 880% centimeter sudden t test 100,000 by to b	Requires linked DNA markers Can hold large pices of chromsoms Detects polymorphic DNA sequence must be altess 50p long Circular C	Is based on recombination Have been used in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Montanisent Vinulant all the three options Heteroxygous condition Symetryic Heteroxygous condition Heteroxygous condition Symetryic Heteroxygous condition Minrablus jalapa Law of thermodynamics Freind Py 2000 000,000,000 computational genomics gene mal-ysis efforts States Networks
Linkage markysis? Addinon hybrids? Physical mapping? Addinon xitys genome isin type? Addinon xitys genome isin type? Addinon xitys genome isin type? Addinon xitys genome is? Addinon xitys genome is? Addinon xitys genome is? Compose insign point nature of addino xitys is? Genome integration nature of addino xitys is? Genome integration nature of addino xitys is? Compose insign point have on years Expression pattern of addino xitys is? This presents Define the enter reported in? Complexe linkage has been reported in? The presents Define the enter reported in? The presents Define the enter reported in? The presents Define the enter reported in? The barrow points in the data is genome is a sequencing of outrie genomes is ? The presents approximatelymucleotide? The presents approximately many is a sequencing of outrie genomes is ? The genomes that data with from a genitic cross involving 2 genes in which a total of 100 progenic are counted. The recombination frequency between these 2 On a genetic maphotisme between genes are measured in ? The human genome counds approximatelymucleotide? The present map of throman genome counds of genomics are measured in ? The present map of throman genome counds of genomics are measured in ? The present map of throman genome counds of genomics are measured in ? The present map of throman genome counds of genomics are measured in ? The present map of throman genome counds of genomics are measured in ? The present map of throman genome counds of genomics are measured in ? The present map of throman genome counds of genomics are measured in ? The present map of counds throw genitic unciper agenes is must be a total of 100 progenic are counted. The recombination frequency between these 2 On a genetic maphotic between these measured in ? The human genome counds of motion in a statistical botal was the set of restriction fragments generated by a particular restriction en	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA boxis integration Non-int	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries saDNA saRNA saRNA saRNA binergating Long lasting Long	Requires linked DNA markers Can hold large fuce of chromosom Detects polymorphic DNA sequence must be atleast of bolog Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Sab Soft Sab Semantic Sab Semantic Sab Semantic Sab Semantic Sab Semantic Sab Sab Sab Sab Sab Sab Sab Sab Sab Sab	Is based on recombination Have been self in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA all the three options all the three options and the three options all the
Linkage multiplications by the second	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Maine Non-integration Non-integration Non-integration Non-integration Non-integration Non-integration Non-integration Non-integration Non-integration Soundie Laflynus odoratus Independent assortment Bacterium 1980 S00,000 S00	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA september 2000 by long stDNA stNNA stNA stNA lobb intergrating Long lasting certain organic chemicals Segregation of alleles Haptoid Human female pismu sativum Law of segregation Human female pismu sativum Law of segregation Veast Veast 23 300,000 32 33 8,0% Statutural genomics tack crosses 23 8,0% Statutural genomics tack crosses 23 8,0% Statutural genomics tack crosses 23 8,0% Statutural genomics tack crosses 23 8,0% Statutural genomics tack crosses 24 8,0% Statutural genomics tack crosses 25 8,0% Statutural genomics tack crosses 26 27 28 20 28 20 20 20 20 20 20 20 20 20 20 20 20 20	Requires linked DNA markers Can hold large fuces of chromoson Detects polymorphic DNA sequence must be atleast 50bp long Circular Circular Circular Circular Circular Circular andiation long lasting potential long lasting radiation Hybrid formation Semantic Mak Drosophila Datura Law of dominance Human mitochondriion 1990 Carcomparative genomics phylogenetic analysis 24 24 23 3000,000,000 c comparative genomics phylogenetic analysis 24 31 28 31 28 31 28 31 28 31 28 31 20	Is based on recombination Have been used in mapping the ; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Otothe Non transient virulant all the three options Heterozyogus condition Symergic all the three options Heterozyogus condition Symergic Heterozyogus condition Mirablus jalapa Heterozyogus condition Symergic Heterozyogus condition Mirablus jalapa Heterozyogus condition Mirablus jalapa Heterozyogus condition Symergic Symergic Symergic Heterozyogus condition Mirablus jalapa Symergic Symergic Symergic Mirablus jalapa Heterozyogus condition Symergic Heterozyogus condition Symergic Allow Condition Symergic Allow Condition Symergic Allow Condition Symergic Heterozyogus condition Heterozyogus
Linkage markysis? Physical mapping? Addeno virus isin type? Addeno virus is genome isin type? Addeno virus is genome isin type? Addeno virus is genome isin type? Addeno virus is? Genome integration nature of addeno virus is? Genome integration nature of addeno virus is? Genome integration nature of addeno virus is? Composed in target of how of the plant? Linkage prevents? Unikage prevents? Unikage prevents? Unikage prevents? The plante short of this genome to be sequenced was that of a?? The horman genome project formally began in what year? The horman genome project formally began in what year? The burnam genome contains approximated using? The genetic maps of human genome contaits of movies are genome site? The genetic maps of human genome contaits of movies are included in genetic and of 100 progenic are counted. The recombination frequency between these 2 On a genetic map of human genome contaits of moviemated in the site? The human genome contaits of moviemated in the genetic maps of human genome contaits of moviemated in the site? The genetic maps of genomes are constated of agenomes in constated of 100 progenic are counted. The recombination frequency between these 2 On a genetic maps of human genome constates of " The formation genetic result from a genetic constate in the integration of the site in the site of restriction frequency between these 2 On a genetic maps of ONA that contains? The human genome contains approximated in the site of restriction frequency between these 2 NortKas are regions of NAM and contains? The human genome contains approximated in the site of restriction frequency between these 2 NortKas are regions of NAM that contains? The human genome contains approximated in that size of restriction frequency between these 2 NortKas are regions of NAM that contains? The human genome contains approximated in that size of restriction frequency between these 2 NortKas are regions of NAM that contains? The human ge	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA Non-integration Non	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA sstDNA sstNA sstNA sstNA sstNA Long lasting certain organic chemicals Segregation of alleles Human female pisum sativum Law of segregation Human female pisum sativum Law of segregation Yeast 1970 3000,000 structural genomics back crosses 23 880% centimeter student tes 100,000 RFLP a variable number of tandem repeate r 100 Caemophilus clegans	Requires linked DNA markers Can hold large fuces of chromsoms Detects polymorphic DNA sequence must be alteased to the sequence must be alteased to the sequence Circular Skb transient Determine Hybrid formation Hybrid Commission Hybrid Circular C	Is based on recombination Is based on recombination Is we been used in mapping the '; Uses RPLPA must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Constants Circular RNA C
Linkage manysis? Addino hybrids? Physical mapping? Addino xitys genome isin type? Addino xitys genome is? Calculate the provide the pro	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Maine South and the second second Maine Maine Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Maine Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Maine Caroline Maine Caroline Maine Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Labyus odoratus Maine Second Second Ecoli genome musus	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries saDNA saRNA saRNA saRNA saRNA saRNA saRNA saRNA loab intergrating intergrating intergrating intergrating comparing and challes samparing and samparing samparing an	Requires linked DNA markers Can hold large fuce of chromsons Detects polymorphic DNA sequence must be atleast of the sequence Circular Mala Drosophila Datura Mala Drosophila Datura Law of dominace Human miticchondriion 1990 3.000,0000 Circometica analysis 24 Low Stat Low STR Long Stat Long Long Long Long Long Long Long Long	Is based on recombination Have been self in mapping the ' Uses RFLPs must be a unique DNA sequence Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA Circular RNA di the three options all the three options all the three options all the three options Hearonzy oparations Hearonzy options Hearonzy options Hearonzy options Hearonzy options Mirability options Mirability options Hearonzy options Mirability
Linkage multiplications by the second	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Non-integration Non-integ	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA seproduced by irradiation of UV sisDNA sisDNA SisDNA SisDNA Long lasting Control of alleles Hanna fermale pismu sativum Lapoid Human fermale pismu sativum Lapoid Segregation of alleles Hapdoid Human fermale pismu sativum Law of segregation Veast Veast 23 SisDNA SisDNA Control Bennets Sister Library Sister Lib	Requires linked DNA markers Can hold large fuce of chromoson Detects polymorphic DNA sequence must be atleast 50b long Circular Circular Circular Circular Circular Circular andiation Bybrid formation Bybrid formation Bybrid formation Bybrid formation Semantic Mak Drosophila Datura Law of dominance Human mitochondriion 1990 Carconplata Datura Law of dominance Human mitochondriion 1990 Carconplata Datura Law of dominance Human mitochondriion 1990 Carconplata Datura Law of dominance Human mitochondriion 1990 Southorne State Law State Comparative genomics phylogenetic analysis 24 Jake Joba Store method I dod score method I dod score method I dod score method I dodo Stra Fine Constanti State Constanti State Constanti Cons	Is based on recombination Have been used in mapping the ; Uses RFLP6 must be a unique DNA sequence. Circular RNA Circular
Linkage marbysis?  Addino hybrids?  Physical mapping?  Addino xinysis genome is in type?  Apha virus genome type is?  Genome integration nature of addino virus is?  Genome integration nature of addino virus is?  Compared inkage has been reported in?  Compared inkage has been reported in?  Compared inkage has been reported in?  The benomes one to be sequenced was that of a?  The benomes one to be sequenced was that of a?  The benomes one contins approximately nucleotids?  The subfield of genomics that deals with genitic mapping and sequencing of entire genomes is ?  The genetic maps of summar contains approximately uncleotids?  The genetic maps of summar contains approximately are sub-tain virus in a uncleotids?  The for to complex genite result from a genitic corso involving 2 genes in which a total of 100 progenic are counted .The recombination frequency between these 2  On a genetic map.of Summar contains approximately	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA have all the second second ddDNA have all the second homosygous condition Non-integration Non-integration Non-integration Non-integration Non-integration Non-integration Non-integration Homosygous condition Somatic Homosygous condition Somatic Lathytus doratus Independent assortment Basterium Justo Somatic Basterium Justo Somatic Basterium Justo Somatic Basterium Justo Nuclional genomics genetic crosses Justo Nucli sequere Loob Nucli sequere Loob Nucli sequere Loob Nucli sequere Loob Nucli sequere Somatic Loob Konise Somatic Loob Nucli sequere Somatic Loob Nucli sequere Somatic Loob Nucli sequere Loob Conise Somatic Recond Basterium Somatic Nucli Somatic Somatic Nucli Somatic Somatic Nucli Somatic Somatic Nucli Somatic Somatic Somatic Nucli Somatic So	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries stDNA seproduced by irradiation of UV sisDNA sisDNA SisDNA SisDNA Long lasting certain organic chemicalls Segregation of alleles Hapdoid Human female pisum sativum Law of segregation Yeast UP70 Souther Sister	Requires linked DNA markers Can hold large fuces of chromsoms Detects polymorphic DNA sequence must be alteast 50bp long Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Skb Transient Hybrid formation Semantic Mak Drosophila Datura Law of dominance Human mitschondrion 1990 Comparative genomics phylogenetic analysis 24 Austransi Stransie Circular Displogenetic analysis 24 a long stretch of a single repeated m 1,000,000 STR a long stretch of a single repeated m 1,000 Haemophilus influenza protoome yeast indertiance of recessive genes seggregation chromosom 16	Is based on recombination Is based on recombination Is we been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular R
Linkage manysis? Addition hybrids? Physical mapping? Addition type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function tautor of algoes transmitter of type is? Common function to medder for three plant? The plant system contains approximately recordstate three plantsmitter of type is? The function for the plant system contains approximately	Is used in physical mapping Are human cell index Requires large number of organisms must occur in multiple alleles daDNA	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries substances and the substances substances and the substances substances and substances substances and substances substances and substances substances and substances substances and substances substances and substances pisum substances substances and substances pisum substances and substances substances and substances and substances and substances and substances and substances and substances a	Requires linked DNA markers Can hold large fuce of chromosom Detects polymorphic DNA sequence must be atleast Circular Circular Circular Circular Circular Circular Circular Circular Datura Potenti bang lassing potenti bang lassing potenti bang lassing Potenti bang lassing Potenti bang lassing Potenti bang lassing Potenti bang Simonica Human mitochondriion 1990 3.000,0000 Carcompetitive genomics phylogenetic analysis 24 L2% Low STR Low STR Long attech of a single repeated m London STR Long stretch of a single repeated m London London London STR Long stretch of a single repeated m London London London London London London STR Long stretch of a single repeated m London	Is based on recombination Is based on recombination Is we been self in mapping the '; Uses RPLP6 must be a unique DNA sequence Circular RNA Circular
Linkage markysis?  Addition hybrids?  Physical mapping?  Addition type is?  Compared mature of addition type is?  Genome integration mature of addition type is?  Genome integration mature of addition type is?  Compared mature of addition type is genomes as mecanized in the plant?  The pherometron of Unitage was first observed in the plant?  The humma genome contains approximately	Is used in physical mapping Are human cell inlos Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA how-integration Non-integ	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries substances and the substances substances and substances and substances and substances and substances and substances and substances and substances substances and substances and substances and subst	Requires linked DNA markers Can hold large picce of chromsons Detects polymorphic DNA sequence must be atleast of the sequence Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Shb transient Dog Lasting Sho Semantic Male Drosophila Datura Law of dominance Homan micochondrison Homan micochondrison Law of dominance Homan micochondrison Datura Law of dominance Homan micochondrison Strate Sho Sho Sha Datura Law of dominance Homan micochondrison Strate Law of cominance Law of dominance Homan micochondrison Strate Law of cominance Law of the semantic semantic semantic semantic labers Law of the semantic homan semantic semantic labers La ong stretch of a single repeated m 10,000 STR Longoptic for a single repeated m 10,000 Strates Long stretch of a single repeated m 10,000 Strates Longenetic Indage , centimorgan chromscome 16 genetic linkage , centimorgan chisma_chromosomal interference Seminori Sho Methol Methol Strates Sho Methol Methol Strates Sho Methol Strates	Is based on recombination Is based on recombination Is we been set in mapping the '; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage markysis? Addino hybrids? Physical mapping? Addino hybrids? Physical mapping? Addino hybrids genome isin type? Addino hybrids genome isin three genome is genome integration that of addino hybrids Expression pattern of addino hybrids is? Composed intage hybrids hybrids genome is? Composed intage hybrids hybrids genome is? Composed intage hybrids hybrids hybrids Expression pattern of addino hybrids Expression pattern of genome is a constraint of a? The human genome contains approximately The dotted of genomics that deadins with genitic mappring and sequencing of entire genome is ? The genetic mapp of futura genome consists of	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Maine and the alleles ddDNA Maine and the alleles Maine and the alleles Maine and the alleles Homozygous condition Somatic Maine Ladynus doratus Independent assortment Bacterium 1980 Maine aller alleles Homozygous condition Bacterium 1980 Maine aller Bacterium 1980 Maine aller Bacterium 1980 Maine aller Bacterium 1980 Maine aller Maine all	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries substa tailout the substantiation of UN substantiation of UN substantiation of the substantiation of the substantiation of the substantiation substantiation of the substantiation of the Supergration of tailedes Human femule pisum sativum Law of supergration Substantiation of the substantiation substantiation of the substantiation substant test to the substantiation of the substantiation substantiation of the substantiation of the substantiation of the substantiation substantiation of the substantiation of the substa	Requires linked DNA markers Can hold large fuces of chromoson Detects polymorphic DNA sequence must be atlass 3 belog long Circular Circular Circular Circular Circular Circular andiation Hybrid formation Bybrid formation Skb transient Datura Law of dominance Human mitochondrion 1990 atura Law of dominance Human mitochondrion 1990 atura Law of dominance Human mitochondrion 1990 atura Comparative genomics phylogenetic analysis 24 Jama Journa La% Journa STR Tal a long stretch of a single repeated m 10,000 Haemophilus influenza proteome Segregion Influenza Genetic of recessive genes segregion Influenza Informan Influenza proteome (binanchennosan Interference Fertilization restores both the diploi Zygnete-n	Is based on recombination Is based on recombination Have been used in mapping the ; Uses RFLPA must be a unique DNA sequence Circular RNA Circular R
Linkage manysis? Addition hybrid segmentsin type? Addition hybrid segmentsin type? Addition hybrid segmentsin type? Addition hybrid segmentsin type? Addition hybrid segments? Canonascincted vitus segments? Canonascincted vitus segments is? Canonascincted vitus setup rest is? The hybrid setup rest is a distributed vitus of a? The harma genome contains approximatelynucleatids? The subfield of genomics that deal vitus is guitice mapping and sequencing of entire genomes is ? The genetic map of projects formall byeque in whaty and sequencing of entire genomes is ? The genetic map of munan genome contains of	Is used in physical mapping Are human cell index Requires large number of organisms must occur in multiple alleles daDNA	Is primarily used with prokaryotes Are produced by inclution of UV Vilizes genomic libraries substances and the substances substances and the substances substances and substances substances and substances substances and substances substances and substances substances and substances substances and substances prism substances and substances prism substances and substances and substances prism substances and substances and substances and substances and substances and substances prismales without substances and substances mucleosome for mosaic, continues and substances and substances prismales without substances and substances prismales without substances and substances prismales without substances and substances motiverse and prism substances and substances and substances prismales without substances and substances and substances prismales without substances and substances and substances and substances and and substances and substances and substances and substances and substances and substances and substances and substances and substances and substances and substances and substanc	Requires linked DNA markers Can hold large fuces of chromsoms Detects polymorphic DNA sequence must be alleased Stop long Circular Skb transient Byb Stop Maic Drosophila Datura Law of dominance Human mitochondrion 1990 Carcunar Law of dominance Human mitochondrion 1990 Circular Statura Law of dominance Human Mitochondrion 1990 Circular	Is based on recombination Is based on recombination Wess beau anight mapping the '; Uses RPLPA must be a unique DNA sequence Circular RNA Circular R
Linkage multiplications by the set of the se	Is used in physical mapping Are human cell index Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA boxis integration Non-int	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries substances and the substances substances and the substances and the substances and the substances a	Requires linked DNA markers Can hold large picce of chromsons Detects polymorphic DNA sequence must be atleast and the sequence Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Sho Sho Sho Sho Sho Sho Sho Sho Sho Sho	Is based on recombination Is based on recombination Have been self in mapping the; Uses RFLPs must be a unique DNA sequence Circular RNA Circular RN
Linkage markysis? Addition hybrids? Physical mapping? Addition hybrids? Physical mapping? Addition hybridsn type? Addition hybridsn type? Addition hybridsnnn	Is used in physical mapping Are human cell inloss Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Maine Source and the second second Maine Laftynus odoratus Independent assortment Bacterium 1980 Soundo Laftynus odoratus Independent assortment Bacterium 1980 Sou.000 Source and Bacterium 1980 Sou.000 Source and Bacterium 1980 Sou.000 Source and Bacterium 1980 Sou.000 Source and Bacterium 1980 Sou.000 Source and Bacterium 1980 Sou.000 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Bacterium 1980 Source and Source and S	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries substa tailout the substantiation of UN substantiation of UN substantiation of the substantiation of the substantiation of alleles Human female piems sativum Lapoid Human female piems sativum Law of segregation of alleles Human female piems sativum Law of segregation Law of sectornoscomes qualitative ratio Loci Cencophilus e legans Konsoline very Law of sectornoscomes Law of sectornoscomes Law of segregation very Law of segregation very Law of the piper segregation Law of sectornoscomes Law of sectornoscomes Law of sectornoscomes Law of sectornoscomes of the piper segregation very Law of the piper segregation very Law of the piper segregation very Law of the piper segregation very Law of the piper segregation v	Requires linked DNA markers Can hold large fuce of chromsons Detects polymorphic DNA sequence must be atleast polymorphic DNA sequence must be atleast polymorphic DNA sequence Turcular Circular Circular Circular Circular Circular andiation Hybrid formation Skb transient Jong Januar Hybrid formation Semantic Mak Drosophila Datura Law of dominance Human mitochordrinon 1990 Carcular Law of dominance Human mitochordrinon 1990 Carcular Januar Law of dominance Human mitochordrinon 1990 Carcular Janun	Is based on recombination Is based on recombination Have been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular RN
Linkage marbysis? Addino in hybrids? Physical mapping? Addeno virus is genome isin type? Addeno virus is genome isin type? Addeno virus is genome isin type? Addeno virus is? Genome integration nature of addeno virus is? Genome integration nature of addeno virus is? Genome integration nature of addeno virus is? Compared indication before the plan? Linkage prevents? Unlike gametes body cells are called? Compared inlikage was firs observed in the plan? Linkage prevents? Unlike gametes body cells are called? Compared inlikage was firs observed in the plan? Linkage is an exception to meddls principle of? The horman genome project formulty began in what year? The horman genome project formulty began in what year? The horman genome consists of? The horman genome consists of The prevents? Unlike gametes body cells are called? Compared in dades with genitic mapping and sequencing of entire genomes is ? The genetic maps of genomes are constated using 12 genes in which a total of 100 progenic are counted .The recombination frequency between these 2 On a genetic map of human genome consists of	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA how integration Transient Viruses Homozygous condition Somatic Lathyrus odoratus Homozygous condition Somatic Lathyrus odoratus Independent assortment Basterium J980 J0000 functional genomics genetic crosses 13 L20% micrometer chi-square J000 VWTR Restriction their square J00 VWTR Restriction L20% L00 Ecoli Ecoli Chromosomal interference chi square Chromosomal interference as a qualitative trait loci Ecoli The same number of chromosomes as quantitative trait loci Ecoli The same number of chromosomes as quantitative trait loci Ecoli Ecoli Ecoli Ecoli	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries surplant and the surplant and the surplant and the surplant and the Surgergation of alleles Hapdoid Human female pismu sativum Law of surgergation Human female pismu sativum Law of surgergation Veast 1970 3,000,000 Structural genomics and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant and the surplant	Requires linked DNA markers Can hold large fuces of chromsoms Detects polymorphic DNA sequence must be alteast 50p long Circular Circular Circular Circular Circular Circular Circular andiation Hybrid formation Skb transient potential long lasting radiation Hybrid formation Semantic Mak Drosophila Datura Law of dominance Human mitcchondrion 1990 Comparative genomics phylogenetic analysis 24 Mac Drosophila Joungoo c comaparative genomics phylogenetic analysis 24 Joungoo c comaparative genomics phylogenetic analysis 24 Jak Joungoo STR a long stretch of a single repeated m 1,0000 Haemophilus influenza proteome formismon 16 genetic linkage, centimorgan chisma,chromosomal linterference Fertilization restores both the diploi Zygote-a Symbiosis the same number of sex chromosom maps Haemophilus influenza Four point crossing over	Is based on recombination Is have been used in mapping the; Usas RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage numbrisk? Addition hybrids? Addition type is? Common factor type is common factor type is? Common factor type is for the law is? The type common of linkage type is a compared type in the law is? The type compared type is the compared type in the law is? The type compared type is the compared type in the law is? The type compared type is the compared type in the law is? The type compared type is the compared type in type in type in? The type compared type is a gravity common of linkage type are constructed using ? The period type of type commons that calls with gravity The period type of type commons that calls with gravity The period type of type commons that calls with gravity and gravity The period type of type commons that calls with gravity and gravity The period type of type compared type is a maximum that is of restriction fragments generated by a particular restriction endomaclease? Which type of DNA marker corresponds to a base paris mutation that is of restriction fragments generated by a particular restriction endomaclease? Which type of DNA marker corresponds to a base paris mutation that is of restriction fragments generated by a particular restriction endomaclease? Which type of DNA marker corre	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA box integration Non-integ	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries substances and the substances substances and the substances substances and substances substances and substances substances and substances substances and substances Substances and substances substances and substances pieum substances and substances pieum	Requires linked DNA markers Can hold large fuce of chromsoms Detects polymorphic DNA sequence must be atleast Assessment Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Sub Sub Software Systematic Sub Semunic Maile Toromotion Hyma formation Systematic Sub Software Maile Toromotion Hyma mitochondrion 1990 3000,000,000 Circumartic Sub Sub Sub Sub Sub Sub Sub Sub Sub Sub	Is based on recombination Is have been used in mapping the; Uses RFLPs must be a unique DNA sequence Circular RNA Circular
Linkage markysis?  Addition hybrids?  Physical mapping?  Addition hybridsin type?  Addition kybrids genome isin travesin traves	Is used in physical mapping Are human cell inlos Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Non-integration Non-integ	Is primarily used with prokaryotes Are produced by inradiation of UV Uilizes genomic libraries must be atleast 1000 bp long sdDNA sd	Requires linked DNA markers Can hold large pice of chromsons Detects polymorphic DNA sequence must be atleast pice of chromsons Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Determination of the comparison of the comparison of the comparison Hybrid formation Hybrid formation Hybrid Committee Hybrid Datura Law of dominance Human mitochondriion 1990 Jatura Law of dominance Human mitochondriion 1990 Comparative genomics phylogenetic analysis 24 1000,000 Comparative genomics phylogenetic analysis 24 1000,000 Hastenophilus influenza protect inflage, centimorgan chromsoome 16 genetic linkage, centimorgan chisma, chromsoomal interference Ferrilization restores both the diploi Zygote-a Symbiotis the same number of sex chromsoom maps Hastenophilus influenza Four point crossing over 22 pair of chromsome Physiology 23 chromsomes	Is based on recombination Is based on recombination Is we been used in mapping the ; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkge manhysis? Addino hybride? Addino ity prive	Is used in physical mapping Are human cell lines Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Non-integration Non-integration Transient Vivanes Homozygous condition Somatic Labyrous dontus Independent assortment Basterium Justo Contrus Independent assortment Basterium Justo Justo Basterium Justo Justo Justo Hasterium Justo Justo Justo Loob Loob E.coli entomosome al inference. chi square Chromosome al inference as addition there and the foremosomes as aquantiative trait loci E.coli The same number of chromosomes as aquantiative trait loci E.coli The same number of chromosomes as aquantiative trait loci E.coli The same number of chromosomes as aquantiative trait loci E.coli Two-point cross 12 pair of chromosomes Karyology 46 chromosomes Karyology	Is primarily used with prokaryotes Are produced by irradiation of UV Uilizes genomic libraries surphated to the provided of the surphate libraries surphates and the provided of the surphates and the surphates and the surphates and the surphates and the surphates and the surphates and the surphates and the surphates and the surphates and the surphates and the surphates	Requires linked DNA markers Can hold large pices of chromsoms Detects polymorphic DNA sequence must be alteast 50p long Circular Circular Circular Circular Circular Circular andiation Hybrid formation Skb Transient Jobs (Comparent) Hybrid formation Semantic Mak Drosophila Datura Law of dominance Hybrid formation Hybrid for the Hybrid Hybrid for the Hybrid Hybrid for the Hybrid Hybrid for the Hybrid Hybrid Hybrid for the Hybrid Hybrid for the Hybrid	Is based on recombination Is have been used in mapping the; Usas RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage manhysis? Addition hybrids? Addition type is? Addition type is	Is used in physical mapping Are human cell index Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA ddDNA Maker Nor integration Nor integration Security Maker Labyuy odornuts Independent assortment Besterium Maker Labyus odornuts Independent assortment Besterium Maker Security 13 12.0% micrometer Consolition 13 12.0% micrometer Security 13 12.0% micrometer Security 13 12.0% Maker Security 13 12.0% Maker Security 13 12.0% Maker Security 13 12.0% Maker Security 13 12.0% Maker Security 13 12.0% Maker Security 13 12.0% Maker Security 13 12.0% Maker Security 14 15 16 16 16 16 16 16 16 16 16 16	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries substances and the substances substances and the substances substances and substances substances and substances substances and substances substances and substances Substances and substances and substances and substances and substances and substances and substances and subs	Requires linked DNA markers Can hold large fuce of chromsoms Detects polymorphic DNA sequence must be atleast and the sequence Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Sab Detection of the sequence Sab Semantic Male Drosophila Semantic Male Drosophila Sab Semantic Male Drosophila Male Dros	Is based on recombination Is have been used in mapping the; Uses RFLPs must be a unique DNA sequence Circular RNA Circular
Linkge mahysis?  Addinon hybrids?  Physical mapping?  Addonon xings genome is in type?  Addonon xings genome is?  Genome integration nature of addono xings  Expression pattern of addono xings is?  Genome integration nature of addono xings  Expression pattern of addono xings  Proving addono addono xings  Expression pattern of addono xings  Expression pattern addono xings  Expression pattern  Expressin pattern	Is used in physical mapping Are human cell inlos Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Non-integration Non-integ	Is primarily used with prokaryotes Are produced by inradiation of UV Uilizes genomic libraries and the attest at 000 bp long saDNA saNA saNA saNA saNA saNA saNA saNA sa	Requires linked DNA markers Can hold large pice of chromsons Detects polymorphic DNA sequence must be atleast pice of chromsons Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Dotomon Hybrid formation Hybrid Cormation Hybrid Cormation Hybrid Committee Hybrid Circular Circular Circular Hybrid Committee Hybrid Circular Ci	Is based on recombination Is have been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkge manhysis? Addition hybrides? Addition hybrides? Addition hybrides? Addition hybrides? Addition hybrides genome isin type? Addition hybrides genome isin type? Addition hybrides genome isin type? Addition hybrides genome isin trype? Addition hybrides genome is? Compared mature of addition virus is? The home genome consolits approximatelymucleotids? The subfield of genomics that deals with genitic mappring and sequencing of entire genomes is ? The genetic map of formang genome consists of? The formang genome consists of? The formang around consists of? The formang area are genome consists of? The formang area area measured in ? The formang area or area area measured in ? The formang area or area or area area measured in ? The formang area or area or area measured in ? The formang area ore	Is used in physical mapping Are human cell inlos Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Source and the second second ddDNA ddDNA Source and the second second source and second second homozypous condition Non-integration Non-integration Non-integration Non-integration Non-integration Homozypous condition Source and Maize Ladynus doratus Independent assortment Bacterium 1980 Source and second ladgendent assortment Bacterium 1980 Source and second sector and second 100 Each micrometer chi-square 100 Each mouse 100 Each mouse 100 Each mouse of disorders addition thertiance of disorders addition the same number of chromosones as yuanitative trait loci Each Each The same number of chromosones as yuanitative trait loci Each Each Novopoint cross 12 pair of chromosones Each Each Novopoint cross 12 pair of chromosones Karyology Animal cell Charlos	Is primarily used with prokaryotes Are produced by inradiation of UV Uilizes genomic libraries substa to 100 bp long subNA substa to 100 bp long subNA substa to 100 bp long subNA substa to 100 bp long substa to 100 bp lo	Requires linked DNA markers Can hold large fuces of chromsomo Detects polymorphic DNA sequence must be atlass 100 plong Circular Circular Circular Circular Circular Circular Circular andiation Hybrid formation Skb transient Joseff and Joseff Hybrid formation Semantic Mak Drosophila Datura Law of dominance Hybrid formation Hybrid formation Hybrid formation Human mitochondrion 1990 at comparative genomics phylogenetic analysis 24 Joseff and Joseff Joseff and Joseff Joseff Andrewski and Joseff Joseff Andrewski and Joseff Joseff Andrewski analysis 24 Joseff Andrewski Analysis 25 Joseff Andrewski Analysis 25 Joseff Andrewski Analysis 25 Joseff Andrewski Analysis 26 Joseff Andrewski Analysis 27 Joseff Andrewski Analysis 27 Joseff Andrewski Analysis 28 Joseff Andrewski Analysis 29 Joseff Andrewski Analysis 20 Joseff Analysis 20	Is based on recombination Is hew been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage manysis? Addition hybrid sequences and the sequences of th	Is used in physical mapping Are human cell index Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA box integration Non-integ	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries substances and the substances substances and the substances substances and substances substances and substances substances and substances substances and substances substances and substances substances and substances and substances and substanc	Requires linked DNA markers Can hold large fuce of chromsoms Detects polymorphic DNA sequence must be atleast Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Sub Sub Sub Sub Sub Sub Sub Sub Sub Sub	Is based on recombination Is have been self in mapping the; Uses RFLPs must be a unique DNA sequence Circular RNA Circular
Linkge manbysis?  Addinon hybrids?  Physical mapping?  Addinon type isin type?  Addinon type is?  Addinon type is?  Addinon type is?  Addinon type is?  Compared type is the observed type is the observed type is the plant?  Linkge preserved type is the observed type is the observed type is the observed type is the plant?  Linkge type is the observed type is t	Is used in physical mapping Are human cell in loss Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA Monitorial and the alleles daDNA Monitorial and the alleles data alleles da	Is primarily used with prokaryotes Are produced by inclusion of UV Uillizes genomic libraries and the attest of the attest saturation of the attest saturation of the attest saturation of attest Saturation of attest Saturation of attest Saturation of attest Human formalies Human formalies Human formation of attest Saturation of attest Human formation of attest Human formation of attest Saturation of attest Saturation of attest Human formation of attest Saturation of attest Human formation of attest Saturation of attest Human for	Requires linked DNA markers Can hold large pice of chromsoms Detects polymorphic DNA sequence must be atleast of the sequence Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Domession Hybrid formation Hybrid Cornation Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Hybrid Commission Commission Commission Commission Hybrid Commission Commission Hybrid Commission C	Is based on recombination Is have been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage manhysis? Project manhysis? Project manhysis? Project mapping? Project mapping? Project mapping? Project mapping? Project mapping? Project mapping m	Is used in physical mapping Are human cell inloss Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Source and the second second second ddDNA Source and second second second Non-integration Non-integ	Is primarily used with prokaryotes Are produced by inradiation of UV Uilizes genomic libraries must be atleast 1000 bp long sdDNA sd	Requires linked DNA markers Can hold large fuces of chromsomo Detects polymorphic DNA sequence must be atleast 500 long Circular Circular Circular Circular Circular Circular Circular andiation Hybrid formation Skb Sumatic Skb Semantic Mak Drosophila Datura Law of dominance Hybrid formation Hybrid formation Hybrid formation Hybrid formation Hybrid Scomettic Mak Drosophila Datura Law of dominance Hybrid formation Hybrid Scomettic Jaura Law of dominance Human mitcohordrion 1990 comparative genomics phylogenetic analysis 24 Jaura Jaura Jaura Jaura Law of dominance Human mitcohordrion 1990 comparative genomics phylogenetic analysis 24 Jaura Jaur	Is based on recombination Is have been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage manipais? Addono hyperball and hyperball an	Is used in physical mapping Are human cell index must occur in multiple alleles daDNA daDNA daDNA daDNA daDNA how integration New integration	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries and the attest at 000 bp long saDNA saRNA sAR	Requires linked DNA markers Can hold large fuce of chromsoms Detects polymorphic DNA sequence must be atleast Circular Sub Sub Sub System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter System Hyter Hyter System Hyter Hyter System Hyter Hy	Is based on recombination Is have been self in mapping the; Uses RFLPs must be a unique DNA sequence Circular RNA Circular
Linkage manhysis? Project mapping? Project mapping? Project mapping? Project mapping mapping? Project mapping mapping? Project mapping mapping? Project mapping mapp	Is used in physical mapping Are human cell in lons Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA how-integration Non-int	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries must be atleast 1000 bp long saDNA assTNA	Requires linked DNA markers Can hold large pice of chromsoms Detects polymorphic DNA sequence must be atleased Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Document Hybrid formation Hybrid formation Hybrid formation Hybrid Samanic Hybrid formation Hybrid Samanic Human mitochondrison Hybrid Comparative genomics phylogenetic analysis 12% Comparative genomics phylogenetic analysis 12% Comparative genomics phylogenetic analysis 12% Comparative genomics phylogenetic analysis 12% Comparative genomics phylogenetic analysis 12% Loss on method 10,000 STR Loss on method 10,000 STR Loss on method 10,000 STR Loss on method 10,000 STR Loss on the formation Comparative show the dipol Zygote-n Symbiosis the same number of sex chromosom Inpa Haemophilus influenza Fortizizator restress both the dipol Zygote-n Symbiosis the same number of sex chromosom Inpa Lasson of the same show the same physiology 22 chromosomes Cytogenetics hetersome Morphan 3 Amenpiohy Normal no of centromere Inversion	Is based on recombination Is have been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage manhysis? Provide mapping? Provide mapping mapping? Provide mapping mapping provide mapping mapping provide mapping mapping provide mapping mapping provide mapping mapping provide mapping prov	Is used in physical mapping Are human cell inlos Requires large number of organisms must occur in multiple alleles ddDNA ddDNA ddDNA Non-integration Non-integ	Is primarily used with prokaryotes Are produced by inradiation of UV Uilizes genomic libraries must be atleast 1000 bp long sdDNA sdDNA sdDNA sdDNA sdDNA sdDNA sdDNA sdDNA sdDNA long tasting certain organic chemicals Segregation of alleles Haptoid Human female pism sativum Lapoid atless Haptoid Human female pism sativum Law of segregation Law of sectors compositions elegans monsile, centimogram centromere, map units homologous chromosomes and alleles Fromales Fromales Physiology Han cell function of the cell H-banded CGH wendal Law of chromosome Diplication Physiology Hospiologiv Four type Four type Hospiologiv Ho	Requires linked DNA markers Can hold large pice of chromsoms Detects polymorphic DNA sequence must be atleast Software of the Circular Circular Circular Circular Circular Circular Circular andiation Hybrid formation Skb Sub Samantic Mak Drosophila Datura Law of dominance Hyman mitochordrion 1990 Carcular Law of dominance Human mitochordrion 1990 Carcular Law of dominance Human mitochordrion 1990 Carcular Law of dominance Human mitochordrion 1990 Carcular Law of dominance Human mitochordrion 1990 Carcular Law of dominance Law of dominance Law of dominance Law of dominance Law of dominance Law of dominance Carcular analysis 24 Januar	Is based on recombination Is have been used in mapping the; Uses RFLPA must be a unique DNA sequence Circular RNA Circular
Linkage analysis? Provide mapping? Provide mapping? Provide mapping? Provide mapping regions is in type? Advance vites grown vites is? Provide mapping the grown is? Provide mapping the grown is? Provide mapping the book not y Provide mapping the book not on y	Is used in physical mapping Are human cell index must occur in multiple alleles daDNA daDNA daDNA daDNA daDNA daDNA daDNA sector in multiple alleles daDNA d	Is primarily used with prokaryotes Are produced by inclusion of UV Vilizes genomic libraries subs at estables 1000 bp long subNA subNA subNA subNA subNA subNA long issific characteristic long issific characteristic long issific characteristic substration of alleles substration of alleles substrati	Requires linked DNA markers Can hold large fuce of chromsoms Detects polymorphic DNA sequence must be atleast Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Datura Mathe Drosophila Datura Mala Drosophila Datura Mala Drosophila Datura Law of dominance Human mitochondrion 1990 3.000,000.000 c comaparative genomics phylogenetic analysis 24 12% inches I od score method 1.000.00 STR a long stretch of a single repeated m 10,000 STR 10,000 STR a long stretch of a single repeated m 10,000 STR a long stretch of a single repeated m 10,000 STR a long stretch of a single repeated m 10,000 Haemophilus influenza proteome yeast inheritance of recessive genes segregation chosmoomal inference Fertilization restress both the dipoli Zygnite- nis influenza Four point crossing over 22 pair of chromosome Physiology 23 chromosomes Cytogenetics Shatterson Shanded HCH Cytogenetics Shatterson Shanded HCH Normal no of centromere Inversion One type Dipology Nationis	Is based on recombination Is heve been self in mapping the; Uses RFLPs must be a unique DNA sequence Circular RNA Circular
Link ge analysis? Provide mapping? Provide mapping Provide mapping? Provide mapping Provide mapping? Provide mapping Provide mapping Provide mapping Provide Provide Mapp	Is used in physical mapping Are human cell in loss Requires large number of organisms must occur in multiple alleles daDNA daDNA daDNA Maine Security and the alleles daDNA Maine Security and the alleles daDNA Maine Security and the alleles daDNA Maine Labyuy ocharatus Independent assortment Besterium Maine Labyuy ocharatus Independent assortment Besterium Maine Carloneromics genetic crosses 13 12,0% muicromal genetic motional security Besterium Maine Carloneromics genetic crosses 13 12,0% muicromal terference di suguare Carloneres, chromosoma in literiference di suguare Carloneres, chromosoma in literiference Carloneres, chromosoma in literiference Secoli mapping Aresonbiant DNA The same number of chromosomes as guanitative trata loci Ecoli Beh chromosoma alleles are pair Maiss Recombinant DNA The same number of chromosomes as guanitative trata loci Ecoli Secoli marca and alles are pair Maiss Ausyotogy Austing Carloneros La Jai or chromosome Ecolomosome Carloneros autoscome tastoscome Cargenson 2 Diploidy Normal no ofderononosome Delection Normal no ofderononosome Normal no ofderonono	Is primarily used with prokaryotes Are produced by inclusion of UV Uilizes genomic libraries must be atleast 1000 bp long saDNA assTNA	Requires linked DNA markers Can hold large pice of chromsoms Detects polymorphic DNA sequence must be atleased splot of the Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Circular Sub Sub Sub Sub Sub Sub Sub Sub Sub Sub	Is based on recombination Is have been self in mapping the; Uses RFLP6 must be a unique DNA sequence Circular RNA Circular

Inversion that involve the centromere are called	Paracentric inversion	Pericentric inversion	Inversion	Centromere
Chromosomal mutation is called	Duplication	Addition	Translocation	Inversion
Types of single inversions are	Two types	Only onentype	Three types	Four types
Reversal in the linear order of genes in a segment without change in the total gene content called	Delection	Translocation	Inversion	Duplication
The term of chromosomal rearrangement is called	Chromosomal deletion	Chromosomal duplication	Chromosomal inversion	Chromosomal translocation
Types of inversions are	Three type	Two types	One type	Four types
Complex inversions are	Independed inversion	Overlapping inversion	Included inversion	All the three options
Inversions occurs more than one in a chromosome is called	single inversion	pericentric inversion	paracentric inversion	complex inversion
is located in a single arm of the chromosome.	Single inversion	Pericentric inversion	Paracentric inversion	Complex inversion
inversions occurs in different regions of the chromosome.	Independed inversion	Included inversion	Overlapping inversion	Complex inversion
Missing a extra or irregular portion of chromosomal DNA is called	chromosome	chromosome normality	chromosome abnormality	standard chromosome
The deletion of chromosome containing gene and is called deficinency.	G & H	C & D	B & C	J & K
A segment of a chromosme is lost from the intercalary region it is termed as	Duplication	Deletion	Inversion	Translocation
Trisomy-21 in man causes a syndrome is called	Turne's syndrome	Klinefelter's syndrome	Down's syndrome	night blindness
A mutation in acodon leads to the substitution of one aminoacid with another what is the name of mutation	Nonsence mutation	Missence mutation	Frameshift mutation	Promoter mutation
Rearrangement of parts between nonhomologus chromosome is called	Chromosomal deletion	Chromosomal duplication	Chromosomal inversion	Chromosomal translocation
how many types of deletion?	two types	three types	four types	five types
The centromere is included in the inverted segment is called	Paracentric inversion	Pericentric inversion	Single inversion	Complex inversion
Non homologous chromosome echange segments is called	Delection	Duplication	Inversion	Translocation
In Monosomic number of chromosomes are present	2n	2n-1	2n+1	2n-2
Arises from nondisinjuction in either meiotic division as	Aneuploidy	Haploidy	Polyploidy	ploidy
Duplication:	less harmful then deletion of same siz	Generally have minimal phenotypic ef	visible in the polytene chromosome	All the three options
Chromosome were first observed by	Waldeyer	Hofrnesister	Oudet	Feulgen
Lampbrush chromosome in oocytes of Amphibia reaches up to microns.	800	300	600	200
The term genome is referred for	Haploid set of chromosome	Diploid set of chromosome	polyploid set of chromosome	Triploid set of chromosome
The inner clear zone is called	Pycnosis	Kinosome	Nucleosome	Autosome
cytochromes were discovered by	David Keilin	Mirabel	Brown	Robert brown
In trisomic number of chromsmes are present	2n	2n-1	2n+1	2n+2
The chromosome responsible for characteristic other than sex are known by which of the following terms?	Ribosome	Lysosome	Autosome	Spermatocytes
what genetic term described the situation when a part of a chromosme is broken off and lost?	Chromosomal deletion	Chromosomal duplication	Chromosomal inversion	Chromosomal translocation
Chromosomal translocation occurs in	Gametogenesis	cytogenesis	chromatin	chromosome
Chromosome abnormalities are shown in microscope the test is called	staining test	Karyotype analysis	cell counting	physiological analysis
Mutation caused disorders	Thalessemia	diphoid fever	sickle cell anemia	Brain tumour
Chromosome 4 is missing is called	Turne's syndrome	Wolf-Hischhorn syndrome	Klinefelter's syndrome	Down's syndrome
Missing of chromosome is called	Chromosomal deletion	Chromosomal duplication	Chromosomal inversion	Chromosomal translocation
Chromosome 5 is missing is called	Cri du chat syndrome	wolf-Hischhorn syndrome	Down's syndrome	Klinefeleter's syndrome
Mutations may be passed from parent to child . This substance ae called	Mutatntus	Mutagens	Mutation	genes
One of the two X chromosome in females is either missing or incomplete is called	Turne's syndrome	Wolf-Hischhorn syndrome	Klinefelter's syndrome	Down's syndrome
was first to locate duplication in drosophilla	Kaurel	bridges	Mirabel	Feulgen

Answer Pachytene Mitotic division Sexual reproduction Gregor johannmendel Garden pea 7 Garden pea Garden pea To Male and female gametes 03:01 22 Bay of segregation and independent assortment genetics 23 allele beritability dominant 1865 and 1866 1900 Taw of deminance faw of segregation independent assortment Recessive 03:01 fl berefatury diterminants Homozypeic fl berefatury diterminants Homozypeic dibybrid cross test Monohybrid 7 purple or white 7 purple or white yellow or green yellow or green Phenotype genetic disorders homologous all the three options yellow monohybrid Yellow gamete domiant homozygous hom 2 genotype looking at it o dominance incomplete dominance over dominance vytoplasmi inherinance extra chromosomal inherinance inherinance inherinance inherinance inherinance both cytoplasmi genes extra chromosomal inherinance both cytoplasmi genes extra chromosomal inherinance inherinance inherinance both cytoplasmi and extramolear inherinance inherinance inherinance both cytoplasmi and extramolear inherinance inherinance inherinance inherinance inherinance inherinance inherinance inherinance inherinance arrectina variation anternalinherinance arrectina variation anternalinherinance arrectina variation inherinance inherinance inherinance inherinance genetic inherinance inherinance arrectina variation arrectinance inherinance genetic inherinance inherinance arrectina variation arrectinance inherinance genetic inherinance inherinance arrectina variation arrectinance inherinance genetic inherinance inherinance arrectinance inherinance genetic inherinance inherinance arrectinance inherinance arrectinance inherinance analysis and therinance analysis and therinance inherinance inherinance arrectinance inherinance in active chromosine andiere inherinance inherinance arrectinance inherinance inherinance arrectinance inherinance inherinance inherinance inherinance inherinance arrectinance inherinance inherinance inherinance arrectinance inherinan Independent assortment 5 3autosomes+X+Y 2n:14 Reduction division 4 5

XX The mule all the three options: the same number of sex chromosomes as humans is on the X chromosome Xr and Y heterorygraus Genotypes AB and C 3 Alisome chick pa Banding pattern of stained chromosomes Requires that the gene have been cloned if in situ hybridization is 1 It is located at a known site on the chromosome AFLP Any changes in the NNA sequence RFLP Physical maps used to identify a specific RNA quantitative trait loci PCR based method RAPD kilobases Gene pair a segment of shared by over 99% of the population al the three options a lib thetire options a lib thetire options a signent of shared by over 99%, of the population al the three options a signent of shared by over 99%, of the population a signent of shared by over 99%, of the population a signent of shared by over 99%, of the population a signent of shared by over 99%, of the population a signent of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population fails the the options is signed and the signet by over 99%, of the population a signet of shared by over 99%, of the population a signet of shared by over 99%, of the population independent association independent associatio 46 12% centimeters Lod score method 1,000,000 RFLP a ariable number of tandem repeated nucleotide sequences 100 Haemophilus influenza proteome yeast inheritance of disorders addition all the three options chromosomes have been sequenced genetic linkage centimorgan chiasma, chromosomal interference all the three options Fernales Gene pair He same number of chromosomes and sex chromosomes quantitative trait loci Haemophilus influenza two point cross two-point cross 23 pairs of chromosome Karyology 46 chromosomes Cytogenetics Plant cell both structure and function G-banded CGH Karyotype autosome Caperson 4 4 Aneuploidy Abnormal no of chromosome Duplication Two types Diploidy All the three options Ploidy Insertion sequence

Pericentric inversion Inversion Two types Chromosomal translocation Two types All the three options Complex inversion Independed inversion Independed inversion Independed inversion Independed inversion Complex inversion Chromosoma beatomility G & H Deletion down's syndrome Missence mutation Chromosomal translocation two types Pericentric inversion Translocation 2n-1 Annapiloidy All the three options Hoffmassiare B00 Diploid Section Diploid Section Kinosome David Kellin 2n-1 Autonome Chromosomal deletion Chromosomal chromosoma chromosomal chromosomal chromosomal chromosomal chromosomal chromosomal chromosomal chromosomal chromosoma chromosomal chromosomal chromosoma chro

to be used