

**Scope:** This syllabus includes current tools for rDNA technology and its applications.

**Objective:** The application of rDNA technology ranges from developing genetically modified organisms, production of novel drugs and vaccines.

### UNIT-I

**Introduction to r-DNA technology:** Basic tools and applications – isolation and purification of nucleic acids, Enzymes used in cloning - restriction enzymes, ligases, polymerases, kinases, phosphatases. Gene recombination and gene transfer – transformation, episomes, plasmids and other cloning vectors (bacteriophage-derived, artificial chromosomes), microinjection, electroporation, ultrasonication.

### UNIT-II

**Selection and screening of recombinant clones:** Probes – radio labeled and non radio-labeled, guessmers and degenerate probes. Sequence dependent and independent screening, southern, northern hybridization, colony and plaque hybridization, *in situ* chromosomal hybridization, chromosome walking, Genome mapping, DNA fingerprinting, Polymerase chain reaction (PCR), RT- (Reverse transcription) PCR.

### UNIT-III

**Expression and characterization of cloned DNA:** Expression vectors, optimization of protein expression in heterologous systems, Fusion proteins, *In vitro* translation systems. Preparation and comparison of Genomic and cDNA library.

### UNIT-IV

**Random and site-directed mutagenesis:** Primer extension and PCR based methods of site directed mutagenesis, Random mutagenesis, Gene shuffling, production of chimeric proteins, Protein engineering concepts and examples (any two).

### UNIT-V

**Applications of Genetic Engineering:** In plants: use of *Agrobacterium tumefaciens* and *A. rhizogenes*, Ti plasmids, Strategies for gene transfer to plant cells, Direct DNA transfer to plants, Gene targeting in plants, Use of plant viruses as episomal expression vectors.

In animals: Production and applications of transgenic mice, role of ES cells in gene targeting in mice, Therapeutic products - blood proteins, human hormones, immune modulators and vaccines (one example each). Ethical, legal and social issues.

### References

1. Clark, D.P., & Pazdernik, N.J. (2009). *Biotechnology-Appling the Genetic Revolution*. USA: Elsevier Academic Press.
2. Brown, T.A., (2006). *Gene Cloning and DNA Analysis* (5th ed.). Oxford: UK, Blackwell Publishing.
3. Primrose, S.B., & Twyman, R.M. (2006). *Principles of Gene Manipulation and Genomics* (7th ed.). Oxford: UK, Blackwell Publishing.
4. Glick, B.R., & Pasternak, J.J. (2003). *Molecular Biotechnology- Principles and Applications of recombinant DNA*. Washington: ASM Press.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
(Deemed to be University Established Under Section 3 of UGC Act 1956)  
Coimbatore – 641 021.

**LECTURE PLAN**  
**Department of Biotechnology**

STAFF NAME: Dr J Anburaj

SUBJECT NAME: Recombinant DNA Technology

SEMESTER: IV

SUB.CODE:17BTU402

CLASS: II B.Sc (BT)

S.No	Lecture Duration Period	Topics to be Covered	Support Material/Page Nos
		<b>UNIT-I</b>	
1	1	isolation and purification of nucleic acids	R <sub>1</sub> . 178-180
2	1	restriction enzymes, ligases	R <sub>1</sub> . 183
3	1	polymerases, kinases, phosphatases	R <sub>1</sub> . 183-185
4	1	transformation, episomes	R <sub>1</sub> . 199
5	1	plasmids	T <sub>1</sub> . 98
6	1	bacteriophage-derived vector	R <sub>1</sub> . 205
7	1	artificial chromosomes vector	R <sub>1</sub> . 205
8	1	microinjection, electroporation, ultrasonication	T <sub>1</sub> . 69-86
	<b>Total No of Hours Planned For Unit 1=8</b>		
		<b>UNIT-II</b>	
1	1	radio labeled and non radio-labeled	T <sub>1</sub> . 69-86
2	1	guessmers and degenerate probes	R <sub>1</sub> . 13-16
3	1	Sequence dependent and independent screening	T <sub>2</sub> . 19-38
4	1	southern, northern hybridization	R <sub>1</sub> . 147-165
5	1	colony and plaque hybridization	T <sub>2</sub> . 93-100
6	1	<i>in situ</i> chromosomal hybridization, chromosome	T <sub>1</sub> . 101-103

		walking	
7	1	Genome mapping, DNA fingerprinting	T <sub>1</sub> . 101-103
8	1	Polymerase chain reaction (PCR), RT- (Reverse transcription) PCR	R <sub>1</sub> . 243
<b>Total No of Hours Planned For Unit II=8</b>			
		<b>UNIT-III</b>	
1	1	Expression vectors	T <sub>1</sub> . 243
2	1	optimization of protein expression	R <sub>1</sub> . 254
3	1	Fusion proteins	R <sub>2</sub> . 87-94
4	1	<i>In vitro</i> translation systems	R <sub>2</sub> . 87-94
5	1	Preparation and comparison of Genomic library	R <sub>2</sub> . 87-94
6	1	Preparation and comparison of cDNA library	R <sub>2</sub> . 219-221
<b>Total No of Hours Planned For Unit III=8</b>			
		<b>UNIT-IV</b>	
1	1	Primer extension and PCR based methods of site directed mutagenesis	R <sub>1</sub> . 181-185
2	1	Random mutagenesis	R <sub>1</sub> . 285-290
3	1	Gene shuffling	R <sub>1</sub> . 192-295
4	1	production of chimeric proteins	R <sub>1</sub> . 301
5	1	Protein engineering	R <sub>1</sub> . 296
<b>Total No of Hours Planned For Unit IV=8</b>			
		<b>UNIT-V</b>	
1	1	In plants: use of <i>Agrobacterium tumefaciens</i> and <i>A. rhizogenes</i>	R <sub>1</sub> . 305
2	1	Ti plasmids	R <sub>1</sub> . 305
3	1	Strategies for gene transfer to plant cells	T <sub>2</sub> . 313-348
4	1	Direct DNA transfer to plants	T <sub>2</sub> . 313-348
5	1	Gene targeting in plants	R <sub>1</sub> . 322

6	1	Use of plant viruses	R <sub>1</sub> . 318
7	1	In animals: Production and applications of transgenic mice	R <sub>2</sub> . 292-294
8	1	role of ES cells in gene targeting in mice	R <sub>2</sub> . 292-294
9	1	blood proteins	R <sub>2</sub> . 145-150
10	1	human hormones	R <sub>2</sub> . 141-143
11	1	immune modulators	R <sub>2</sub> . 128-130
12	1	vaccines	R <sub>2</sub> . 113-119
13	1	Ethical, legal and social issues	R <sub>2</sub> . 178-186
<b>Total No of Hours Planned for unit V=8</b>			
Total Planned Hours	<b>40</b>		

## REFERENCES

1. Clark, D.P., & Pazdernik, N.J. (2009). *Biotechnology-Appling the Genetic Revolution*. USA: Elsevier Academic Press.
2. Brown, T.A., (2006). *Gene Cloning and DNA Analysis* (5th ed.). Oxford: UK, Blackwell Publishing.
3. Primrose, S.B., & Twyman, R.M. (2006). *Principles of Gene Manipulation and Genomics* (7th ed.). Oxford: UK, Blackwell Publishing.
4. Glick, B.R., & Pasternak, J.J. (2003). *Molecular Biotechnology- Principles and Applications of recombinant DNA*. Washington: ASM Press.

UNIT-I

SYLLABUS

**UNIT-I**

**Introduction to r-DNA technology:** Basic tools and applications – isolation and purification of nucleic acids. Enzymes used in cloning - restriction enzymes, ligases, polymerases, kinases, phosphatases. Gene recombination and gene transfer – transformation, episomes, plasmids and other cloning vectors (bacteriophage-derived, artificial chromosomes), microinjection, electroporation, ultrasonication.

**Recombinant DNA (rDNA) molecules**

- are DNA sequences that result from the use of laboratory methods (molecular cloning) to bring together genetic material from multiple sources, creating sequences that would not be found in biological organisms.

**Recombinant DNA is possible because**

- DNA molecules from all organisms share the same chemical structure;
- they differ only in the sequence of nucleotides within that identical overall structure. Consequently, when DNA from a foreign source is linked to host sequences that can drive DNA replication and then introduced into a host organism, the foreign DNA is replicated along with the host DNA.

Recombinant DNA molecules are sometimes called **chimeric DNA**, because they are usually made of material from two different species, like the mythological chimera.

- The DNA sequences used in the construction of recombinant DNA molecules can originate from any species.
- For example, plant DNA may be joined to bacterial DNA, or human DNA may be joined with fungal DNA.
- In addition, DNA sequences that do not occur anywhere in nature may be created by the chemical synthesis of DNA, and incorporated into recombinant molecules.
- Using recombinant DNA technology and synthetic DNA, literally any DNA sequence may be created and introduced into any of a very wide range of living organisms.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: I

BATCH-2017-2021

- Proteins that result from the expression of recombinant DNA within living cells are termed recombinant proteins.
- When recombinant DNA encoding a protein is introduced into a host organism, the recombinant protein will not necessarily be produced.
- Expression of foreign proteins requires the use of specialized expression vectors and often necessitates significant restructuring of the foreign coding sequence.
- It is important to note that recombinant DNA differs from genetic recombination in that the former results from artificial methods in the test tube, while the latter is a normal biological process that results in the remixing of existing DNA sequences in essentially all organisms.

### Creating recombinant DNA

- Molecular cloning is the laboratory process used to create recombinant DNA.
- It is one of two widely-used methods (along with polymerase chain reaction, PCR) used to direct the replication of any specific DNA sequence chosen by the experimentalist.
- The fundamental difference between the two methods is that molecular cloning involves replication of the DNA within a living cell, while PCR replicates DNA in the test tube, free of living cells.
- Formation of recombinant DNA requires a cloning vector, a DNA molecule that will replicate within a living cell.
- Vectors are generally derived from plasmids or viruses, and represent relatively small segments of DNA that contain necessary genetic signals for replication, as well as additional elements for convenience in inserting foreign DNA, identifying cells that contain recombinant DNA, and, where appropriate, expressing the foreign DNA.
- The choice of vector for molecular cloning depends on the choice of host organism, the size of the DNA to be cloned, and whether and how the foreign DNA is to be expressed.
- The DNA segments can be combined by using a variety of methods, such as restriction enzyme/ligase cloning or Gibson assembly.

In standard cloning protocols, the cloning of any DNA fragment essentially involves **seven steps**:

(1) Choice of host organism and cloning vector,

- (2) Preparation of vector DNA,
- (3) Preparation of DNA to be cloned,
- (4) Creation of recombinant DNA,
- (5) Introduction of recombinant DNA into the host organism,
- (6) Selection of organisms containing recombinant DNA,
- (7) Screening for clones with desired DNA inserts and biological properties.

### **1.1. Isolation and purification of nucleic acids**

Three major types of techniques, or combinations of them, are employed in the isolation of nucleic acids:

- [1] differential solubility,
  - [2] absorption methods,
  - [3] density gradient centrifugation.
- The choice of method will depend on the type of DNA being isolated and the application.
  - A major goal of nucleic acid isolation is the removal of proteins.
  - The separation of nucleic acids from proteins is generally easily accomplished due to their different chemical properties.
  - In particular, the highly charged phosphate backbone makes the nucleic acids rather hydrophilic as compared to proteins which are more hydrophobic.

#### **Most nucleic acid isolation protocols involve a**

- [1] cell lysis step,
- [2] enzymatic treatments,
- [3] differential solubility (eg., phenol extraction or absorption to a solid support),
- [4] precipitation.

#### **[1] Cell Lysis**

- Nucleic acids must be solubilized from cells or other biological material.
- This solubilization is usually carried out under denaturing conditions such as: SDS, alkali, boiling or chaotropic agents.

- These denaturing conditions efficiently solubilize the nucleic acids and generally do not adversely affect them.
- In addition, the denaturing conditions promote the removal of proteins during the subsequent steps and inhibit the activity of nucleases which will degrade the nucleic acids.

## **[2] Enzymatic Treatment**

- Another approach in the isolation of nucleic acids is to degrade unwanted components.

For example, inclusion of proteases (eg., proteinase K) in the lysate will promote the removal of proteins.

- Proteinase K is still active at 55°C in the presence of 0.5% SDS.
- The elevated temperature and SDS improve solubility and inhibit any DNase activity that may be present in the lysate.
- Nucleases can also be used to remove unwanted nucleic acids.

For example, many DNA extraction protocols include a RNase treatment step, and visa versa.

- It is important that the RNase be free of DNase activity.

DNase-free RNase is easily prepared by boiling commercial RNase for 10 minutes.

- The stability of RNase makes the preparation of RNase-free DNase more difficult.
- RNase-free DNase should be purchased from a reliable vendor or tested before it use.

## **[3] Phenol Extraction**

- Phenol is an organic solvent that is used to separate proteins from nucleic acids.
- Proteins are hydrophobic and partition in the organic phase.
- Nucleic acids are highly charged and partition in the aqueous phase.

The advantages are of phenol extraction are that

- it is easy to carry out and can be adapted to many applications.
- It is also easily applied over a wide range of volumes (40  $\mu$ l to several ml).
- In particular, phenol extraction is widely used for the isolation of high molecular weight genomic DNA.

Phenol extraction is accomplished by mixing the sample with an equal volume of phenol which has been previously saturated with a Tris buffer at pH 8 containing EDTA and NaCl.

- The phenol should be molecular biology grade phenol should and store at -20<sup>o</sup> until preparing the saturated solution.
- The saturated solution is stored at 4<sup>o</sup>.
- Phenol is easily oxidized, as evidenced by yellowing, and the oxidation products can break DNA. Oxidized phenol should be discarded.
- Depending on the application, the two phases are completely mixed by vortexing, or gently mixed (eg., high molecular weight DNA).
- The phases are separated by centrifugation and the upper aqueous phase, which contains the nucleic acids, is retained.
- Proteins will often be visible as flocculent material at the top of the phenol phase.
- The two phases need to be carefully separated in that the nucleic acids and proteins tend to be at the interface.
- Leaving too much of the aqueous layer behind will lead to undue loss of material and aspirating too close to the interface can include protein.
- The aqueous phase can be re-extracted with phenol to remove more protein.
- Phenol is a hazardous waste material that needs to be disposed of properly.
- A common variation of phenol extraction is a mixture of phenol:chloroform: isoamyl alcohol (25:24:1).
- The more organic chloroform removes lipids, denatures more protein and mixes less with the aqueous phase leading to more efficient extraction.

#### **[4] Ethanol Precipitation**

- Nucleic acids can be precipitated from dilute solutions with ethanol.
- This precipitation can be a concentration step or a means to change buffers, especially after phenol extraction.
- Typically either sodium acetate or potassium acetate, pH 5.0-5.5, is added to a final concentration of approximately 0.3 M.

- The sodium and acidic pH will neutralize the highly charged phosphate backbone and to promote hydrophobic interactions.
- Two-to-two and a half volumes of ethanol are added and the sample is incubated as - 20°C.
- If the nucleic acids are small in size and/or in low concentrations an extended incubation (several hours to overnight) is needed.
- The precipitated DNA is collected by centrifugation.
- The pellet is rinsed with 70% ethanol to remove any excess salt, dried and dissolved in the appropriate buffer.
- A variation is to substitute ammonium acetate if the 'hard' salts are a problem.
- Another modification is to use an equal volume of isopropanol (instead of 2-2½ volumes of ethanol) which minimizes the increase in sample volume.

#### **1.1.1. Isolation of High Molecular Weight Genomic DNA**

- High molecular weight chromosomal DNA is usually isolated by multiple rounds of phenol extraction and enzyme treatments as discussed above.
- Shear forces, which can break long DNA molecules, need to be avoided during all steps and samples should never be vortexed.
- Therefore, the phenol extraction is carried with gentle rocking for several hours.
- These precautions against shear forces are not necessary in the isolation of low molecular weight DNA.
- Another common modification at the ethanol precipitation step is 'spool out' the high molecular weight genomic DNA on the end of a sealed Pasteur pipet.
- The precipitated DNA is wrapped around the end of the pipet is then allowed to partially dry and then dissolved in the appropriate buffer.
- This minimizes the contamination with RNA and low molecular weight DNA fragments.

#### **1.1.2. Plasmid Minipreps and Adsorption Methods**

- Historically, phenol extractions were used for the isolation of most forms of nucleic acids.

- It is now more common to use techniques based upon adsorption chromatography for the isolation of smaller DNA molecules, such as plasmids.
- Various kits are available for the rapid isolation of small quantities of plasmid DNA.
- The procedure consists of solubilizing the bacteria in an alkali solution followed by neutralization with sodium acetate.
- The neutralization results in the precipitation of some of the protein and the genomic DNA which is removed by centrifugation.
- The soluble material is then mixed with a resin in the presence of chaotropic agents (usually guanidine hydrochloride).
- The resins are usually either based on silica or diatomaceous earth.
- Under these conditions DNA binds to the matrix, but proteins and RNA do not.
- The DNA is eluted in a low salt buffer.
- These methods are rapid and yield a highly purified plasmid DNA which can generally be used directly in most applications without further processing.
- Another common application for an adsorption method is the isolation of DNA fragments following gel electrophoresis.
- In this case the agarose gel piece containing the DNA is dissolved in NaI, a chaotropic salt, and the DNA adsorbed to silica.
- The DNA is then eluted with a low salt buffer and sometimes gentle heating.

## **1.2. Isolation of RNA**

Most RNA isolation protocols also involve phenol extractions and are similar to DNA isolations. However, there are some differences and special considerations.

In particular,

- precautions against RNase activity must be taken.
- RNase is an extremely stable and active enzyme.
- Gloves should be worn at all times and sterile plasticware should be used whenever possible to avoid introducing exogenous RNase to the sample.
- Glassware needs to be treated with DEPC-water and autoclaved to inactivate any RNase.

- Buffers should be prepared from DEPC-water or RNase inhibitors included.
- The cell lysis and solubilization of RNA will typically be carried out in guanidine salts(especially guanidine thiocyanate).
- Guanidine is a strong chaotropic agent and will inhibit RNases.
- The strong denaturing effect of this salt will also promote better phenol extraction.
- The phenol extraction is the same as the DNA isolation except that the phenol is usually saturated with a buffer of pH 5-6.
- The lower pH will result in some DNA partitioning in the organic phase.
- If DNA contamination is a problem, it is possible to purchase RNase-free DNase.
- Long term storage is best as a precipitate in 70% ethanol at  $-20^{\circ}\text{C}$ .

### **1.2.1. Precipitation of RNA with LiCl**

- LiCl has been used to selectively precipitate RNA.
- Large RNAs (rRNA, mRNA) are insoluble at high ionic strength, whereas small RNAs (tRNA and 5S rRNA) and DNA generally remain soluble.
- Following either phenol or guanidine extraction, an equal volume of 8 M LiCl is added.
- The sample is mixed vigorously and incubated at  $-20^{\circ}\text{C}$ .
- The precipitate is collected by centrifugation and reprecipitated if necessary.

### **Affinity chromatography**

- Most eukaryotic mRNA contains a stretch of A residues at its 3' end which added post-transcriptionally.
- It is possible to isolate mRNA by affinity chromatography on oligo dT columns.
- The RNA solution is passed over an oligo-dT column under conditions which promote base pairing. Only RNA with a polyA tail binds.
- The polyA RNA is eluted under conditions (usually low salt and high temperature) which breaks the base pairing.

### **DENSITY GRADIENT CENTRIFUGATION**

- Density gradient centrifugation can also be used in the analysis and isolation of nucleic acids. Double-stranded DNA, single-stranded DNA, RNA and proteins all have different densities and therefore can be separated by isopycnic (i.e., equilibrium) centrifugation.
- CsCl is the standard medium for the density gradient centrifugation of nucleic acids and are especially useful for the purification of large amounts of highly purified DNA.
- The gradients are carried out in the presence of ethidium bromide which fluoresces when bound to DNA.
- The DNA bands are detected by illumination with ultraviolet light and easily recovered with a syringe and needle by puncturing the wall of the disposable tube and aspirating the DNA.
- The CsCl can be removed by dialysis or by precipitating the DNA.
- The %G:C content affects the density of DNA.
- This can result in multiple bands on CsCl gradients if DNA composed different G:C compositions are centrifuged.
- For example, minor bands, called satellite DNA, are often observed when total DNA from an organism is analyzed by CsCl gradients.
- These satellite bands are usually due to highly repetitive DNA or organellar DNA.
- Mitochondrial DNA is noted for its high A:T content.
- These satellite DNA bands can be purified from the genomic DNA by density gradient centrifugation.
- Nucleic acids can also be separated according to size by rate zonal centrifugation on sucrose gradients.
- However, this method is not widely used since gel electrophoresis is generally a more convenient method for the size fractionation of nucleic acids.

## **2. Enzymes used in rDNA technology**

Based on the type of reaction that they catalyze, five classes of DNA manipulative enzymes are available

1. Nucleases – enzymes that cut, shorten or degrade nucleic acid.
2. Ligases – join nucleic acid molecules together.

3. Polymerases – makes copies of nucleic acid molecules.
4. Modifying enzymes – remove or add chemical groups.
5. Topoisomerases – introduce or remove supercoils from covalently closed circular DNA.

## 2.1. Nuclease

A **nuclease** is an enzyme capable of cleaving the phosphodiester bonds between the nucleotide subunits of nucleic acids.

In the late 1960s, **Stuart Linn** and **Werner Arber** isolated examples of the two types of enzymes responsible for phage growth restriction in *Escherichia coli* (*E. coli*) bacteria.

One of these enzymes added a methyl group to the DNA, generating methylated DNA called "methylase, the other enzyme cleaves unmethylated DNA at a wide variety of locations along the length of the molecule called as "restriction nuclease". These enzymatic tools were important to scientists who were gathering the tools needed to "cut and paste" DNA molecules.

### 2.1.1. Types of nucleases

1. Structure specific nuclease
2. Sequence specific nuclease

#### Sequence specific nuclease

**H.O. Smith, K.W. Wilcox, and T.J. Kelley, in 1968**, isolated and characterized the first restriction nuclease whose functioning depended on a specific DNA nucleotide sequence. Working with *Haemophilus influenzae* bacteria, this group isolated an enzyme, called *HindII*, that always cut DNA molecules at a particular point within a specific sequence of six base pairs.

#### Types of Sequence specific nuclease

1. Exonuclease
2. Endonuclease

#### 1. Exonuclease

- A nuclease that functions by removing nucleotides from the ends of the DNA molecule is called an exonuclease.

## **2. Endonuclease**

- A restriction endonuclease functions by "scanning" the length of a DNA molecule.
- Once it encounters its particular specific recognition sequence, it will bind to the DNA molecule and makes one cut in each of the two sugar-phosphate backbones.
- The positions of these two cuts, both in relation to each other, and to the recognition sequence itself, are determined by the identity of the restriction endonuclease used to cleave the molecule in the first place.
- Different endonucleases yield different sets of cuts, but one endonuclease will always cut a particular base sequence the same way, no matter what DNA molecule it is acting on.
- Once the cuts have been made, the DNA molecule will break into fragments.

### **2.1.2. Restriction Endonucleases –**

- Endonucleases are enzymes that produce internal cuts, called cleavage, in DNA molecules. Many endonucleases cleave DNA molecules at random sites.
- But a class of endonucleases cleaves DNA only within or near those sites, which have specific base sequences; such endonucleases are known as restriction endonucleases, and the sites recognised by, them are called recognition sequences or recognition sites.
- The recognition sequences are different and specific for the different restriction endonucleases or restriction enzymes.
- Restriction enzymes were discovered due to and named after the phenomenon of host restriction of bacterial phages.

The presence restriction enzymes was postulated by **W. Arber** during 1960s, while the first true restriction endonuclease was isolated in **1970**.

**Smith, Nathans and Arber** were awarded the Nobel Prize for Physiology and Medicine in **1978** for the discovery of endonucleases.

- Restriction endonucleases are indispensable for DNA cloning and sequencing.
- They serve as the tools for cutting DNA molecules at predetermined sites, which is the basic requirement for gene cloning or recombinant DNA technology.

### **Types of Restriction Endonucleases –**

There are three distinct types of restriction endonucleases.

#### **Type I restriction endonucleases –**

- are complex endonucleases, and
- have recognition sequences of about 15 bp;
- they cleave the DNA about 1000 bp away from the 5'-end of the sequence "TCA" located within the recognition site, e.g., EcoK, EcoB etc.

#### **Type II restriction endonucleases**

- are remarkably stable and induce cleavage either, in most cases, within their recognition sequences or very close to them.
- More than 350 different type II endonucleases with over 100 different recognition sequences are known.
- They require  $Mg^{2+}$  ions for cleavage.
- The first type II enzyme to be isolated was **Hind II in 1970**.
- Only type II restriction endonucleases are used for restriction mapping and gene cloning.

#### **Type III restriction endonucleases**

- are intermediate between the type I and type II enzymes;
- they cleave DNA in the immediate vicinity of their recognition sites, e.g., EcoPI, EcoP15, Hind III etc.

### **2.1.3. Nomenclature of Restriction Endonucleases –**

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: I

BATCH-2017-2021

The nomenclature of restriction endonucleases follows a general pattern.

- (1) The first letter of the name of genus in which a given enzyme is first discovered is written in capital.
- (2) This is followed by the first two letters of species name of the organism. These three letters are generally written in italics, e.g., Eco from *Escherichia coli*, Hin from *Haemophilus influenzae*, etc.
- (3) Strain or type identification is depicted next in Roman. e.g., Ecol;.
- (4) When an organism produces more than one enzyme, they are identified by sequential Roman numerals, e.g., the different enzymes produced by *H. influenzae* strain Rd are named Hind II, Hind III, etc.

### Some Restriction Endonucleases -

Restriction endonuclease	Source (organism and strain)	Recognition sequence
AluI	<i>Arthrobacter luteus</i>	AG/CT TC/GA
BamHI	<i>Bacillus amyloliquefaciens</i> H	G/GATCCCCTAG/G
EcoRI	<i>Escherichia coli</i> Ry13	G/AATTCCTTAA/G
HindIII	<i>H. influenzae</i> Rd	A/AGCTTTTTCGA/A
Sau3A	<i>Staphylococcus aureus</i> 3A	/GATCCTAG/
TaqI	<i>Thermus aquaticus</i> YTI	T/CGAAGC/T

### Recognition Sequences For Type II Endonucleases –

- The recognition sequences for Type II endonucleases form palindromes with rotational symmetry.
- In a palindrome, the base sequence in the second half of a DNA strand is the mirror image of the sequence in its first half; consequently, the complementary DNA strand of a double helix also shows the same situation.

- But in a palindrome with rotational symmetry, the base sequence in the first half of one strand of a DNA double helix is the mirror image of the second half of its complementary strand.
- Thus in such palindromes, the base sequence in both the strands of a DNA duplex reads the same when read from the same end (either 5' or 3') of both the strands.
- Most of the type II restriction endonucleases have recognition sites of 4, 5 or 6 bp (base pairs), which are predominantly GC-rich.
- Longer palindromic target sequences are also known, and so are nonpalindromic ones (specific for some enzymes).
- Some restriction enzymes have ambiguities in their recognition sites, e.g. EcoRII, so that they may recognise upto 4 different target sequences.

#### **Cleavage Pattern of Type II Restriction Endonucleases –**

Most type II restriction endonucleases cleave the DNA molecules within their specific recognition sequences, but some produce cuts immediately outside the target sequence, e.g., NlaIII, Sau3A, etc.

#### **These cuts are either**

- (1) staggered or
  - (2) even, depending on the enzyme.
- Most enzymes produce staggered cuts in which the two strands of a DNA double helix are cleaved at different locations; this generates protruding (3'- or 5'-) ends, i.e., one strand of the double helix extends some bases beyond the other:
  - Due to the palindromic (symmetrical) nature of the target sites, the two protruding ends generated by such a cleavage by a given enzyme have complementary base sequence.
  - As a result, they readily pair with each other; such ends are called cohesive or sticky ends.

- An important consequence of this fact is that when fragments generated by a single restriction enzyme from different DNAs are mixed, they join together due to their sticky ends.
- Therefore, this property of the restriction enzymes is of great value for the construction of recombinant DNAs.
- Some restriction enzymes, on the other hand, cut both the strands of a DNA molecule at the same site so that the resulting termini or ends have blunt or flush ends in which the two strands end at the same point.
- The blunt cut ends also can be effectively utilized for construction of recombinant DNAs following one of several strategies.

### **3. DNA ligase (EC 6.5.1.1) -**

- is a specific type of enzyme that repairs single-stranded discontinuities in double stranded DNA molecules, in simple words strands that have double-strand break (a break in both complementary strands of DNA).
- Purified DNA ligase is used in gene cloning to join DNA molecules together.
- The alternative, a single-strand break, is fixed by a different type of DNA ligase using the complementary strand as a template, but still requires DNA ligase to create the final phosphodiester bond to fully repair the DNA.
- DNA ligase has applications in both DNA repair and DNA replication.
- In addition, DNA ligase has extensive use in molecular biology laboratories for Genetic recombination experiments

#### **3.1. History**

- The first DNA ligase was purified and characterized in 1967.
- The common commercially available DNA ligases were originally discovered in bacteriophage T4, *E. coli* and other bacteria.

#### **3.2. Ligase mechanism**

The mechanism of DNA ligase is to form two covalent phosphodiester bonds between 3' hydroxyl ends of one nucleotide, ("acceptor") with the 5' phosphate end of another ("donor").

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402


UNIT: I

BATCH-2017-2021

ATP is required for the ligase reaction, which proceeds in three steps:

- (1) adenylation (addition of AMP) of a residue in the active center of the enzyme, pyrophosphate is released;
- (2) transfer of the AMP to the 5' phosphate of the so-called donor, formation of a pyrophosphate bond;
- (3) formation of a phosphodiester bond between the 5' phosphate of the donor and the 3' hydroxyl of the acceptor.

Ligase will also work with blunt ends, although higher enzyme concentrations and different reaction conditions are required.



**A pictorial example of how a ligase works (with sticky ends)**

In mammals, there are **four specific types of ligase**.

- **DNA ligase I:** ligates the nascent DNA of the lagging strand after the Ribonuclease H has removed the RNA primer from the Okazaki fragments.
- **DNA ligase II:** alternatively spliced form of DNA ligase III found in non-dividing cells.
- **DNA ligase III:** complexes with DNA repair protein XRCC1 to aid in sealing DNA during the process of nucleotide excision repair and recombinant fragments.
- **DNA ligase IV:** complexes with XRCC4. It catalyzes the final step in the non-homologous end joining DNA double-strand break repair pathway. It is also required for V(D)J recombination, the process that generates diversity in immunoglobulin and T-cell receptor loci during immune system development.

### 3.3. Applications of DNA ligase in molecular biology research

- DNA ligases have become an indispensable tool in modern molecular biology research for generating recombinant DNA sequences.
- For example, DNA ligases are used with restriction enzymes to insert DNA fragments into plasmids.
- Most experiments use T4 DNA Ligase (isolated from bacteriophage T4), which is most active at 25°C.
- However, for optimal ligation efficiency with cohesive-ended fragments ("sticky ends"), the optimal enzyme temperature needs to be balanced with the melting temperature  $T_m$  (also the annealing temperature) of the sticky ends being ligated.
- If the ambient temperature exceeds  $T_m$ , the homologous pairing of the sticky ends would not be stable because the high temperature disrupts hydrogen bonding.
- Ligation reaction is most efficient when the sticky ends are already stably annealed, disruption of the annealing ends would therefore results in low ligation efficiency.
- The shorter the overhang, the lower the  $T_m$ , typically a 4-base overhang has a  $T_m$  of 12-16°C.
- Since blunt-ended DNA fragments have no cohesive ends to anneal, the melting temperature is not a factor to consider within the normal temperature range of the ligation reaction. However, the higher the temperature, the less chance that the ends to be joined will be aligned to allow ligation (molecules move around the solution more at higher

temperatures). The limiting factor in blunt end ligation is not the activity of the ligase but rather the number of alignments between DNA fragment ends that occur.

- The most efficient ligation temperature for blunt-ended DNA would therefore be the temperature at which the greatest number of alignments can occur.
- Therefore, the majority of blunt-ended ligations are carried out at 14-16°C overnight.

### 3.4. DNA polymerase

- A **DNA polymerase** is an enzyme that helps catalyze in the polymerization of deoxyribonucleotides into a DNA strand.
- DNA polymerases are best known for their feedback role in DNA replication, in which the polymerase "reads" an intact DNA strand as a template and uses it to synthesize the new strand.
- This process copies a piece of DNA.
- The newly polymerized molecule is complementary to the template strand and identical to the template's original partner strand.
- DNA polymerases use magnesium ions as cofactors.
- Human DNA polymerases are 900-1000 amino acids long.

#### 3.4.1. Function of DNA polymerase

- DNA polymerase can add free nucleotides to only the 3' end of the newly forming strand. This results in elongation of the new strand in a 5'-3' direction.
- No known DNA polymerase is able to begin a new chain (*de novo*).
- DNA polymerase can add a nucleotide onto only a preexisting 3'-OH group, and, therefore, needs a primer at which it can add the first nucleotide. Primers consist of RNA and/or DNA bases.
- In DNA replication, the first two bases are always RNA, and are synthesized by another enzyme called primase.
- An enzyme known as a helicase is required to unwind DNA from a double-strand structure to a single-strand structure to facilitate replication of each strand consistent with the semiconservative model of DNA replication.

- Error correction is a property of some, but not all, DNA polymerases.
- This process corrects mistakes in newly synthesized DNA.
- When an incorrect base pair is recognized, DNA polymerase reverses its direction by one base pair of DNA.
- The 3'-5' exonuclease activity of the enzyme allows the incorrect base pair to be excised (this activity is known as *proofreading*).
- Following base excision, the polymerase can re-insert the correct base and replication can continue.

### 3.4.2. Various DNA polymerases are extensively used in molecular biology experiments.

#### DNA Polymerase I (or Pol I)

- is an enzyme that participates in the process of DNA replication in prokaryotes.
- It is composed of 928 amino acids, and is an example of a processive enzyme - it can sequentially catalyze multiple polymerisations.
- Discovered by Arthur Kornberg in 1956,
- it was the first known DNA polymerase (and, indeed, the first known of any kind of polymerase).
- It was initially characterized in *E. coli*, although it is ubiquitous in prokaryotes.
- In *E. coli* and many other bacteria, the gene which encodes Pol I is known as *polA*.

#### Pol I possesses three enzymatic activities:

1. A 5' → 3' (forward) DNA polymerase activity, requiring a 3' primer site and a template strand
2. A 3' → 5' (reverse) exonuclease activity that mediates proofreading
3. A 5' → 3' (forward) exonuclease activity mediating nick translation during DNA repair.
4. In the replication process, DNA Polymerase I removes the RNA primer (created by Primase) from the lagging strand and fills in the necessary nucleotides between the Okazaki fragments in 5' → 3' direction, proofreading for mistakes as it goes.

5. It is a template-dependent enzyme - it only adds nucleotides that correctly base pair with an existing DNA strand acting as a template.

### **Research applications of DNA polymerase I**

1. DNA polymerase I obtained from *E. coli* is used extensively for molecular biology research.
2. However, the 5' → 3' exonuclease activity makes it unsuitable for many applications.
3. Fortunately this undesirable enzymatic activity can be simply removed from the holoenzyme to leave a useful molecule called the Klenow fragment, widely used in molecular biology.
4. Exposure of DNA polymerase I to the protease subtilisin cleaves the molecule into a smaller fragment, which retains only the DNA polymerase and proofreading activities.

### **DNA polymerase II (also known as DNA Pol II or Pol II)**

- is a prokaryotic DNA polymerase most likely involved in DNA repair.
- The enzyme is 90 kDa in size and
- is coded by the polB gene.
- DNA Pol II can synthesize DNA new base pairs at an average rate of between 40 and 50 nucleotides/second.
- Strains lacking the gene show no defect in growth or replication.
- Synthesis of **Pol II** is induced during the stationary phase of cell growth.
- This is a phase in which little growth and DNA synthesis occurs.
- It is also a phase in which the DNA can accumulate damage such as short gaps, which act as a block to DNA Pol III.
- Under these circumstances, Pol II helps to overcome the problem because it can reinitiate DNA synthesis downstream of gaps.
- Pol II has a low error rate but it is much too slow to be of any use in normal DNA synthesis.
- Pol II differs from Pol I in that it lacks a 5' → 3' exonuclease activity, and cannot use a nicked duplex template.

### DNA polymerase III holoenzyme

- is the primary enzyme complex involved in prokaryotic DNA replication.
- It was discovered by **Thomas Kornberg** and **Malcolm Gefter** in **1970**.
- The complex has high processivity (i.e. the number of nucleotides added per binding event) and, specifically referring to the replication of the *E.coli* genome, works in conjunction with four other DNA polymerases (Pol I, Pol II, Pol IV, and Pol V).
- Being the primary holoenzyme involved in replication activity,
- the DNA Pol III holoenzyme also has proofreading capabilities that correct replication mistakes by means of exonuclease activity working 3'→5'.
- DNA Pol III is a component of the replisome, which is located at the replication fork.

The replisome is composed of the following:

- **2 DNA Pol III enzymes**, each comprising  $\alpha$ ,  $\epsilon$  and  $\theta$  subunits.
  - the  $\alpha$  subunit has the polymerase activity.
  - the  $\epsilon$  subunit as 3'-5' exonuclease activity.
  - the  $\theta$  subunit stimulates the  $\epsilon$  subunit's proofreading.
- **2  $\beta$  units** which act as sliding DNA clamps, they keep the polymerase bound to the DNA.
- **2  $\tau$  units** which acts to dimerize two of the core enzymes ( $\alpha$ ,  $\epsilon$ , and  $\theta$  subunits).
- **1  $\gamma$  unit** which acts as a clamp loader for the lagging strand Okazaki fragments, helping the two  $\beta$  subunits to form a unit and bind to DNA. The  $\gamma$  unit is made up of 5  $\gamma$  subunits which include 3  $\gamma$  subunits, 1  $\delta$  subunit, and 1  $\delta'$  subunit. The  $\delta$  is involved in copying of the lagging strand.
- **X and  $\Psi$**  which form a 1:1 complex and bind to  $\gamma$  or  $\tau$ .

### Activity of DNA polymerase III

1. synthesizes base pairs at a rate of around 1000 nucleotides per second.
2. DNA Pol III activity begins after strand separation at the origin of replication.

3. Because DNA synthesis cannot start *de novo*, an RNA primer, complementary to part of the single-stranded DNA, is synthesized by primase (an RNA polymerase)

### Taq DNA polymerase

- is a thermostable DNA polymerase named after the thermophilic bacterium *Thermus aquaticus* from which it was originally isolated by **Thomas D. Brock** in **1965**.
- It is often abbreviated to "**Taq Pol**" (or simply "**Taq**"), and is frequently used in polymerase chain reaction (PCR), a method for greatly amplifying short segments of DNA.
- *T. aquaticus* is a bacterium that lives in hot springs and hydrothermal vents, and *Taq* polymerase was identified as an enzyme able to withstand the protein-denaturing conditions (high temperature) required during PCR.
- Therefore it replaced the DNA polymerase from *E. coli* originally used in PCR.
- *Taq*'s optimum temperature for activity is 75–80°C,

with a half-life of greater than 2 hours at 92.5°C, 40 minutes at 95°C and 9 minutes at 97.5°C, and can replicate a 1000 base pair strand of DNA in less than 10 seconds at 72°C.

### Taq's drawbacks

- is its relatively low replication fidelity.
- It lacks a 3' to 5' exonuclease proofreading activity, and
- has an error rate measured at about 1 in 9,000 nucleotides.
- Some thermostable DNA polymerases have been isolated from other thermophilic bacteria and archaea, such as *Pfu* DNA polymerase, possessing a proofreading activity, and are being used instead of (or in combination with) *Taq* for high-fidelity amplification.
- *Taq* makes DNA products that have A (adenine) overhangs at their 3' ends.

This may be useful in TA cloning, whereby a cloning vector (such as a plasmid) that has a T (thymine) 3' overhang is used, which complements with the A overhang of the PCR product, thus enabling ligation of the PCR product into the plasmid vector.

### 4. **RNA polymerase (RNAP or RNAPol)**

- is an enzyme that produces RNA.

- In cells, RNAP is necessary for constructing RNA chains using DNA genes as templates, a process called transcription.
- RNA polymerase enzymes are essential to life and are found in all organisms and many viruses.
- In chemical terms, RNAP is a nucleotidyl transferase that polymerizes ribonucleotides at the 3' end of an RNA transcript.

#### 4.1. History of RNA polymerase

- RNAP was discovered independently by **Sam Weiss, Audrey Stevens, and Jerard Hurwitz** in **1960**.
- The 2006 Nobel Prize in Chemistry was awarded to **Roger Kornberg** for creating detailed molecular images of RNA polymerase during various stages of the transcription process.
- RNAP accomplishes *de novo* synthesis.
- It is able to do this because specific interactions with the initiating nucleotide hold RNAP rigidly in place, facilitating chemical attack on the incoming nucleotide.
- Such specific interactions explain why RNAP prefers to start transcripts with ATP (followed by GTP, UTP, and then CTP).
- In contrast to DNA polymerase, RNAP includes helicase activity, therefore no separate enzyme is needed to unwind DNA.

#### Products of RNAP include:

- Messenger RNA (mRNA)—template for the synthesis of proteins by ribosomes.
- Non-coding RNA or "RNA genes"—a broad class of genes that encode RNA that is not translated into protein. The most prominent examples of RNA genes are transfer RNA (tRNA) and ribosomal RNA (rRNA), both of which are involved in the process of translation. However, since the late 1990s, many new RNA genes have been found, and thus RNA genes may play a much more significant role than previously thought.
  - Transfer RNA (tRNA)—transfers specific amino acids to growing polypeptide chains at the ribosomal site of protein synthesis during translation.

- Ribosomal RNA (rRNA)—a component of ribosomes.
- Micro RNA—regulates gene activity.
- Catalytic RNA (Ribozyme)—enzymatically active RNA molecules.

## 4.2. RNA polymerase in bacteria

In bacteria, the same enzyme catalyzes the synthesis of mRNA and ncRNA.

RNAP is a relatively large molecule. The core enzyme has 5 subunits (~400 kDa):

- $\alpha_2$ : The two  $\alpha$  subunits assemble the enzyme and bind regulatory factors. Each subunit has two domains:  $\alpha$ CTD (C-Terminal domain) binds the UP element of the extended promoter, and  $\alpha$ NTD (N-terminal domain) binds the rest of the polymerase. This subunit is not used on promoters without an UP element.
- $\beta$ : this has the polymerase activity (catalyzes the synthesis of RNA), which includes chain initiation and elongation.
- $\beta'$ : binds to DNA (nonspecifically).
- $\omega$ : restores denatured RNA polymerase to its functional form in vitro. It has been observed to offer a protective/chaperone function to the  $\beta'$  subunit in *Mycobacterium smegmatis*. Now known to promote assembly.<sup>[6]</sup>
- In order to bind promoter-specific regions, holoenzyme requires another subunit, sigma ( $\sigma$ ).
- The sigma factor greatly reduces the affinity of RNAP for nonspecific DNA while increasing specificity for certain promoter regions, depending on the sigma factor.
- That way, transcription is initiated at the right region.
- The complete holoenzyme therefore has 6 subunits:  $\alpha_2\beta\beta'\sigma\omega$  (~480 kDa).
- The structure of RNAP exhibits a groove with a length of 55 Å (5.5 nm) and a diameter of 25 Å (2.5 nm). This groove fits well the 20 Å (2 nm) double strand of DNA. The 55 Å (5.5 nm) length can accept 16 nucleotides.
- When not in use, RNA polymerase binds to low-affinity sites to allow rapid exchange for an active promoter site when one opens.
- RNA polymerase holoenzyme, therefore, does not freely float around in the cell when not in use.

### **4.3. RNA polymerase in eukaryotes**

Eukaryotes have several types of RNAP, characterized by the type of RNA they synthesize:

1. **RNA polymerase I** - synthesizes a pre-rRNA 45S (35S in yeast), which matures into 28S, 18S and 5.8S rRNAs which will form the major RNA sections of the ribosome.
2. **RNA polymerase II** - synthesizes precursors of mRNAs and most snRNA and microRNAs. This is the most studied type, and due to the high level of control required over transcription a range of transcription factors are required for its binding to promoters.
3. **RNA polymerase III** - synthesizes tRNAs, rRNA 5S and other small RNAs found in the nucleus and cytosol.
4. **RNA polymerase IV** - synthesizes siRNA in plants.
5. **RNA polymerase V** - synthesizes RNAs involved in siRNA-directed heterochromatin formation in plants.

There are other RNA polymerase types in mitochondria and chloroplasts. And there are RNA-dependent RNA polymerases involved in RNA interference.

### **4.4. RNA polymerase in archaea**

- Archaea have a single RNAP that is closely related to the three main eukaryotic polymerases (Pol I,II,III).
- Thus, it has been speculated that the archaeal polymerase resembles the ancestor of the specialized eukaryotic polymerases.

### **4.5. RNA polymerase in viruses**

- Many viruses also encode for RNAP.
- the most widely studied viral RNAP is found in bacteriophage T7.
- The single-subunit T7 RNA polymerase is related to that found in mitochondria and chloroplasts, and shares considerable homology to DNA polymerase.
- It is believed that most viral polymerases therefore evolved from DNA polymerase and are not directly related to the multi-subunit polymerases described above.

- The viral polymerases are diverse, and include some forms that can use RNA as a template instead of DNA.
- This occurs in negative strand RNA viruses and dsRNA viruses, both of which exist for a portion of their life cycle as double-stranded RNA.
- However, some positive strand RNA viruses, such as polio, also contain these RNA-dependent RNA polymerases.

## 5. Reverse transcriptase –

In the fields of molecular biology and biochemistry, a **reverse transcriptase**, also known as **RNA-dependent DNA polymerase**,

- is a DNA polymerase enzyme that transcribes single-stranded RNA into single-stranded DNA.
- It also is a **DNA-dependent DNA polymerase** which synthesizes the second strand of DNA of the double helix DNA once the RNA has been reverse transcribed, and degraded via its RNaseH activity, into a single strand cDNA.
- Normal transcription involves the synthesis of RNA from DNA; hence, reverse transcription is the *reverse* of this.

**Well studied reverse transcriptases include:**

- **HIV-1 reverse transcriptase** from human immunodeficiency virus type 1 (PDB 1HMY)
- **M-MLV reverse transcriptase** from the Moloney murine leukemia virus
- **AMV reverse transcriptase** from the avian myeloblastosis virus
- **Telomerase reverse transcriptase** that maintains the telomeres of eukaryotic chromosomes

### 5.1. History of reverse transcriptase

Reverse transcriptase was discovered by **Howard Temin** and independently by **David Baltimore** in **1970**. The two shared the 1975 Nobel Prize in Physiology or Medicine with Renato Dulbecco for their discovery.

## 5.2. Function of reverse transcriptase in viruses

- The enzyme is encoded and used by reverse-transcribing viruses, which use the enzyme during the process of replication.
- Reverse-transcribing RNA viruses, such as retroviruses, use the enzyme to reverse-transcribe their RNA genomes into DNA, which is then integrated into the host genome and replicated along with it.
- Reverse-transcribing DNA viruses, such as the hepadnaviruses, can allow RNA to serve as a template in assembling, and making DNA strands.
- HIV infects humans with the use of this enzyme. Without reverse transcriptase, the viral genome would not be able to incorporate into the host cell, resulting in the failure of the ability to replicate.

### 5.2.1. Process of reverse transcription

- Reverse transcriptase creates single-stranded DNA from an RNA template.
- In virus species with reverse transcriptase lacking DNA-dependent DNA polymerase activity, creation of double-stranded DNA can possibly be done by host-encoded DNA polymerase  $\delta$ , mistaking the viral DNA-RNA for a primer and synthesizing a double-stranded DNA by similar mechanism as in primer removal, where the newly synthesized DNA displaces the original RNA template.
- The process of reverse transcription is extremely error-prone and it is during this step that mutations may occur. Such mutations may cause drug resistance.

### Retroviral Reverse Transcription

- Retroviruses, also referred to as class VI ssRNA-RT viruses,
- are RNA reverse transcribing viruses with a DNA intermediate.
- Their genomes consist of two molecules of positive sense single stranded RNA with a 5' cap and 3' polyadenylated tail. Examples of retroviruses include *Human Immunodeficiency Virus* (HIV) and *Human T-Lymphotropic virus* (HTLV).

Creation of double-stranded DNA occurs in the cytosol<sup>[5]</sup> as a series of steps:

1. A specific cellular tRNA acts as a primer and hybridizes to a complementary part of the virus genome called the primer binding site or PBS
2. Complementary DNA then binds to the U5 (non-coding region) and R region (a direct repeat found at both ends of the RNA molecule) of the viral RNA
3. A domain on the reverse transcriptase enzyme called RNase H degrades the 5' end of the RNA which removes the U5 and R region
4. The primer then 'jumps' to the 3' end of the viral genome and the newly synthesised DNA strands hybridizes to the complementary R region on the RNA
5. The first strand of complementary DNA (cDNA) is extended and the majority of viral RNA is degraded by RNase H
6. Once the strand is completed, second strand synthesis is initiated from the viral RNA
7. There is then another 'jump' where the PBS from the second strand hybridizes with the complementary PBS on the first strand
8. Both strands are extended further and can be incorporated into the hosts genome by the enzyme integrase

#### **In eukaryotes**

- Self-replicating stretches of eukaryotic genomes known as retrotransposons utilize reverse transcriptase to move from one position in the genome to another via a RNA intermediate. They are found abundantly in the genomes of plants and animals.
- Telomerase is another reverse transcriptase found in many eukaryotes, including humans, which carries its own RNA template; this RNA is used as a template for DNA replication.

#### **In prokaryotes**

- Reverse transcriptases are also found in bacterial Retron msr RNAs, distinct sequences which code for reverse transcriptase,
- and are used in the synthesis of msDNA.
- In order to initiate synthesis of DNA, a primer is needed. In bacteria, the primer is synthesized during replication.

### **5.2.2. Structure of reverse transcriptase**

- Reverse transcriptase enzymes include an RNA-dependent DNA polymerase and a DNA-dependent DNA polymerase, which work together to perform transcription.
- In addition to the transcription function, retroviral reverse transcriptases have a domain belonging to the RNase H family which is vital to their replication.

### **5.3. Applications of reverse transcriptase**

#### **Antiviral drugs**

- As HIV uses reverse transcriptase to copy its genetic material and generate new viruses (part of a retrovirus proliferation circle), specific drugs have been designed to disrupt the process and thereby suppress its growth.
- Collectively, these drugs are known as reverse transcriptase inhibitors and include the nucleoside and nucleotide analogues zidovudine (trade name Retrovir), lamivudine (Epivir) and tenofovir (Viread), as well as non-nucleoside inhibitors, such as nevirapine (Viramune).

#### **Molecular biology**

- Reverse transcriptase is commonly used in research to apply the polymerase chain reaction technique to RNA in a technique called reverse transcription polymerase chain reaction (RT-PCR).
- The classical PCR technique can be applied only to DNA strands, but, with the help of reverse transcriptase, RNA can be transcribed into DNA, thus making PCR analysis of RNA molecules possible.
- Reverse transcriptase is used also to create cDNA libraries from mRNA.
- The commercial availability of reverse transcriptase greatly improved knowledge in the area of molecular biology, as, along with other enzymes, it allowed scientists to clone, sequence, and characterise DNA.
- Reverse transcriptase has also been employed in insulin production.
- By inserting eukaryotic mRNA for insulin production along with reverse transcriptase into bacteria, the mRNA can insert itself into the prokaryote's genome, and large amounts

of insulin can be created, sidestepping the need to harvest pig pancreas and other such traditional sources.

- Inserting eukaryotic DNA (instead of mRNA) into bacteria would not work because it is fragmented, with introns, and would not transcribe successfully using the bacteria's ribosomes.

## 6. Polynucleotide kinase (or PNK)

- is a T7 bacteriophage (or T4 bacteriophage) enzyme that catalyzes the transfer of a gamma-phosphate from ATP to the free hydroxyl end of the 5' DNA or RNA.
- The resulting product could be used to end-label DNA or RNA, or in a ligation reaction.

## 7. Alkaline phosphatase (ALP, ALKP) (EC 3.1.3.1)

- is a hydrolase enzyme responsible for removing phosphate groups from many types of molecules, including nucleotides, proteins, and alkaloids.
- The process of removing the phosphate group is called *dephosphorylation*.
- As the name suggests, alkaline phosphatases are most effective in an alkaline environment. It is sometimes used synonymously as **basic phosphatase**.

### In bacteria,

- alkaline phosphatase is located in the periplasmic space, external to the cell membrane.
- Since this space is much more subject to environmental variation than the actual interior of the cell, bacterial alkaline phosphatase is comparatively resistant to inactivation, denaturation, and degradation, and also has a higher rate of activity.
- The optimal pH for the activity of the *E. coli* enzyme is 8.0 while the bovine enzyme optimum pH is slightly higher at 8.5.

### Common alkaline phosphatases used in research include:

- **Shrimp alkaline phosphatase (SAP)**, from a species of Arctic shrimp (*Pandalus borealis*)

- **Calf-intestinal alkaline phosphatase (CIP)**
- **Placental alkaline phosphatase (PALP)** and its C terminally truncated version that lacks the last 24 amino acids (constituting the domain that targets for GPI membrane anchoring)
  - the secreted alkaline phosphatase (SEAP)

#### **Use in research**

- Typical use in the lab for alkaline phosphatases includes removing phosphate monoester to prevent self ligation.
- Alkaline phosphatase has become a useful tool in molecular biology laboratories, since DNA normally possesses phosphate groups on the 5' end.
- Removing these phosphates prevents the DNA from ligating (the 5' end attaching to the 3' end), thereby keeping DNA molecules linear until the next step of the process for which they are being prepared; also, removal of the phosphate groups allows radiolabeling (replacement by radioactive phosphate groups) in order to measure the presence of the labeled DNA through further steps in the process or experiment.
- Another important use of alkaline phosphatase is as a label for enzyme immunoassays.
- One common use in the dairy industry is as a marker of pasteurisation in cows' milk.
- This molecule is denatured by elevated temperatures found during pasteurisation, and can be tested for via colour change of a para-Nitrophenylphosphate substrate in a buffered solution (Aschaffenburg Mullen Test).
- Raw milk would typically produce a yellow colouration within a couple of minutes, whereas properly pasteurised milk should show no change.

#### **8. Terminal deoxynucleotidyl transferase**

**Terminal deoxynucleotidyl transferase (TdT), also known as DNA nucleotidylexotransferase (DNTE) or terminal transferase,**

- is a specialized DNA polymerase expressed in immature, pre-B, pre-T lymphoid cells, and acute lymphoblastic leukemia/lymphoma cells.
- TdT adds N-nucleotides to the V,D, and J exons during antibody gene recombination enabling the phenomenon of junctional diversity.

- In humans, terminal transferase is encoded by the *DNTT* gene.
- TdT is notably absent in fetal liver HSCs significantly impairing junctional diversity in B-cells during the fetal period.

### 8.1. Function of TdT

- TdT catalyses the addition of nucleotides to the 3' terminus of a DNA molecule.
- Unlike most DNA polymerases it does not require a template.
- The preferred substrate of this enzyme is a 3'-overhang, but it can also add nucleotides to blunt or recessed 3' ends.
- Cobalt is a necessary cofactor, however the enzyme catalyzes reaction upon Mg and Mn administration *in vitro*.

### 8.2. Uses of TdT

- Terminal transferase has applications in molecular biology.
- It can be used in RACE to add nucleotides which can then be used as a template for a primer in subsequent PCR.
- It can also be used to add nucleotides labeled with radioactive isotopes, for example in the TUNEL assay (Terminal deoxynucleotidyl transferase dUTP Nick End Labeling) for the demonstration of apoptosis (which is marked, in part, by fragmented DNA).
- Also used in the immunofluorescence assay for the diagnosis of acute lymphoblastic leukemia.
- In immunohistochemistry, antibodies to TdT can be used to demonstrate the presence of immature T and B cells and multipotent haematopoietic stem cells, which possess the antigen, while mature lymphoid cells are always TdT-negative.
- While TdT-positive cells are found in small numbers in healthy lymph nodes and tonsils, the malignant cells of acute lymphoblastic leukaemia are also TdT positive, and the antibody can therefore be used as part of a panel to diagnose this disease and to distinguish it from, for example, small cell tumours of childhood.

## **9. Deoxyribonuclease (Dnase)**

- is any enzyme that catalyzes the hydrolytic cleavage of phosphodiester linkages in the DNA backbone.
- Thus, deoxyribonucleases are one type of nuclease.
- A wide variety of deoxyribonucleases are known, which differ in their substrate specificities, chemical mechanisms, and biological functions.

### **9.1. Modes of action**

- Some DNases cleave only residues at the ends of DNA molecules (exodeoxyribonucleases, a type of exonuclease).
- Others cleave anywhere along the chain (endodeoxyribonucleases, a subset of endonucleases).
- Some are fairly indiscriminate about the DNA sequence at which they cut, while others, including restriction enzymes, are very sequence-specific.
- Some cleave only double-stranded DNA; others are specific for single-stranded molecules; and still others are active toward both.
- DNase enzymes can be inhaled using a nebuliser by cystic fibrosis sufferers.
- DNase enzymes help because white blood cells accumulate in the mucus, and, when they break down, they release DNA, which adds to the 'stickiness' of the mucus. DNase enzymes break down the DNA, and the mucus is much easier to clear from the lungs.

### **9.2. Types of deoxyribonucleases**

The two main types of DNase found in metazoans are known as

1. deoxyribonuclease I and
2. deoxyribonuclease II.

Other types of DNase include Micrococcal nuclease.

### **9.3. Assay of deoxyribonucleases**

- DNA absorbs UV light with a wavelength of maximal absorbance near 260 nm.

- This absorption is due to the pi electrons in the aromatic bases of the DNA.
- In dsDNA, or even regions of RNA where double-stranded structure occurs, the bases are stacked parallel to each other, and the overlap of the base molecular orbitals leads to a decrease in absorbance of UV light.
- This phenomenon is called the hyperchromic effect.
- When DNase liberates nucleotides from dsDNA, the bases are no longer stacked as they are in dsDNA, so that orbital overlap is minimized and UV absorbance increases.
- This increase in absorbance underlies the basis of Kunitz unit of DNase activity.
- One Kunitz unit is defined as the amount of enzyme that causes an increase in absorbance at 260 nm of 0.001 per mL when acting upon highly polymerized DNA at 25 °C and pH 5.0 under specified conditions.
- A standard enzyme preparation should be run in parallel with an unknown because standardization of DNA preparations and their degree of polymerization in solution is not possible.

## **10. Ribonuclease (RNase)**

- is a type of nuclease that catalyzes the degradation of RNA into smaller components.
- Ribonucleases can be divided into endoribonucleases and exoribonucleases,
  - comprise several sub-classes within the EC 2.7 (for the phosphorolytic enzymes) and 3.1 (for the hydrolytic enzymes) classes of enzymes.

### **10.1. Function**

- All organisms studied contain many RNases of many different classes, showing that RNA degradation is a very ancient and important process.
- As well as cleaning of cellular RNA that is no longer required,

- RNases play key roles in the maturation of all RNA molecules, both messenger RNAs that carry genetic material for making proteins, and non-coding RNAs that function in varied cellular processes.
- In addition, active RNA degradation systems are a first defense against RNA viruses, and provide the underlying machinery for more advanced cellular immune strategies such as RNAi.
- Some cells also secrete copious quantities of non-specific RNases such as A and T1.
- RNases are, therefore, extremely common, resulting in very short lifespans for any RNA that is not in a protected environment.
- It is worth noting that all intracellular RNAs are protected from RNase activity by a number of strategies including 5' end capping, 3' end polyadenylation, and folding within an RNA protein complex (ribonucleoprotein particle or RNP).
- RNases play a critical role in many biological processes, including angiogenesis and self-incompatibility in flowering plants (angiosperms).
- Also, RNases in prokaryotic toxin-antitoxin systems are proposed to function as plasmid stability loci, and as stress-response elements when present on the chromosome.
- Similar to restriction enzymes, which cleave highly specific sequences of double-stranded DNA, a variety of endoribonucleases that recognize and cleave specific sequences of single-stranded RNA have been recently classified.
- RNases play a critical role in many biological processes, including angiogenesis and self-incompatibility in flowering plants (angiosperms).
- Also, RNases in prokaryotic toxin-antitoxin systems are proposed to function as plasmid stability loci, and as stress-response elements when present on the chromosome.

## **10.2. Major types of endoribonucleases**

- **RNase A**

is an RNase that is commonly used in research.

RNase A (e.g., bovine pancreatic ribonuclease A) is one of the hardiest enzymes in common laboratory usage; one method of isolating it is to boil a crude cellular extract until all enzymes other than RNase A are denatured.

It is specific for single-stranded RNAs.

It cleaves 3' end of unpaired C and U residues, leaving a 3'-phosphorylated product, via a 2',3'-cyclic monophosphate.

- **RNase H**

is a ribonuclease that cleaves the RNA in a DNA/RNA duplex to produce ssDNA.

RNase H is a non-specific endonuclease and catalyzes the cleavage of RNA via a hydrolytic mechanism, aided by an enzyme-bound divalent metal ion. RNase H leaves a 5'-phosphorylated product.

- **RNase I**

cleaves 3'-end of ssRNA at all dinucleotide bonds leaving a 5' hydroxyl, and 3' phosphate, via a 2',3'-cyclic monophosphate intermediate.

- **RNase III**

is a type of ribonuclease that cleaves rRNA (16s rRNA and 23s rRNA) from transcribed polycistronic RNA operon in prokaryotes.

It also digests double strands RNA (dsRNS)-Dicer family of RNase, cutting pre-miRNA (60–70bp long) at a specific site and transforming it in miRNA (22–30bp), that is actively involved in the regulation of transcription and mRNA life-time.

- **RNase L**

- is an interferon-induced nuclease that, upon activation, destroys all RNA within the cell

- **RNase P**

- is a type of ribonuclease that is unique in that it is a ribozyme – a ribonucleic acid that acts as a catalyst in the same way as an enzyme.
- Its function is to cleave off an extra, or precursor, sequence on tRNA molecules.
- RNase P is one of two known multiple turnover ribozymes in nature (the other being the ribosome).

- A form of RNase P that is a protein and does not contain RNA has recently been discovered.
- **RNase PhyM**
  - is sequence specific for single-stranded RNAs. It cleaves 3'-end of unpaired A and U residues.
- **RNase T1**
  - is sequence specific for single-stranded RNAs.
  - It cleaves 3'-end of unpaired G residues.
- **RNase T2**
  - is sequence specific for single-stranded RNAs.
  - It cleaves 3'-end of all 4 residues, but preferentially 3'-end of As.
- 
- **RNase U2**
  - is sequence specific for single-stranded RNAs.
  - It cleaves 3'-end of unpaired A residues.
- **RNase V1**
  - is non-sequence specific for double-stranded RNAs.
  - It cleaves base-paired nucleotide residues.
- **RNase V**

### 10.3. Major types of exoribonucleases

- **Polynucleotide Phosphorylase (PNPase)** functions as an exonuclease as well as a nucleotidyltransferase.
- **RNase PH** functions as an exonuclease as well as a nucleotidyltransferase.
- **RNase II** is responsible for the processive 3'-to-5' degradation of single-stranded RNA.
- **RNase R** is a close homolog of RNase II, but it can, unlike RNase II, degrade RNA with secondary structures without help of accessory factors.
- **RNase D** is involved in the 3'-to-5' processing of pre-tRNAs.
- **RNase T** is the major contributor for the 3'-to-5' maturation of many stable RNAs.
- **Oligoribonuclease** degrades short oligonucleotides to mononucleotides.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: I

BATCH-2017-2021

- **Exoribonuclease I** degrades single-stranded RNA from 5'-to-3', exists only in eukaryotes.
- **Exoribonuclease II** is a close homolog of Exoribonuclease I.

### Reference

Unit topics	Book Reference	Page Number
<b>Unit – I</b>		
Nucleases		52 to 56
Restriction endonucleases		
DNA Ligase		56 to 58
DNA polymerase I, II, and III	Gene cloning – An	57 to 58
<i>Taq</i> DNA polymerase	Introduction, Third	
RNA polymerase	edition, T.A. Brown,	
Polynucleotide kinase	1998	59
Alkaline phosphatase		
Terminal deoxynucleotidyl transferase		
Isolation and purification of nucleic acids		27 to 51

UNIT-II

SYLLABUS

**UNIT-II**

**Selection and screening of recombinant clones:** Probes – radio labeled and non radio-labeled, guessmers and degenerate probes. Sequence dependent and independent screening, southern, northern hybridization, colony and plaque hybridization, *in situ* chromosomal hybridization, chromosome walking, Genome mapping, DNA fingerprinting, Polymerase chain reaction (PCR), RT- (Reverse transcription) PCR.

**Cloning Vectors**

- A cloning vector is a DNA molecule that has the ability to replicate in an appropriate host cell, and into which the DNA insert is integrated for cloning.
- Therefore, a vector must have an origin of DNA replication (denoted as ori) that functions in the host cell.
- Any extrachromosomal small genome, e.g., plasmid, phage and virus, may be used as a vector.

**1.1. Properties of Good Vector**

1. It should be able to replicate autonomously. When the objective of cloning is to obtain a large number of copies of the DNA insert, the vector replication must be under relaxed control so that it can generate multiple copies of itself in a single host cell.
  2. It should be easy to isolate and purify.
  3. It should be easily introduced into the host cells, Le., transformation of the host with the vector should be easy
  4. The vector should have suitable marker genes that allow easy detection and/or selection of the transformed host cells.
  5. When the objective is gene transfer, it should have the ability to integrate either itself or the DNA insert it carries into the genome of the host cell.
- . The cells transformed with the vector containing the DNA insert (recombinant DNA) should be identifiable be selectable from those transformed by the unaltered vector.

7. A vector should contain unique target sites for as many restriction enzymes as possible into which the DNA insert can be integrated.

8. When expression of the DNA insert is desired, the vector should contain at least suitable control elements, e.g., promoter, operator and ribosome binding sites.

It should be kept in mind that

- (1) the DNA molecules used as vectors have coevolved with their specific natural host species, and hence are adapted to function well in them and in their closely related species. Therefore, the choice of vector depends largely on the host species into which the DNA insert of gene is to be cloned.
- (2) most naturally occurring vectors do not have all the required functions; therefore, useful vectors have been created by joining together segments performing specific functions (called modules) from two or more natural entities.

## 1.2 Cloning and Expression Vectors

All vectors used for propagation of DNA inserts in a suitable host are called **cloning vectors**. But when a vector is designed for the expression of, i.e., production of the protein specified by, the DNA insert, it is termed as **expression vector**.

- As a rule, such vectors contain at least the regulatory sequences, i.e., promoters, operators, ribosomal binding sites, etc., having optimum function in the chosen host.
- When an eukaryotic gene is to be expressed in a prokaryote, the eukaryotic coding sequence has to be placed after prokaryotic promoter and ribosome binding site since the regulatory sequences of eukaryotes are not recognised in prokaryotes.
- In addition, eukaryotic genes, as a rule, contain introns (noncoding regions) present within their coding regions.
- These introns must be removed from the DNA insert to enable the proper expression of eukaryotic genes since prokaryotes lack the machinery needed for their removal from the RNA transcripts.
- When eukaryotic genes are isolated as cDNA, they are intron-free and, hence, suitable for expression in prokaryotes.

### 1.3. *E.coli* vectors

- Bacteria are the hosts of choice for DNA cloning.
- Among them, *E. coli* occupies a prominent position since cloning and isolating DNA inserts for structural analysis is the easiest in this host.
- Therefore, the initial cloning experiments are generally carried out in *E. coli*.
- The *E. coli* strain K12 is the most commonly used;
- it has several substrains, e.g., C600, RRI, HB101, etc., each of which has some specific features important in cloning.
- For example, the substrain RRI has, in addition to certain other features, the mutation *hsdR*, which inactivates the restriction enzyme endogenous to *E.coli* K12; this minimises the degradation of recombinant DNA introduced into it.

### 1.4. Properties of Good Host

A good host should have the following features:

- (1) be easy to transform,
- (2) support the replication of recombinant DNA,
- (3) be free from elements that interfere with replication of recombinant DNA,
- (4) lack active restriction enzymes, e.g., *E. coli* K12 substrain HB101,
- (5) should not have methylases since these enzymes would methylate the replicated recombinant DNA. which, as a result, would become resistant to useful restriction enzymes, and
- (6) be deficient in normal recombination function so that the DNA insert is not altered by recombination events.

***E. coli* supports several types of vectors, which can be grouped as follows:**

- (1) plasmids,
- (2) bacteriophages (both natural),
- (3) cosmids,
- (4) phasmids,
- (5) shuttle vectors

## 2. Plasmid vector

- Plasmids are widely used as cloning vector,
- Plasmids are replicons which are stably inherited in an extrachromosomal state.
- Most plasmids exist as double-stranded circular DNA molecules.
- If both strands of DNA are intact circles the molecules are described as covalently closed circles or CCC DNA.
- If only one strand is intact, then the molecules are described as open circles or OC DNA.
- When isolated from cells, covalently closed circles often have a deficiency of turns in the double helix,
- such that they have a supercoiled configuration.
- Plasmids are widely distributed throughout the prokaryotes, vary in size from less than  $1 \times 10^6$  daltons to greater than  $200 \times 10^6$ , and are generally dispensable.
- Plasmids to which phenotypic traits have not yet been ascribed are called *cryptic* plasmids.

Plasmids can be categorized into one of two major type – depending upon whether or not they carry a set of transfer genes, called the *tra* genes, which promote bacterial conjugation.

1. conjugative
2. non-conjugative

Plasmids can also be categorized on the basis of copies per cell being maintained as

1. *relaxed* plasmids - multiple copies per cell
2. *stringent* plasmids - a limited number of copies per cell

**pBR322** - An ideal plasmid vector must have the following functions:

- (1) minimum amount of DNA,
- (2) relaxed replication control,
- (3) at least two selectable markers,
- (4) only one (unique) recognition site for at least one restriction endonuclease, and

- (5) for easy selection of the recombinant DNA, this unique restriction site must be located within one of the two selectable markers.

The name pBR denotes the following:

p signifies plasmid,

B is from Boliver, and


R is from Rodriguez, the two initials of the scientist who developed pBR322.

### **pBR322**

- is the most popular and
- most widely used plasmid of 4363 bp;
- It has the replication module of *E. coli* plasmid Col El.
- This module has been incorporated in many other plasmid vectors since it permits plasmid replication even when chromosome replication and cell division are inhibited by amino acid starvation or chloramphenicol.
- Under such conditions, each cell accumulates several thousand copies of the plasmid so that one litre of bacterial culture easily yields a milligram of plasmid DNA.
- It has two selectable markers (tetracycline, tetr, and ampicillin, amp', resistance genes), and unique recognition sites for 12 different restriction enzymes (two unique sites, PstI and PvuI, are located within the amp' gene, and 4, e.g., BamHI, SalI, etc., are within tetr gene).

The presence of restriction sites within the markers tetr and ampr permits an easy selection for cells transformed with the recombinant pBR322.

- Insertion of the DNA fragment into the plasmid using restriction enzyme PstI or PvuI places the DNA insert within the gene amp'; this makes amp' nonfunctional.
- Bacterial cells containing such a recombinant pBR322 will be unable to grow in the presence of ampicillin, but will grow on tetracycline.
- Similarly, when restriction enzyme BamHI or SalI is used, the DNA insert is placed within the gene tetr making it nonfunctional.



- Bacterial cells possessing such a recombinant pBR322 will, therefore, grow on ampicillin but not on tetracycline.
- This feature allows an easy selection of a single bacterial cell having recombinant pBR322 from among 10<sup>8</sup> other types of cells.
- Transformed *E. coli* cells are first plated on an agar medium containing the antibiotic within the resistance gene for which the DNA fragment is not inserted, i.e., for which the bacterial cells having the recombinant DNA are expected to be resistant.
- This eliminates nontransformed bacterial cells; the resulting bacterial colonies will possess either recombinant or unaltered pBR322.
- The colonies so obtained are then replicaplated on agar plates containing the other antibiotic (within the resistance gene for which the DNA insert is placed);
- all the colonies that develop on this plate will contain the unaltered pBR322.
- Therefore, the antibiotic sensitive colonies are identified and recovered from the master plate; these colonies will have the recombinant pBR322. This entire process may take up to 2 days.

**pUC18-19 –**

- pUC gets its name from University of California.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT


COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: II

BATCH-2017-2021

- It is a derivative of pBR322 and is much smaller (-2.7 kb);
- it has all the essential parts of pBR322, e.g.,
  - (1) ampicillin resistance gene and
  - (2) Col E1 origin.
  - (3) The second selectable marker is due to *E. coli* gene *lacZα* encoding the a fragment of  $\beta$ -galactosidase, the enzyme that hydrolyses lactose.
  - (4) The *E. coli* strains, e.g., JM103, JM109, used as hosts for the pUC series vectors have the *lacZα* deleted from their *lacZ* genes.



- When pUC enters such an *E. coli* cell, the host genome and the plasmid encode for different parts of the  $\beta$ -galactosidase enzyme, which interact with each other to produce the active enzyme enabling these cells to hydrolyse lactose.  $\beta$ -galactosidase also hydrolyses X-gal (5-Bromo-4-chloro-3-indolyl-p-D-galactoside) to yield a blue dye.
- Therefore appropriate *lacZ<sup>-</sup>* *E. coli* cells transformed by the pUC vectors behave as *lacZ<sup>+</sup>* and produce blue coloured colonies on a X-gal containing medium.
- A poly linker sequence located within the *lacZα* provides several (10 in case of pUC18/pUC19) unique restriction sites for DNA insertion.
- The polylinker sequence by itself does not interfere with *lacZα* expression.
- But when a DNA insert is placed within it, *lacZα* expression is prevented.
- Vectors pUC18 and pUC19 are identical, except for the orientation of the polylinker sequence, which is oriented in the opposite directions in the two vectors.
- The unique restriction sites used for integration of DNA inserts into pUC vectors interrupt the *lacZα* fragment so that appropriate

- E. coli cells possessing recombinant pUC DNA are  $\beta$ -galactosidase deficient and, as a result,
  - produce white colonies on X-gal medium.
  - Therefore, appropriate E.coli cells transformed with pUC recombinant DNA are grown on ampicillin, X-gal and IPG (isopropyl-  $\beta$  D-thiogalactoside; it serves as inducer of  $\beta$  -galactosidase, while X-gal itself can not) containing medium to eliminate non transformed cells.
- The white colonies are selected as they contain the recombinant DNA (in contrast, blue colonies will contain the unaltered vector). The other vectors in pUC series are pUC 8, pUC 9, pUC 12, pUC 13, etc.

**The pUC series vectors offer the following advantages over pBR322:**

- (1) each E. coli cell produces up to 700 copies without any treatment,
- (2) cells containing recombinant DNA are selected in a single step,
- (3) the sites for DNA insert integration are confined to the poly linker, which permits the use of two restriction enzymes to open the vector, and
- (4) they also allow sequencing of the DNA insert.

### **3. Bacteriophage Vectors**

- Bacteriophages are viruses that attack bacteria.
- Most phages lyse the bacterial cells they infect (lytic phages).
- But many others can choose to follow either a lytic or a lysogenic cycle; in the latter situation, the phage chromosome integrates into the bacterial chromosome and multiplies with the latter as prophage (temperate or lysogenic phages).
- The prophage may dissociate from the bacterial chromosome and follow the lytic cycle.

Several bacteriophages are used as cloning vectors,

- the most commonly used E. coli phages being  $\lambda$  (lambda) and M13 phages.

Plasmid vectors have to be introduced into bacterial cells, which are then cloned and selected for the recovery of recombinant DNA.

In contrast, the phage vectors are directly tested on an appropriate bacterial lawn (a continuous bacterial growth on an agar plate) where each phage particle forms a plaque (a clear bacteria-free zone in the bacterial lawn).

**Phage vectors present two advantages over plasmid vectors.**

- (1) They are more efficient than plasmids for cloning of large DNA fragments; the largest cloned insert size in a  $\lambda$  vector is just over 24 kb, while that for plasmid vectors it is less than 15 kb. In addition,
- (2) it is easier to screen a large number of phage plaques than bacterial colonies for the identification of recombinant plaques/clones.

**Lambda ( $\lambda$ ) Phage Vectors –**

- The  $\lambda$  genome (total 48,502 bp) contains an
  - origin of replication,
  - genes for head and tail proteins and
  - enzymes for DNA replication, lysis and lysogeny, and
  - single-stranded protruding cohesive ends of 12 bases (5' GGGCGGCGACCT; the other end is complementary to it, i.e., CCCGCCGCTGGA 5').
  - The  $\lambda$  genome remains linear in the phage head, but within E. coli cells the two cohesive ends anneal to form a circular molecule necessary for replication.
  - The sealed cohesive ends are called cos sites, which are the sites of cleavage during and are necessary for packaging of the mature phage DNA into phage heads.
  - The  $\lambda$  DNA must be larger than 38 kb and smaller than 52 kb to be packaged into phage particles.
  - The genes for lysogeny are located in the segment between 20 and 38 kb; the whole or a part of this segment is deleted to create  $\lambda$  vectors to
- (1) accommodate larger DNA inserts and
  - (2) to ensure that the recombinant phage is always lytic.


Several vectors were produced from wild type  $\lambda$  genome by mutation and recombination in vivo as well as by recombinant DNA techniques.

**These vectors have the following two basic features.**

- (1) The vector itself can be propagated as phage in *E. coli* cells enabling preparation of vector DNA.
- (2) They contain restriction sites, which allow the removal of the lysogenic segment and also provide insertion site for the DNA fragment.
- (3) During annealing and ligation of the DNA insert with the  $\lambda$  vector, two or more recombinant DNAs may join end-to-end producing a concatemer, which is the proper precursor for packaging of  $\lambda$  genome into phage heads.

#### **Phage M13 Vectors –**

- These vectors are used for obtaining single-strand copies of cloned DNA, which are especially suited for DNA sequencing.
- They are derived from the 6.4 kb genome of the *E. coli* filamentous bacteriophage M13.
- This phage has a single-stranded linear DNA genome in phage particles, which converts into a double stranded circular replicative intermediate within the host cells.
- M13 infects only F<sup>+</sup> cells; it does not kill the cells, but forms turbid plaques due to growth retardation of infected cells.
- Ordinarily, the double stranded form is used to produce recombinant molecules since single-stranded DNAs are not cleaved by type II restriction endonucleases; this form is readily isolated from M13-infected *E. coli* cells.



- However, the single-stranded form of M13 is used to

recover single-stranded copies of the DNA inserts; this form of vector is available from the phage particles abundant in the growth medium.

### Properties of M13 Vectors –

M13 genome has been used to produce M13mp series of vectors, e.g., M13mp 8, M13mp 9, etc.

The desirable features of M13 vectors are as follows:

- (1) very large inserts can be cloned since packaging does not depend on genome size (as is the case with  $\lambda$  vectors).
- (2) Pure single-strand copies of double-strand DNA inserts are obtained in abundance.
- (3) Since DNA inserts are accepted in either of the orientations (which is also the case for plasmid and  $\lambda$  vectors), some recombinant clones will produce single-strand copies of one strand of the DNA double-strand, while others would produce copies of the complementary strand of the DNA insert.

The phage particles in a single plaque, as a rule, will yield copies of the same single-strand. This property is very useful for a precise DNA sequencing (using both the strands of a DNA molecule) and for the synthesis of specific radio-labelled DNA probes.

- (4) Bacterial cells infected by these vectors remain viable as in the case of plasmid vectors; this allows easy maintenance of the vector. Finally,
- (5) they form plaques like  $\lambda$  phage vectors making selection of the recombinant DNAs rather easy, and
- (6) the recombinant DNA is obtained within stable bacteriophage particles.

### Two types of $\lambda$ phage vectors

1. Insertional vectors
2. Replacement vectors


**The  $\lambda$  insertional vectors** – accept less DNA than the replacement type, the foreign DNA is simply inserted into a region of the phage genome with appropriate restriction sites.

Example -  $\lambda$ gt10,  $\lambda$ charon16A

In  $\lambda$  replacement vectors – a central region of DNA not essential for lytic growth is removed. This creates two DNA fragments, called right and left arms. The central stuffer fragment is replaced by inserting foreign DNA between the arms to form a functional recombinant  $\lambda$  phage.

Example –  $\lambda$ embl,  $\lambda$ ZAP

*General schemes used for cloning in  $\lambda$  insertion and  $\lambda$  replacement vectors*




#### 4. Cosmid Vectors

Cosmids are essentially plasmids that contain a minimum of 250 bp of  $\lambda$  DNA, which includes

- (1) the cos site (the sequence yielding cohesive ends) and
- (2) sequences needed for binding of and cleavage by terminase so that under appropriate conditions they are packaged in vitro into empty  $\lambda$  phage particles.

A typical cosmid has

- (1) replication origin,
  - (2) unique restriction sites and
  - (3) selectable markers from the plasmid; therefore, selection strategy for obtaining the recombinant DNA is based on that for the contributing plasmid.
- Cosmid vectors are constructed using recombinant DNA techniques.
  - The cosmid vectors are opened by the appropriate restriction enzyme at a unique site, are then mixed with DNA inserts prepared by using the same enzyme and annealed.
  - Among the several types of products, long concatemers are present, which are the appropriate precursors for packaging in  $\lambda$  particles.
  - This procedure selects for long DNA inserts since for packaging the distance between two cos sites must be between 38 and 52 kb.
  - Cosmids can accommodate upto 45 kb long DNA inserts.
  - Packaged cosmids infect host cells like  $\lambda$  particles, but once inside the host they replicate and propagate like plasmids



**The typical features of cosmids areas follows:**

- (1) they can be used to clone, DNA inserts of upto 45 kb.
- (2) They can be packaged into  $\lambda$  particles that infect host cells, which is many-fold more efficient than plasmid transformation.
- (3) Selection for recombinant DNA is based on the procedure applicable to the plasmid making up the cosmid.

- (4) Finally, these vectors are amplified and maintained in the same manner as the contributing plasmid.


## **5. Phasmid Vectors**

- These vectors are shortened linear  $\lambda$  genomes containing DNA replication and lytic functions plus the cohesive ends of the phage; their middle nonessential segment is replaced by a linearized plasmid with intact replication module.
- In practice, a plasmid vector contains several tandem copies of the plasmid to make it longer than 38 kb, the minimum size needed for packaging in  $\lambda$  particles.
- During construction of the recombinant DNA, one or more copies of the plasmid are deleted from and the DNA insert is integrated into the vector, but generally one copy of the plasmid is retained in the recombinant DNA.
- Phasmids, both recombinant and unaltered, are packaged in  $\lambda$  particles in vitro and used for infection of appropriate *E. coli* cells.
- If a plasmid lacks the  $\lambda$  gene *el*, which produces the lysis repressor, it multiplies like a phage and produces plaques on a bacterial lawn. But if *el* gene is present, the plasmid replicates like a plasmid.
- Further, a plasmid may contain a mutant *cI* gene, which produces a temperature sensitive CI protein (inactive at higher temperatures); such vectors replicate as plasmids at lower temperatures, but behave like phage at higher temperatures. This feature is quite useful in some experiments.

## **6. Vectors For Plants**

- Plants cells do not contain any plasmid.
- But two plasmids, called pTi and pRi, and present in the bacteria *Agrobacterium tumefaciens* and *A. rhizogenes*, respectively, provide a naturally occurring transformation system.
- These plasmids transfer a part of their DNA, called T-DNA, into the genomes of most dicot and some monocot plants.
- These plasmids, especially the Ti plasmid, have been used to develop a variety of vectors.

- In addition, genomes of many plant viruses are being developed as vectors.



- The purpose of plant vectors is almost always a stable transformation ordinarily in the form of integration in plant genomes.
- But in the case of virus vectors, the objective is to produce large quantities of the protein encoded by the DNA insert.

## 7. Animal viral vectors

### 7.1. Retroviruses

- are one of the maintype of current gene therapy approaches.
- The recombinant retroviruses such as the Moloney murine leukemia virus have the ability to integrate into the host genome in a stable fashion.
- They contain a reverse transcriptase that allows integration into the host genome.
- They have been used in a number of FDA-approved clinical trials such as the SCID-X1 trial.
- Retroviral vectors can either be replication-competent or replication-defective.

- Replication-defective vectors are the most common choice in studies because the viruses have had the coding regions for the genes necessary for additional rounds of virion replication and packaging replaced with other genes, or deleted.
- These virus are capable of infecting their target cells and delivering their viral payload, but then fail to continue the typical lytic pathway that leads to cell lysis and death.
- Conversely, replication-competent viral vectors contain all necessary genes for virion synthesis, and continue to propagate themselves once infection occurs.
- Because the viral genome for these vectors is much lengthier, the length of the actual inserted gene of interest is limited compared to the possible length of the insert for replication-defective vectors. Depending on the viral vector, the typical maximum length of an allowable DNA insert in a replication-defective viral vector is usually about 8–10 kB.

**Drawback to use of retroviruses** the Moloney retrovirus involves the requirement for cells to be actively dividing for transduction.

- As a result, cells such as neurons are very resistant to infection and transduction by retroviruses.
- There is concern that insertional mutagenesis due to integration into the host genome might lead to cancer or leukemia.
- This concern remained theoretical until gene therapy for ten SCID-X1 patients using Maloney murine leukemia virus resulted in two cases of leukemia caused by activation of the LMO2 oncogene due to nearby integration of the vector.

## 7.2 Shuttle Vectors

- These vectors have been designed to replicate in cells of two different species; therefore, they contain two origins of replication, one specific for each host species, as well as those genes necessary for their replication and not provided by the host cells.
- These vectors are created by recombinant techniques.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: II

BATCH-2017-2021

- Some of them can be grown in two different prokaryotic species, while others can propagate in a prokaryotic species, usually *E. coli*, and a eukaryotic one, e.g. yeast, plants, animals.
- Since these vectors can be grown in one host and then moved into another without any extra manipulation, they are called shuttle vectors.

A shuttle vector designed to replicate in *E. coli* and *Streptomyces* has been constructed as follows:

(1) the modules for DNA replication in *Streptomyces* and methylenomycin A resistance are derived from a streptomyces plasmid, and

(2) the replication module for maintenance in *E. coli* and a gene for antibiotic resistance are taken from an *E. coli* plasmid.

- This shuttle vector allows the initial cloning of *Streptomyces* DNA inserts in *E. coli* and their subsequent functional tests in *Streptomyces*.
- Shuttle vectors have been designed to specifically satisfy this need, i.e., the initial cloning of DNA inserts in *E. coli* and subsequent functional tests in the species to which the DNA inserts belong.

Most of the eukaryotic vectors are, in fact, shuttle vectors.

### UNIT-III

### SYLLABUS

#### **UNIT-III**


**Expression and characterization of cloned DNA:** Expression vectors, optimization of protein expression in heterologous systems, Fusion proteins, *In vitro* translation systems. Preparation and comparison of Genomic and cDNA library.

## 1. Cloning strategies

Any cell-based cloning procedure has four essential parts:

- a method for generating the DNA fragment for cloning;
- a reaction that inserts that fragment into the chosen cloning vector;
- a means for introducing that recombinant vector into a host cell wherein it is replicated; and
- a method for selecting recipient cells that have acquired the recombinant

### Generalized overview of cloning strategies



### Linkers

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT


COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III


BATCH-2017-2021

- Blunt ligation is most usefully applied to joining blunt-ended fragments via *linker* molecules; in an early example of this,
- **Scheller *et al.* (1977)** synthesized self-complementary decameric oligonucleotides, which contain sites for one or more restriction endonucleases.
- The molecule can be ligated to both ends of the foreign DNA to be cloned, and then treated with restriction endonuclease to produce a sticky-ended fragment, which can be incorporated into a vector molecule that has been cut with the same restriction endonuclease.
- Insertion by means of the linker creates restriction enzyme target sites at each end of the foreign DNA
- and so enables the foreign DNA to be excised and recovered after cloning and amplification in the host bacterium.



## 2. Adaptors

- It may be the case that the restriction enzyme used to generate the cohesive ends in the linker will also cut the foreign DNA at internal sites.
- In this situation the foreign DNA will be cloned as two or more subfragments.
- One solution to this problem is to choose another restriction enzyme, but there may not be a suitable choice if the foreign DNA is large and has sites for several restriction enzymes.
- Another solution is to methylate internal restriction sites with the appropriate modification methylase. Alternatively, a general solution to the problem is provided by chemically synthesized adaptor molecules which have a *preformed* cohesive end.
- Consider a blunt-ended foreign DNA containing an internal *Bam*HI site (Figure), which is to be cloned in a *Bam*HI-cut vector.
- The *Bam* adaptor molecule has one blunt end bearing a 5' phosphate group and a *Bam* cohesive end which is not phosphorylated.
- The adaptor can be ligated to the foreign DNA ends.
- The foreign DNA plus added adaptors is then phosphorylated at the 5' termini and ligated into the *Bam*HI site of the vector.
- If the foreign DNA were to be recovered from the recombinant with *Bam*HI, it would be obtained in two fragments.
- However, the adaptor is designed to contain two other restriction sites (*Sma*I, *Hpa*II), which may enable the foreign DNA to be recovered intact.
- Note that the only difference between an adaptor and a linker is that the former has cohesive ends and
- the latter has blunt ends.




### 3. Homopolymer tailing

- A general method for joining DNA molecules makes use of the annealing of complementary homopolymer sequences.
- Thus, by adding oligo(dA) sequences to the 3' ends of one population of DNA molecules and oligo(dT) blocks to the 3' ends of another population, the two types of molecule can anneal to form mixed dimeric circles (Figure).
- An enzyme purified from calf thymus, terminal deoxynucleotidyltransferase, provides the means by which the homopolymeric extensions can be synthesized, for if presented with a single deoxynucleotide triphosphate it will repeatedly add nucleotides to the 3' OH termini of a population of DNA molecules.
- DNA with exposed 3' OH groups, such as arise from pretreatment with phage  $\lambda$  exonuclease or restriction with an enzyme such as *Pst*I, is a very good substrate for the transferase.
- Typically, 10–40 homopolymeric residues are added to each end.

One of the earliest examples of the construction of recombinant molecules,

- the insertion of a piece of  $\lambda$  DNA into SV40 viral DNA, made use of homopolymer tailing.
- In their experiments, the single-stranded gaps which remained in the two strands at each join were repaired *in vitro* with DNA polymerase and DNA ligase so as to produce



covalently closed circular molecules.

- The recombinants were then transfected into susceptible mammalian cells.
- Subsequently, the homopolymer method, using either dA.dT or dG.dC homopolymers was used extensively to construct recombinant plasmids for cloning in *E. coli*.

#### 4. Cloning of genomic DNA

There are **two major strategies** for isolating sequences from complex sources such as genomic DNA.

The **first strategy** is

a cell-based cloning strategy, is to divide the source DNA into manageable fragments and *clone everything*.

Such a collection of clones, representative of the entire starting population, is known as a *library*. Then *screen the library* to identify our clone of interest, using a procedure that discriminates between the desired clone and all the others.

The **second strategy** is

to selectively amplify the target sequence directly from the source DNA using the polymerase chain reaction (PCR), and then clone this individual fragment.

### 5.1. Genomic Library

- A genomic library is a collection of plasmid clones or phage lysates containing recombinant DNA molecules so that the sum total of DNA inserts in this collection, ideally, represents the entire genome of the concerned organism. However, inspite of all the care taken in the production of genomic libraries.

Certain DNA fragments should be expected to be under or over represented or even missing. There are several possible reasons for this, and at present they can not be taken care of.

### 5.2. Construction of A Genomic Library

- For preparation of a genomic library, the total genomic DNA of an organism is extracted.
- The DNA is broken into fragments of appropriate size either by mechanical shearing (this generates blunt ended fragments), sonication, or by using a suitable restriction endonuclease for partial digestion of the DNA.
- Complete digestion is avoided since it generates fragments that are too heterogeneous in size.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT


COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III

BATCH-2017-2021

- For partial digestion, restriction enzymes having 4-base (tetrameric) recognition sequences are employed in preference to those having 6-base (hexameric) target sites. This is because a given 4-base recognition site is expected to occur every 44 (= 256) base pairs in a DNA molecule, while a 6 base target site would occur only after every 46 (=



4096) base pairs. (It is assumed here that the arrangement of the 4 bases in DNA molecules is random).

- Therefore, the fragments produced in partial digests with enzymes having 4 base recognition sites are more likely to be of appropriate size for cloning than those generated by enzymes having 6 base recognition sites.
- Single or mixed digestions with the enzymes AluI, HaeIII or Sau3A have been used for constructing genomic libraries.
- The use of restriction enzymes has the advantage that the same set of fragments are obtained from a DNA each time a specific enzyme is used, and many of the enzymes; produce cohesive ends.
- The partial digests of genomic DNA are subjected to agarose gel electrophoresis or sucrose gradient centrifugation for separation from the mixture of fragment of appropriate size.
- These fragments are then inserted into a suitable vector for cloning.
- This constitutes the shotgun approach to gene cloning. In principle, any vector can be used, but  $\lambda$  vectors and cosmids have been the most commonly used since DNA inserts of upto 23-25 kb (kilobase pairs) can be cloned in these vectors. The vectors containing the inserts are cloned in a suitable bacterial host.

### **Genomic DNA libraries *Producing representative genomic libraries in $\lambda$ cloning vectors***

let us suppose that to clone a single-copy gene from the human genome.

#### **How might this be achieved?**

- simply digest total human DNA with a restriction endonuclease, such as *EcoRI*, insert the fragments into a suitable phage- $\lambda$  vector and then attempt to isolate the desired clone.

#### **How many recombinants have to screen in order to isolate the right one?**

- Assuming *EcoRI* gives, on average, fragments of about 4 kb,
- and given that the size of the human haploid genome is  $2.8 \times 10^6$  kb, we can see that over  $7 \times 10^5$  independent recombinants must be prepared and screened in order to have a reasonable chance of including the desired sequence.

**There are two problems with the above approach.**

- First, the gene may be cut internally one or more times by *EcoRI* so that it is not obtained as a single fragment.
- These problems can be overcome by cloning *random* DNA fragments of a large size (for  $\lambda$  replacement vectors, approximately 20 kb).
- Since the DNA is randomly fragmented, there will be no systematic exclusion of any sequence.
- Furthermore, clones will overlap one another, allowing the sequence of very large genes to be assembled and giving an opportunity to 'walk' from one clone to an adjacent one.
- Because of the larger size of each cloned DNA fragment, fewer clones are required for a complete or nearly complete library.

**How many clones are required?**

Let  $n$  be the size of the genome relative to a single cloned fragment. Thus, for the human genome ( $2.8 \times 10^6$  kb) and an average cloned fragment size of 20 kb,  $n = 1.4 \times 10^5$ .

The number of independent recombinants required in the library must be greater than  $n$ , because sampling variation will lead to the inclusion of some sequences several times and the exclusion of other sequences in a library of just  $n$  recombinants.

Clarke and Carbon (1976) have derived a formula that relates the probability ( $P$ ) of including any DNA sequence in a random library of  $N$  independent recombinants:

$$N = \frac{\ln(1 - P)}{\ln\left(1 - \frac{1}{n}\right)}$$

Therefore, to achieve a 95% probability ( $P = 0.95$ ) of including any particular sequence in a random human genomic DNA library of 20 kb fragment size:

$$N = \frac{\ln(1 - 0.95)}{\ln\left(1 - \frac{1}{1.4 \times 10^5}\right)} = 4.2 \times 10^5$$

Notice that a considerably higher number of recombinants is required to achieve a 99% probability, for here  $N = 6.5 \times 10^5$ .

### **How can appropriately sized random fragments be produced?**

Various methods are available.

- Random breakage by mechanical shearing is appropriate because the average fragment size can be controlled, but insertion of the resulting fragments into vectors requires additional steps.
- The more commonly used procedure involves restriction endonucleases. In the strategy devised by Maniatis *et al.* (1978)
- the target DNA is digested with a mixture of *two* restriction enzymes.
- These enzymes have tetranucleotide recognition sites, which therefore occur frequently in the target DNA and in a complete double-digest would produce fragments averaging less than 1 kb.
- However, only a partial restriction digest is carried out, and therefore the majority of the fragments are large (in the range 10–30 kb).
- Given that the chances of cutting at each of the available restriction sites are more or less equivalent, such a reaction effectively produces a random set of overlapping fragments.
- These can be size-fractionated, e.g. by gel electrophoresis, so as to give a random population of fragments.

## **5. Cloning of cDNA**

### **cDNA Library**

- A cDNA library is a population of bacterial transformants or phage lysates in which each mRNA isolated from an organism or tissue is represented as its cDNA insertion in a plasmid or a phage vector.
- The frequency of a specific cDNA in such a library would ordinarily depend on the frequency of the concerned mRNA in the tissue/organism in question.

## 6.2. Preparation of cDNA

- cDNA is the copy or complementary DNA produced by using mRNA (usually) as a template. In fact, any RNA molecule can be used to produce cDNA.
- DNA copy of an RNA molecule is produced by the enzyme reverse transcriptase (RNA dependent DNA polymerase; discovered by **Temin and Baltimore** in 1970) generally obtained from avian myeloblastosis virus (AMV).
- This enzyme performs similar reactions as DNA polymerase, and has an absolute requirement for a primer with a free 3' -OH.
- When eukaryotic mRNA is used as a template, a poly T oligonucleotide (more specifically, oligodeoxynucleotide) is conveniently used as the primer since these mRNAs have a poly-A tail at their 3' ends.
- But special tricks are required to utilize primers for other RNAs, e.g., prokaryotic mRNA, rRNA, RNA virus genomes, etc.

For example, a poly A tail may be added to 3' end of the RNA to make it analogous to eukaryotic mRNA (oligo-T is now used as primer); this reaction is catalyzed by the enzyme poly A polymerase.

The appropriate oligonucleotide primer (oligo- T for eukaryotic mRNA) is annealed with the mRNA; this primer will base-pair to the 3'-end of mRNA.

Reverse transcriptase extends the 3'-end of the primer using mRNA molecule as a template. This produces a RNA.DNA hybrid molecule, the DNA strand being the cDNA.

The RNA strand is digested either by RNase H or alkaline hydrolysis; this frees the single-stranded cDNA.

- Curiously, the 3'-end of this cDNA serves as its own primer and provides the free 3'-OH required for the synthesis of its complementary strand; therefore, a primer is not required for this step.
- The complementary strand of cDNA single strand is synthesized by either the reverse transcriptase itself or by E. coli DNA polymerase; this generates a hairpin loop in the cDNA. The hairpin loop is cleaved by a single strand specific nuclease to yield a regular DNA duplex.

### **6.3. Problems in cDNA Preparation**

- Usually the double strand cDNA preparations are always a mixture of different kinds of molecules due to problems in copying of the RNA and also because even highly purified mRNAs are never absolutely pure.
- Physical and chemical methods are incapable of resolving these mixtures.
- Therefore, the cDNA mixture itself is used for cloning and the desired cDNA is identified and isolated in pure form from the appropriate bacterial clone.

### **6.4. Isolation of mRNA**

For isolation of mRNA, total RNA is first extracted from a suitable organism/tissue. The amount of desired mRNA in this sample is then increased by using one of the several procedures, some of which are listed below.

1. Chromatography on poly U sepharose or oligo T cellulose enriches the preparation with mRNAs of all kinds.
2. When the protein produced by a gene is known, it is purified and used to produce antibodies specific to it. These antibodies are used to precipitate the polysomes (mRNAs associated with ribosomes. and newly synthesized polypeptide chains), and the mRNA is isolated from them.
3. Some genes are expressed only in specific tissues, e.g., seed storage proteins in developing seeds, chicken ovalbumin gene in oviduct, etc. Therefore, mRNA preparations from such tissues are exceptionally rich in the concerned mRNA or may even contain only this mRNA. Use of cDNA is absolutely essential when the expression of an eukaryotic gene is required in a prokaryote, e.g., a bacterium. This is because eukaryotic genes have introns, which must be

removed from their transcripts to yield mature mRNA, and bacteria do not possess the enzymes necessary for removal of introns.

## **7. Transformation of rDNAs to host cell**

Transfer of rDNA into a suitable host cell is an important step in recombinant DNA technology looking for maintenance and expression of a foreign DNA in that cell.

The rDNA is safe in the cell and replicates independently of the chromosomal DNA of the cell. While doing so, the desired foreign gene present in the rDNA expresses its characters in the host cell. The host cell may be a bacterium or plant cell or animal cell. The cell containing an rDNA is known as transformed cell or transformant or recombinant.

Scientists have developed several methods to transfer genes into different types of host cells. The right method to be selected however depends on the type of gene cloning vector and nature of host cell, i.e. whether it is a bacterial cell or plant cell or animal cell.

The rDNAs can be introduced into host cells by *Transformation, Biolistics, Transduction, Electroporation, Transfection, Liposome fusion, Microinjection*.

### **1.1. Gene Transfer into Bacterial Cells**

The rDNA can be delivered into the bacterial cells by *Transformation, transduction and electroporation*. These methods are briefly described below.

#### **Transformation**

- Direct intake of DNA fragments in the medium by bacterial cells is called transformation. This method was first adopted to transfer rDNA into *E.coli* cells by Mandell and Higa in 1979.
- The ability of bacterial cells to intake DNAs from the medium is said to be competence.
- Competence can be increased by physical or chemical treatment.
- Competent bacterial cells can intake rDNAs with the size less than 15 kbp from the culture.
- To introduce rDNAs into *E.coli* cells, the rDNA is added to the bacterial culture and the culture is treated with 50mM calcium chloride ( $\text{CaCl}_2$ ) solution at room temperature.
- $\text{CaCl}_2$  adheres the rDNAs onto the surface of *E.coli* cells.

- It modifies the bacterial cell wall to intake rDNAs.
- The bacteria culture is then heated gently upto 42°C to induce *E.coli* cells to intake the rDNAs.
- *E.coli* cells intake the rDNA and become recombinants.
- Transformation method is also used to transfer rDNA into Cyanobacteria, microalgae, yeast and fungi.

### **Transduction**

- Introduction of a foreign DNA into a bacterial cell through a genetically modified virus or bacteriophage is called transduction.
- rDNAs of 38-50kbp size can be introduced into bacteria using this technique, but the bacteria should be the natural host of the virus.
- Phage derived vectors and cosmids have two cos-sites, one on either side of the DNA.
- These two cos-sites help the DNA to get packed inside the protein coat of lambda phage.
- The desired DNA is inserted into the viral vector DNA to construct an rDNA.
- The rDNA is packed in the protein coat of virus to get infective recombinant virus particles.
- When these virus particles are allowed to infect bacteria in a culture, the virus infects its rDNA into the bacterium.
- Thus, the bacterium receives foreign DNA in the rDNA from virus infection.

### **Electroporation**

- Electroporation is a process of changing the permeability of the cell membrane of cells to uptake macromolecules or organelles in the medium.
- It is done with an electrical instrument called electroporator.
- The electroporation unit consists of an electroporator creates electric current and sends it through the electrodes.
- It generates 50-240 volts through the electrodes. the aluminium electrodes conduct the electric pulse through the medium in the cuvette.
- The electrodes are fixed on the inner side of the cuvette.
- The cuvette is a small vessel of 10-50 ml capacity made of plastic or glass.

- The cell and DNA solution are mixed together and pipetted into the cuvette.
- The electroporator is setup to produce a current of 50-240 volts for 1-10 minutes.
- The electric current induces some permeability to the cell membrane and hence the cells uptakes the DNA from the medium.
- Electroporation is useful to transfer bacteria, fungi, plant cells and animal cells. The frequency of uptake can be increased by adding colcemid to the reaction mixture. To introduce rDNAs into E.coli cells an electric pulse of 2.5 kilo volt/cm is generated with the electrodes for 4.6 milliseconds. it is a rapid method to create more number of recombinants within a short time.

## 1.2. Gene transfer in to plant cell

A few important methods are available to delivered rDNA into plant cells and tissue.

1. Transfection,
2. electroporation,
3. ultrasonication,
4. partical bombardment,
5. agro infection and
6. liposome fusion.

These methods are discussed below:

### Transfection

- The direct intake of naked DNA from the culture medium by eukaryotic cells, is called **transfection** or **direct transformation** it is very similar to transformation in bacteria. Transfection takes place in plant cells and animal cell in vitro.
- Transformation in eukaryotic cells refers to differentiation of normal cells into cancer cells. Hence the word transfection is used to explain the same process in eukaryotes.
- young plants can uptake DNA fragments added to the neutrient solution by their route system. However, only a few cells get the DNA while most cells are been non-recombinants. Hence this method is not useful for gene manipulation.

- J.Paszkowski et al. (1984) made the isolated protoplast of tobacco to uptake rDNA having kanamycin resistance gene ,from the culture medium.They used PEG to induce competence in the protoplast.
- The protoplast took the rDNA from the medium and grew well in the medium containing kanamycin.
- The plants regenerate from the recombinants protoplast were resistant to kanamycin.this method provides the rDNA to all cells of the plant.
- How ever , large DNAs cannot be introduced into plant cells by this method.

### Electroporation

- Electroporation is a process of changing the permeability of cell membrane of cells to uptake macromolecules or organelles in medium.
- It is done with an **electrical instrument** called **electroporator**.
- The electroporation unit consist of an **electroporator**, **two aluminium electro** and a **cuvette**.
- The electroporator creates electric current and sends it through the electrodes.
- It generates 50 to 240 volts through the electrodes.
- The alluminium electrodes conduct the electric pulse through the medium into the cuvette.
- The electrodes are fixed on the inner side of the cuvette.
- The cuvette is a small vessel of 10 to 50 ml capacity made of plastic or glass .
- The cell and DNA solution are mixed together and pipette into the cuvette.
- The electroporator is set up to produce a current of 50 to 240 volts per 1 to 10 minutes.
- The electric current induces some permeability to the cell membrane and hence the cells uptake the DNA from the medium.
- Electroporation is useful to transform bacteria, fungi, plant cells and animal cells

The rDNA and protoplast suspension are mixed together and kept in the cuvette. Then a single electric pulse of about 4 to 8 kilo volts/cm is generated between two electrodes for 4 to 5 milli seconds. The electric pulse forms transient pores in the cell membrane and makes it permeable to the rDNAs.

The protoplast uptakes one or more copies of rDNA from the suspension by **pinocytosis**. The protoplast can uptake rDNA of 3 to 140kbp size.

The frequency of uptake can be further increased by adding colcemid to the reaction mixture.

### **Limitation**

1. Electroporation method cannot be adapted to introduce rDNAs into intact plants.
2. Animal cells are sensitive to electric treatment.
3. If the electric strength is too strong ,plant protoplast may loose their viability.

### **Ultrasonication**

- The method of subjecting plant cells to sound waves at the frequency of 20 to 35.1KHz is called as ultrasonication.
- These ultrasonic waves cannot be audible by human ears, but they do some changes on the surface of plant cells and cell membrane.
- Ultrasonic waves are generated by electronic devices called ultrasonicator.
- The ultrasonic method of gene delivery was discovered by wyber et al in 1999.
- Ultrasonication method is employed in the delivery of desired genes into seedlings, cotyledons, leaflets, immature embryos, callus and embryogenic cell suspensions.
- When a plant material is exposed to ultrasonic waves, the permeability of cell wall and plasma membrane is altered suitably as the cells uptake DNA or macromolecules.
- Plant cell transformation using ultrasonication is easy and simple to perform in ordinary laboratory. However, sonication could cause cell damage and rupture which are not likely during plant transformation.
- Hence, it is necessary to optimise the condition to facilitated uptake of DNA without damaging the cells.

### **Mechanism of DNA uptake**

- **Y.Leu et.al (2005)** explained the mechanism of DNA uptake by cells during sonication.
- When the cells are sonicated, the ultrasonic waves increases the temperature of the medium and produces acoustic cavitation of the cell surface.
- This cavitation increases the permeability of the cell wall and cell membrane.

- Meantime, the rate of mass transfer also increases in the medium.
- This increased mass transfer activity forces the DNA or macromolecule towards the cavitation so that the DNA can penetrate the cell membrane to reach the cell.
- Inside the cell, sonication is stopped the cell starts to regain its original structure by resynthesis of cell wall and membrane.

### **Ultrasonicators**

Many types of ultrasonicators are available for plant transformation.

They are of two types – **Bath sonicators and probe sonicators.**

#### **Bath sonicators**

- The bath sonicators are small automated cabinets equipped with ultra sound producing devices. They can produce ultra sounds with wavelength upto 55KHz.
- They are used to transform cotyledons, leaf tissues, shoot apex and meristem.

#### **Probe sonicators**

- The probe sonicator has one or two probes which produce ultrasonic waves.
- This type of wavelengths upto 20KHz.
- The probe sonicators are employed for the transformation of calli and cell suspensions.
- Probe of the sonicator is inserted in the tube containing plant material and the machine is switched on to produce ultrasound.
- It needs less time of sonication than bath sonicator.

### **Vessels for Sonication**

- Generally, borosilicate tubes (13x100mm) are preferred for the sonication of cotyledons and embryos in bath sonicators.
- Leaf tissues, meristem, apices and seedlings are kept in polypropylene tubes (50 ml) for sonication

### **Sonication**

- The plant material to be transformed is taken in a borosilicate tube containing liquid nutrient medium.

- Then, probe of the ultrasonicator is placed on the surface of the medium at a fixed position.
- This altered permeability makes the cells to uptake DNA from the medium.

#### Duration of Sonication

- Ultrasonication method has been used to deliver foreign genes into plants such as tobacco, wheat, sugar beet, maize, rice, etc.
- Monocot, dicot and gymnosperm plants can be transformed with this method.
- The duration of sonication, however, varies depending on the plant specimen to be transformed in the experiment-
  - Plant cells/tissues are sonicated for 2 seconds to 5 minutes.
  - Cotyledons are sonicated for 1-300seconds.
  - Leaf tissues are sonicated for 10 to 60 seconds.
  - Shoot tips, apices and meristem are sonicated for 5 to 300 seconds.
  - Embryonic suspensions are sonicated for 5 to 120 seconds.
  - Seedlings are sonicated for 10 to 100 seconds.

After this prescribed time point, the machine is switched off to stop sonication,

#### Transformation:

- The foreign gene is cloned in plasmid carrying genetic markers such as **cat**, **nptII** and **Gus** by using suitable restriction and DNA ligase.
- The rDNA may be added to the plant material in the suspension **before sonication or during sonication or after sonication**.
- The sonicated plant material intake the DNA from the solution to become **transformed** material in 1 hour.

#### Particle bombardment gun method:

- Shooting the plant or animal cells by DNA coated tungsten particle for getting DNAs into the cells, is called **particle bombardment** or **biolistics**.

- This method is also called microprojectile bombardment. It is a suitable method for introducing rDNA into plant cells, fungal cells, animal cells and cell organelles such as chloroplast and mitochondria.
- The rDNA is mixed with **tungsten or gold particles** of 1 to 4  $\mu\text{m}$  diameter and it is treated with  **$\text{CaCl}_2$  or spermidine or PEG**.
- **$\text{CaCl}_2$**  precipitates the rDNA on to the metal particles.
- The DNA coated tungsten or gold particles are said to be **microprojectiles**.
- The microprojectiles are fired into the plant cells or animal cells with the speed of 300 to 600 m/s(meter/sec) using an instrument called **microprojectile gun or particle gun or shot gun**.
- Gun powder, helium or electric power is used to provide propelling force to the gun to guide the microprojectile into the cells.
- The microprojectiles penetrate into the cell wall and cell membrane and deliver the rDNA into the cells.
- The rDNAs get integrated with the cell DNA and hence recombinant cells are formed.
- Biolistics can be adopted to introduce rDNA into intact plant cells having cell walls.
- It is a useful method to transfer novel genes into immature embryos, embryonic calli, root sections and epidermal cells of many monocot and dicot plants.
- Biolistics is an effective method to transfer rDNA into mammalian cells and organs.
- The skin and ears of live forms are shot with microprojectiles coated with rDNAs for getting the DNA into the cells.

**Pollen transformation through particle bombardment:**

- Delivery of rDNA into pollen grains using a particle bombardment gun is called **pollen transformation**.
- Pollen grains do not intake naked DNAs from the culture by means of transfection because of the presence of thick pollen wall.
- Again, gene transfer through callus transformation is time consuming and requires special tissue culture techniques and skilled persons.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III

BATCH-2017-2021

- Therefore, scientists have developed pollen transformation through particle bombardment gun to transfer desired genes to plants.

In 1997, **Ramaiah** and **Skinner** developed transgenic alfalfa plant by pollen transformation through particle bombardment. This method involves the following steps:

1. Mature pollen grains are collected from the flowers of alfalfa plant (*Medicago sativa*) by teasing the stamens with a needle.
2. The pollen grains are kept in a test tube and the tube is kept on ice for 1 hour to reduce the metabolic activities of the pollens.
3. Then the pollen grains are kept suspended in 13% sucrose solution for 10 minutes. The sucrose solution activates the pollen grains to start germination and it makes the pollen wall permeable to microprojectiles during bombardment.
4. The desired DNA is cloned in **pBI 121 plasmid** using proper restriction enzyme and DNA ligase. The pBI121 has GUS reporter gene as a genetic marker useful for screening the recombinants.
5. The gene cloned PBI 121 is mixed with tungsten or gold particles 1-4µm diameter and CaCl<sub>2</sub> precipitates the rDNA onto the metal particles to form DNA coated particles called **microprojectiles**.
6. The microprojectiles are loaded in a particle bombardment gun.
7. The pollen grains (in step 3) are place in a petridish and the microprojectiles are shoot into pollen by pressing the firing pin of the particle bombardment gun.
8. The suspension containing the pollen grains are taken in a test tube is kept on ice for 1 hr.
9. Then the suspension is centrifuged to remove the sucrose solution. The pellet of pollen thus obtained is again suspended in 10% sucrose solution for 30 minutes.
10. After removing the sucrose solution, the pollen grains are taken on a fine brush and dusted on the tip of pistil of emasculated flower. The transformed pollens fertilize the eggs to form transformed seeds.
11. Seeds are obtained from the artificially pollinated flowers and sown in separate rows to grow into seedlings in a green house.

12. After 30 days, the seedlings are tested for GUS activity to indicate transformants. To test GUS activity, protoplast isolated from each and every seedling and treated with 5-bromo 4-chloro-3-indolyl- $\beta$ -D-glucuronic acid. Protoplasts of transformants become blue coloured during incubation. This can be observed by viewing the protoplasts under a microscope.

13. The GUS positive seedlings are the genetically manipulated plants which give transgenic product.

Nearly, 30% of seedling raised from the seeds give rise to transgenic plants while the others remain untransformed ones. The transformed plants retain GUS activity for ten generations and then it may decrease slowly. The reason for low GUS activity after ten generations is that the plasmid may be lost spontaneously during repeated cell divisions. So transgenic plants raised through pollen transformation can only be maintained for about 10 generations.

### Agroinfection

- Transfer of foreign DNA to plant cells by *Agrobacterium* is called **Agroinfection**.
- *Agrobacterium* cannot pick up recombinant Ti plasmid from the nutrient medium.
- Therefore, the rDNA is introduced in to *E.coli*, it is transferred to *Agrobacterium* by three parental mating.
- The resulting recombinant *Agrobacteria* are co-cultured with plant cells or tissues for two days.
- During this time, the *agrobacteria* infects the plant cells and deliver their recombinant Ti plasmid in to the cells.
- The T-DNA integrated with the cell DNA and hence recombinant plant cells are formed.
- After proper selection and screening , plants are regenerated from the recombinant cells.

### Limitation

Argoinfection takes place only in dicot plants.

### Liposome fusion

- Liposomes are small spherical vesicles made of phospholipids.

- The rDNA to be delivered into plant cell is entrapped in liposomes which are then allowed to fuse with protoplast of that plant cell using PEG.
- When the liposome fuses with the cell membrane of the plant protoplast, the rDNA contained in it is delivered into the protoplasm of the cell.
- Several genes cloned in plasmid vectors have been transferred to protoplast of carrot, tobacco, petunia, tomato, etc. by using liposomes.

Gene transferred through liposome fusion **has following advantages**

- Liposomes do not show any toxicity to protoplast.
- They protect the rDNA from the cellular nuclease enzyme.
- The rDNA can be stored in liposome for a long time.
- Liposomes can be used for different types of cells.
- Preparation of liposomes with rDNAs is easy.
- Risk potential is very low.

### **1.3. Gene transfer into animals**

- The rDNA can be delivered into animal cells by  
transduction,  
electroporation,  
particle bombardment,  
retroviral method  
microinjection.

These methods are discussed below :

#### **Transduction**

- The direct intake of naked DNA from the culture medium by eukaryotic cells, is called **transfection**.
- It is very similar to transformation in bacteria.
- Transfection takes place in plant cells and animal cells invitro.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III

BATCH-2017-2021

- Foreign DNA can be introduced directly into animal cells and organs by transfection. Example – *He La cells, leukaemia L1210 cells, liver cells, lymphocytes, oocytes of mammals and vertebrates.*
- The DNA is dissolved in **phosphate buffer**. Then **calcium chloride** solution is added.
- This leads to the formation of **calcium phosphate**.
- it combines with DNA to form **calcium phosphate DNA precipitate**.
- This precipitate is added to the cells. The precipitate particles adhere with the cell surface.
- The cell engulfs the particles with (DNA) by phagocytosis.
- The DNA, entering the cell, is integrated with the cells genome.

### Liposome mediated gene transfer

- **The liposome** is a small spherical vesicle made of phosphor lipids. It is used to transfer genes to animal cells.
- It is formed when a lipid is agitated with water. It contains many concentric layers of phosphor lipids.
- In the liposome. Polar heads face outward and the non polar tail face to the centre. If there is more than one lipid layer, the adjacent layers are separated by water layer.
- The size of the liposomes varies from 25nm to a few microns in diameter.
- Ribosomes fuses with cell membrane and discharges its contents into the cell. Hence it is used as a gene transfer system.
- The rDNA, water and phosphatidyl choline are mixed together in a test tube and the tube is shaken well.
- During this process, the lipid bilayers develop around the rDNA present in water and form a liposome. In the liposome in the tube are added to a culture of animal cells.
- Liposomes fuse with cell membrane and discharge their contents into the cells.
- The rDNA present in the contents integrates with the cell DNA.
- However, the DNA may be destroyed in the cytoplasm before reaching the nucleus.
- The safe delivery of rDNA into the nucleus, the liposome is impregnated with some nuclear proteins and then another lipid layer is developed around it.

- The outer lipid layer fuses with the cell membrane and the rest is released into the cell.
- As the inner lipid layer has nucleoprotein, cellular enzyme do not attach it .
- The inner lipid layer fuses with nuclear membrane and discharges its contents into the nucleus.
- The frequency of liposome fusion can be increased by adding addition of polyethylene glycol(PEG) .
- Liposomes are also used to deliver certain drugs or rDNAs or enzymes in to cells of particular organs.

#### Advantages of liposome

- Any large DNA can be delivered into the cells using liposomes.
- Liposomes never disrupts the integrity of cells.
- They protect foreign DNAs from cellular nuclease enzymes.
- Liposomes with desired properties can be made by impregnating them with specific proteins. Liposomes never interfere with the human system.

#### Particle bombardment (Biolistics)

- *Shooting the plant or animal cells by DNA coated gold or tungsten particles for introducing DNA into the cells is called **particle bombardment or biolistics**.*
- This method is also called as *microprojectile bombardment*.
- By this method rDNA can be introduced in to plant cells, fungal cells, animal cells and cell organelles such as chloroplast and mitochondria.
- In this method ,no vector is introduced in to the rDNA in to the cells. The rDNA is literally shot in to cells.
- The instrument used to shoot the DNA in to cell are called **gene gun or microprojectile gun**
- The gene gun is doing the work of vector.

The gene gun consists of following components

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III

BATCH-2017-2021

- Gas acceleration tube
  - Firing pin
  - A blank charger
  - Macro projectile
  - Micro projectile
  - Stopping screen
  - Target cell
- All these components are enclosed in a vacuum chamber.
  - The rDNA is mixed with tungsten or gold particle of microscopic size and it is treated with  $\text{CaCl}_2$  or spermidine or PEG.
  - $\text{CaCl}_2$  precipitate the rDNA on to the metal particles.
  - The rDNA coated tungsten or gold particle said to be microprojectile.
  - The microprojectiles are positioned on a holder about 5 cm above the cells, within the chamber of the gun.
  - The chamber is sealed and helium gas is pumped into a small compartment situated above the gold particles.
  - Once a predetermined pressure is reached, the gas is released and the gold particles are shot into the cells.
  - The gold particle acts as **bullets**, penetrating the cells wall and delivering the rDNA into the cells, interior.
  - The microprojectiles are fired in to the plant cells or animal cells with the speed of 300-600 m/s using the gene gun.
  - **Gun powder, pressurized helium gas or electric power** is used to provide propelling force for the gun to drive the microprojectiles into the cells.
  - The rDNA gets intergrated with the cell DNA and hence recombinant cells are formed.
  - Biolistics can be adopted to introduce rDNA into intact plant cell having cell walls.
  - It is a useful method to transfer novel genes into immature embryos, embryonic calli, root section and epidermal cells of many monocot and dicot plants.
  - Biolistics is an effective method to transfer rDNAs into mammalian cells and organs.

- The skin and ears of live mouse are shot with microprojectiles coated with rDNAs for getting the DNAs into the cells.

### **Virus vector method**

- *Introduction of foreign DNA into eggs with the help of a retrovirus vector is retroviral infection.*
- Small foreign genes, less than 8 kbp, can be transferred to receipt cells through retrovirus vector. Murine leukaemia virus(MuLV) is the most common retrovirus used to introduce genes into animal cells.
- The rDNA is constructed and introduced into a mouse cell line by using calcium phosphate mediated transfection.
- The cell line is then infected with a mutant MuLV which has no signal for invivo packaging.
- RNA of the helper virus produces viral capsid.
- The rDNA produces recombinant viral RNA. The latter gets package in the viral capsid to form infective recombinant MuLV particles
- Eggs of mouse are collected and fertilized invitro to produce zygotes. On reaching 8-celled stage ,the mouse embryos are infected with the recombinant MuLV. Inside the infected cell the recombinant RNA is reverse transcribed into double stranded DNA. The DNA integrates with the cell DNA and hence recombinant cells are formed. thus , the virus delivers the rDNA into the animal cells.
- The mouse embryos are then transferred to a medium containing neomycin . The embryos that have recombinant cells alone survive in the medium and the others die.

### **Non viral method or microinjection**

- Microinjection refers to injection of DNAs or cell organelles directly into cell using a injection needle.
- By this method DNAs, RNAs, proteins or cell organelles into animal cells, eggs, zygotes and plant protoplast.
- A Stereoscopic dissecting microscope, Micropipette, Injection is needed.

- A Fertilized egg is transferred on to microscopic slide under the microscope. The cell is held in position using a sucking pipette. One end of sucking pipette is positioned on the surface of cell and a gentle suction pressure is applied on its other end.
- The rDNA is sucked into the glass injection needle. It is gently inserted into the zygote by viewing through the microscope.
- The rDNA is delivered into the zygote and the needle is drawn back carefully.
- The rDNA gets integrated onto the needle of the zygote.
- The microinjected embryos are cultured *in-vitro* upto the morula or blastocyst stage and then implanted into a female to produce transgenic organism.
- The surviving embryos are transferred into the uterus of a surrogate mother. These embryos develop into *transgenic mice*.

**Limitations:**

1. rDNAs can be microinjected only into a few cells at a time.
2. Microinjection needs technical skill experienced workers.

It needs a micromanipulator.

**Screening and expression of rDNA**

**1. Identification of the Desired Clone**

Identification of the bacterial colony containing the desired DNA fragments from among those making up the library.

**1.1. Screening strategies**

The identification of a specific clone from a DNA library can be carried out by

- exploiting the sequence of the clone
- the structure/function of its expressed product.

**Exploiting the sequence of the clone**

- applies to any type of library, genomic or cDNA, and
- can involve either nucleic acid hybridization or the PCR.

**1.2. Sequence-dependent screening**

methods are

1. Screening by hybridization
2. Chromosome walking
3. Screening by PCR

### 1.2.1. Screening by hybridization


Nucleic acid hybridization is the most commonly used method of library screening because

- it is rapid,
- it can be applied to very large numbers of clones and, in the case of cDNA libraries,
- can be used to identify clones that are not full-length (and therefore cannot be expressed).

**Grunstein and Hogness** (1975) developed a screening procedure

- to detect DNA sequences in transformed colonies by hybridization *in situ* with radioactive RNA probes.
  - Their procedure can rapidly determine which colony among thousands contains the target sequence.
1. The colonies to be screened are first replica-plated on to a nitrocellulose filter disc that has been placed on the surface of an agar plate prior to inoculation (Fig).
  2. A reference set of these colonies on the master plate is retained.
  3. The filter bearing the colonies is removed and treated with alkali so that the bacterial colonies are lysed and the DNA they contain is denatured.
  4. The filter is then treated with proteinase K to remove protein and leave denatured DNA bound to the nitrocellulose, for which it has a high affinity, in the form of a 'DNA print' of the colonies.
  5. The DNA is fixed firmly by baking the filter at 80°C.
  6. The defining, labelled RNA is hybridized to this DNA and the result of this hybridization is monitored by autoradiography.
  7. A colony whose DNA print gives a positive autoradiographic result can then be picked from the reference plate.

### Grunstein–Hogness method for detection of recombinant clones by colony hybridization




**Benton and Davis** (1977) devised a method called *plaque lift*, in which

1. the nitrocellulose filter is applied to the upper surface of agar plates, making direct contact between plaques and filter.
2. The plaques contain phage particles, as well as a considerable amount of unpackaged recombinant DNA.
3. Both phage and unpackaged DNA bind to the filter and can be denatured, fixed and hybridized.

#### Advantages

- several identical DNA prints can easily be made from a single-phage plate:
- this allows the screening to be performed in duplicate,
- increased reliability,
- allows a single set of recombinants to be screened with two or more probes.

The Benton and Davis (1977) procedure is the most widely applied method of library screening, successfully applied in thousands of laboratories to the isolation of recombinant phage by nucleic acid hybridization.



In  
place

of RNA probes, DNA or synthetic oligonucleotide probes can be used.

A number of alternative labelling methods are also available that avoid the use of radioactivity.

These methods involve the incorporation of chemical labels into the probe, such as digoxigenin or biotin, which can be detected with a specific antibody or the ligand streptavidin, respectively.

## **Hybridization probes & Labelling**

### **DNA and RNA probes**

- In molecular biology, a **hybridization probe** is a fragment of DNA or RNA of variable length (usually 100-1000 bases long), which is used in DNA or RNA samples to detect the presence of nucleotide sequences (the DNA target) that are complementary to the sequence in the probe.
- The probe thereby hybridizes to single-stranded nucleic acid (DNA or RNA) whose base sequence allows probe-target base pairing due to complementarity between the probe and target.
- The labeled probe is first denatured (by heating or under alkaline conditions such as exposure to sodium hydroxide) into single stranded DNA (ssDNA) and then hybridized to the target ssDNA (Southern blotting) or RNA (northern blotting) immobilized on a membrane or in situ.
- To detect hybridization of the probe to its target sequence, the probe is tagged (or labelled) with a molecular marker of either radioactive or (more recently) fluorescent molecules; commonly used markers are  $^{32}\text{P}$  (a radioactive isotope of phosphorus incorporated into the phosphodiester bond in the probe DNA) or Digoxigenin, which is non-radioactive antibody-based marker.
- DNA sequences or RNA transcripts that have moderate to high sequence similarity to the probe are then detected by visualizing the hybridized probe via autoradiography or other imaging techniques. Normally, either X-ray pictures are taken of the filter, or the filter is placed under UV light.
- Detection of sequences with moderate or high similarity depends on how stringent the hybridization conditions were applied — high stringency, such as high hybridization temperature and low salt in hybridization buffers, permits only hybridization between nucleic acid sequences that are highly similar, whereas low stringency, such as lower temperature and high salt, allows hybridization when the sequences are less similar.

Hybridization probes used in DNA microarrays refer to DNA covalently attached to an inert surface, such as coated glass slides or gene chips, and to which a mobile cDNA target is hybridized.

Depending on the method the probe may be synthesized using phosphoramidite method or generated and labeled by PCR amplification or cloning (older methods).

In order to increase the in vivo stability of the probe RNA is not used, instead RNA analogues may be used, in particular morpholino. Molecular DNA- or RNA-based probes are now routinely used in screening gene libraries, detecting nucleotide sequences with blotting methods, and in other gene technologies like microarrays.

### ***Probe design***

A great advantage of hybridization for library screening is that it is extremely versatile.

In hybridization is **very stringent conditions** can be used

- so that only sequences identical to the probe are identified.
- This is necessary, for example, to identify genomic clones corresponding to a specific cDNA or to identify overlapping clones in a chromosome walk.

Alternatively, **less stringent conditions** can be used

- to identify both identical and related sequences.
- This is appropriate where a probe from one species is being used to isolate a homologous clone from another species.

Probes corresponding to a conserved functional domain of a gene may also cross-hybridize with several different clones in the same species at lower stringency, and this can be used to identify members of a gene family.

Hybridization thus has the potential to isolate any sequence from any library *if a probe is available*.

If a suitable DNA or RNA probe cannot be obtained from an existing cloned DNA,

- an alternative strategy is to make an oligonucleotide probe by chemical synthesis.
- This requires some knowledge of the amino acid sequence of the protein encoded by the target clone.

### 1.2.2. Chromosome walking

- The principle of chromosome walking is that overlapping clones will hybridize to each other, allowing them to be assembled into a contiguous sequence.
- This can be used to isolate genes whose function is unknown but whose genetic location is known, a technique known as *positional cloning*.

To begin a chromosome walk, it is necessary to have in hand a genomic clone that is known to lie very close to the suspected location of the target gene.

For example - In humans,

- this could be a restriction fragment length polymorphism that has been genetically mapped to the same region.
- This clone is then used to screen a genomic library by hybridization, which should reveal any overlapping clones.
- These overlapping clones are then isolated, labeled and used in a second round of screening to identify further overlapping clones,
- and the process is repeated to build up a contiguous map.

### 1.2.3. Screening by PCR

The PCR is widely used to isolate specific DNA sequences from uncloned genomic DNA or cDNA, but it also a useful technique for library screening.

As a screening method,

- PCR has the same versatility as hybridization, and the same limitations.
- It is possible to identify any clone by PCR but only if there is sufficient information about its sequence to make suitable primers.

To isolate a specific clone, PCR is carried out with gene-specific primers that flank a unique sequence in the target.

- Instead of plating the library out on agar, as would be necessary for screening by hybridization, pools of clones are maintained in multiwell plates.
- Each well is screened by PCR and positive wells are identified.

- The clones in each positive well are then diluted into a series in a secondary set of plates and
- screened again.
- The process is repeated until wells carrying homogeneous clones corresponding to the gene of interest have been identified.

There are also several applications where the use of *degenerate primers* is favourable.

- A degenerate primer is a mixture of primers, all of similar sequence but with variations at one or more positions.
- This is analogous to the use of degenerate oligonucleotides as hybridization probes, and the primers are synthesized in the same way.
- A common circumstance requiring the use of degenerate primers is when the primer sequences have to be deduced from amino acid sequences (Lee *et al.* 1988).
- Degenerate primers may also be employed to search for novel members of a known family of genes or to search for homologous genes between species.

## 2. Exploitation of structural or functional property of the clone

### Screening expression libraries - (expression cloning)

If a DNA library is established using expression vectors, each individual clone can be expressed to yield a polypeptide.

Expression libraries are useful because they allow a range of alternative techniques to be employed, each of which **exploits some structural or functional property of the gene product**.

This can be important in cases where

- the DNA sequence of the target clone is completely unknown and
- there is no strategy available to design a suitable probe or set of primers.

Methods of screening are –

1. *Immunological screening*
2. *South-western and north-western screening*
3. *Screening with alternative ligands*

## 2.1. Immunological screening

Immunological screening involves the use of antibodies that specifically recognize antigenic determinants on the polypeptide synthesized by a target clone.

- This is one of the most versatile expression cloning strategies, because it can be applied to any protein for which an antibody is available.
- The molecular target for recognition is generally an *epitope*, a short sequence of amino acids that folds into a particular three-dimensional conformation on the surface of the protein.
- Epitopes can fold independently of the rest of the protein and therefore often form even when the polypeptide chain is incomplete or when expressed as a fusion with another protein.
- Importantly, many epitopes can form under denaturing conditions, when the overall conformation of the protein is abnormal.

The first immunological screening techniques were developed in the late 1970s, when expression libraries were generally constructed using plasmid vectors.

- The method of **Broome and Gilbert (1978)** was widely used at the time.

This method exploited the fact that antibodies adsorb very strongly to certain types of plastic, such as polyvinyl, and that IgG antibodies can be readily labelled with  $^{125}\text{I}$  by iodination *in vitro*.

1. As usual, transformed cells were plated out on Petri dishes and allowed to form colonies.
2. In order to release the antigen from positive clones, the colonies were lysed, e.g. using chloroform vapour or by spraying with an aerosol of virulent phage (a replica plate is required because this procedure kills the bacteria).
3. A sheet of polyvinyl that had been coated with the appropriate antibody was then applied to the surface of the plate, allowing antigen–antibody complexes to form.
4. The sheet was then removed and exposed to  $^{125}\text{I}$ -labelled IgG specific to a *different* determinant on the surface of the antigen (i.e. a determinant not involved in the initial binding of the antigen to the antibody-coated sheet (Fig. 6.1 4)).
5. The sheet was then washed and exposed to X-ray film.

6. The clones identified by this procedure could then be isolated from the replica plate.
7. 'sandwich' technique is applicable only where two antibodies recognizing different determinants of the same protein are available.
8. However, if the protein is expressed as a fusion, antibodies that bind to each component of the fusion can be used, efficiently selecting for recombinant molecules.
  - it is much more convenient to use bacteriophage- $\lambda$  insertion vectors, because these have a higher capacity and the efficiency of *in vitro* packaging allows large numbers of recombinants to be prepared and screened.
  - Immunological screening with phage- $\lambda$  cDNA libraries was introduced by **Young and Davies** (1983) using the expression vector  $\lambda$ gt11, which generates fusion proteins with  $\beta$ -galactosidase under the control of the *lac* promoter.

**In the original technique,**

screening was carried out using colonies of induced lysogenic bacteria, which required the production of replica plates, as above.

A simplification of the method is possible by directly screening plaques of recombinant phage.

**In this procedure (Fig),**


1. the library is plated out at moderately high density (up to  $5 \times 10^4$  plaques/ cm<sup>2</sup> plate), with *E. coli* strain Y1090 as the host.
2. This *E. coli* strain overproduces the *lac* repressor and ensures that no expression of cloned sequences (which may be deleterious to the host) takes place until the inducer isopropyl- $\beta$ -D-thiogalactoside (IPTG) is presented to the infected cells.
3. Y1090 is also deficient in the *lon* protease, hence increasing the stability of recombinant fusion proteins.
4. Fusion proteins expressed in plaques are absorbed on to a nitrocellulose membrane overlay, and this membrane is processed for antibody screening.
5. When a positive signal is identified on the membrane, the positive plaque can be picked from
6. the original agar plate (a replica is not necessary) and the recombinant phage can be isolated.

The original detection method using **iodinated antibodies** has been replaced by more convenient methods using non-isotopic labels, which are also more sensitive and have a lower background of nonspecific signal.

- Generally, these involve the use of unlabelled primary antibodies directed against the polypeptide of interest, which are in turn recognized by secondary antibodies carrying an enzymatic label.
- As well as eliminating the need for isotopes, such methods also incorporate an amplification step, since two or more secondary antibodies bind to the primary antibody.

Typically, the secondary antibody recognizes the species-specific constant region of the primary antibody and is conjugated to either horseradish peroxidase or alkaline phosphatase, each of which can in turn be detected using a simple colorimetric assay carried out directly on the nitrocellulose filter.

- Polyclonal antibodies, which recognize many different epitopes, provide a very sensitive probe for immunological screening, although they may also cross-react to proteins in the expression host.



- Monoclonal antibodies and cloned antibody fragments can also be used, although the sensitivity of such reagents is reduced because only a single epitope is recognized.

## 2.2. South-western and north-western screening

A closely related approach to immunochemical screening has been used for the screening and isolation of clones expressing sequence specific DNA-binding proteins.

### Procedure

1. a plaque lift is carried out to transfer a print of the library on to nitrocellulose membranes.
2. the screening is carried out, without using an antibody, by incubating the membranes with a radiolabelled *double stranded* DNA oligonucleotide probe, containing the recognition sequence for the target DNA-binding protein.
3. This technique is called ***south-western screening***, because it combines the principles of Southern and western blots.
4. It has been particularly successful in the isolation of clones expressing cDNA sequences corresponding to certain mammalian transcription factors.

Limitation of this technique is that,

- since individual plaques contain only single cDNA clones, transcription factors that function only in the form of heterodimers or as part of a multimeric complex do not recognize the DNA probe and the corresponding cDNAs cannot be isolated.
- the procedure can also be successful only in cases where the transcription factor remains functional when expressed as a fusion polypeptide.

More recently, a similar technique has been used to isolate sequence specific RNA-binding proteins, in this case using a single-stranded RNA probe.

By analogy to the above, this is termed ***north-western screening*** and has been successful in a number of cases

## 2. 3. Screening with alternative ligands

- As well as DNA and RNA, a whole range of alternative 'ligands' can be used to identify polypeptides that specifically bind certain molecules (for example, as an alternative to south-western screening).
- Such techniques are not widely used because they generally have a low sensitivity and their success depends on the preservation of the appropriate interacting domain of the protein when exposed on the surface of a nitrocellulose filter.
- the yeast two-hybrid system and its derivatives provide versatile assay formats for many specific types of protein–protein interaction, with the advantage that such interactions are

tested in living cells, so the proteins involved are more likely to retain their functional interacting domains.

### **3. Basic techniques in screening recombinat DNA**

#### **3.1. Agarose gel electrophoresis**

- The progress of the first experiments on cutting and joining of DNA molecules was monitored by velocity sedimentation in sucrose gradients. However, this has been entirely replaced by gel electrophoresis.
- Gel electrophoresis is not only used as an analytical method, it is routinely used preparatively for the purification of specific DNA fragments.
- The gel is composed of polyacrylamide or agarose.
  - Agarose is convenient for separating DNA fragments ranging in size from a few hundred base pairs to about 20 kb.
  - Polyacrylamide is preferred for smaller DNA fragments.
- The migration of the DNA molecules through the pores of the matrix must play an important role in molecular-weight separations since the electrophoretic mobility of DNA in free solution is independent of molecular weight.
- An agarose gel is a complex network of polymeric molecules whose average pore size depends on the buffer composition and the type and concentration of agarose used.
- DNA movement through the gel was originally thought to resemble the motion of a snake (reptation).
- DNA molecules display elastic behaviour by stretching in the direction of the applied field and then contracting into dense balls.
- The larger the pore size of the gel, the greater the ball of DNA which can pass through and hence the larger the molecules which can be separated.
- Once the globular volume of the DNA molecule exceeds the pore size, the DNA molecule can only pass through by reptation.
- This occurs with molecules about 20 kb in size and it is difficult to separate molecules larger than this without recourse to pulsed electrical fields.

- The bands of DNA in the gel are stained with the intercalating dye ethidium bromide and as little as 0.05 µg of DNA in one band can be detected as visible fluorescence when the gel is illuminated with ultraviolet light.
- In addition to resolving DNA fragments of different lengths, gel electrophoresis can be used to separate different molecular configurations of a DNA molecule.
- Gel electrophoresis can also be used for investigating protein–nucleic acid interactions in the so-called *gel retardation* or *band shift* assay.
- It is based on the observation that binding of a protein to DNA fragments usually leads to a reduction in electrophoretic mobility.
- The assay typically involves the addition of protein to linear double-stranded DNA fragments, separation of complex and naked DNA by gel electrophoresis and visualization.
- Marker DNA fragments of known sizes are run in a separate lane; this permits an accurate determination of the size of an unknown DNA molecule by interpolation. The gels are stained with the intercalating dye ethidium bromide which, gives visible fluorescence on illumination of the gel with UV light; as little as 0.05 µg of DNA in one band can be detected by using this dye.

### 3.2. Nucleic acid blotting

Nucleic acid labelling and hybridization on membranes have formed the basis for a range of experimental techniques central to recent advances in our understanding of the organization and expression of the genetic material.

These techniques may be applied in the isolation and quantification of specific nucleic acid sequences and in the study of their organization, intracellular localization, expression and regulation.

A variety of specific applications includes the diagnosis of infectious and inherited disease.

*Blotting* describes the immobilization of sample nucleic acids on to a solid support, generally nylon or nitrocellulose membranes.

The blotted nucleic acids are then used as ‘targets’ in subsequent hybridization experiments. The main blotting procedures are:

### 3.2.1. Southern Hybridization

The name of this technique is derived from the following:

- (1) the name of its inventor, E.M. Southern, and
  - (2) the DNA-DNA hybridization that forms its basis. It is also called Southern blotting since the procedure for transfer of DNA from the gel to the nitrocellulose filter resembles blotting.
1. This technique has since been extended to the analysis of RNA (northern blotting) and proteins (western blotting); these names are only jargon terms, i.e., reverse of Southern being northern and so on, and do not reflect any functional or historical significance.
  2. This approach is useful when few DNA fragments with considerable length differences are to be separated and studied. This approach also separates the closed circular (supercoiled), nicked (relaxed) and linear configurations of a single DNA molecule.
  3. In many situations, it is critical to detect and identify DNA fragments in a sample that are complementary to a given DNA sequence, e.g., to demonstrate the presence of the gene in question in transgenics, to detect and study RFLP (restriction fragment length polymorphism), etc.

In Southern hybridization, a sample of DNA containing fragments of different sizes is subjected to electrophoresis using either polyacrylamide or agarose gel.

The DNA sample may either be subjected to mechanical shearing or to restriction endonuclease digestion in order to generate the fragments.

**This is achieved by Southern hybridization in which the following steps are performed.**

1. The restriction fragments of DNA present in agarose gel (after electrophoresis) are denatured into single-stranded form by alkali treatment.
2. They are then transferred onto a nitrocellulose filter membrane; this is done by placing the gel on top of a buffer saturated filter paper, then laying the nitrocellulose filter membrane on the top of gel, and finally placing some dry filter papers on top of this membrane. The buffer moves, due to capillary action, from the bottom filter paper through the gel carrying with it the denatured DNA present in the gel; the DNA becomes trapped in the

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III

BATCH-2017-2021


nitrocellulose membrane as the buffer phases through it. This process is known as blotting and takes several hours to complete. The relative positions of the bands on the membrane remain the same as those in the gel and there is a minimal loss in their resolution (sharpness).

3. The nitrocellulose membrane is now removed from the blotting stack, and the DNA is permanently immobilized on the membrane by baking it at 80°C in vacuo.
  4. Single-stranded DNA has a high affinity for nitrocellulose filter membrane. (Note that RNA lacks this affinity). Therefore, the baked membrane is treated, with a solution containing 0.2% each of Ficoll (an artificial polymer of sucrose), polyvinylpyrrolidone and bovine serum albumin; this mixture is often supplemented with an irrelevant nucleic acid, e.g., tRNA (pretreatment). This treatment prevents nonspecific binding of the radioactive probe (to be used in the next step) probably by attaching macromolecules to all the free binding sites on the membrane. Often the above mixture is included in the hybridization reaction itself.
  5. The pretreated membrane is placed in a solution of radioactive, single-stranded DNA or an oligodeoxynucleotide (a DNA segment having few to several nucleotides) called probe. The name probe signifies the fact that this DNA molecule is used to detect and identify the DNA fragment in the gel/membrane that is complementary to the probe. The conditions during this step are chosen so that the probe hybridizes with the complementary DNA on the membrane to the greatest extent with a low nonspecific binding on the membrane; this step is known as hybridization reaction.
- Usually, the initial hybridization reaction is carried out under conditions of relatively low stringency of hybridization to permit a high rate of hybridization; this is followed by a series of post hybridization washes of increasing stringency, i.e., higher temperature or, more commonly, lower ionic strength, with a view to eliminate the pairing of radioactive probe to related sequences and to allow only perfectly complementary pairing.
6. After the hybridization reaction, the membrane is washed to remove the unbound probes.
  7. The membrane is now placed in close contact with an X-ray film and incubated for a desired period to allow images due to the radioactive probes to be formed on the film. The film is then developed to reveal distinct band(s) indicating positions in the gel of the

DNA fragments that are complementary to the radioactive probe used in the study.

It should be kept in mind that electrophoresis of sheared or restricted DNA produces a smear in which the fragments are distributed in a continuum according to their size, and there are no distinct bands. The distinct bands are produced by the hybridization reaction of the selected probe with one or few fragment sequences present in the gel.

The Southern blotting technique is extremely sensitive. It can be used to map the restriction sites around a single copy gene sequence in any genome (even of man). It is used for DNA fingerprinting, preparation of RFLP maps, detection and identification of the transferred genes in transgenic individuals, etc.



Recently some new membrane materials, e.g., nylon membranes, have been developed which have the following advantageous features:

- (1) They are physically robust in comparison to nitrocellulose filter membranes,
- (2) both DNA and RNA become cross-linked to them by a brief exposure to UV light, which
- (3) saves the time needed for baking in vacuo in the case of nitrocellulose membranes, and
- (4) the same membrane blot, e.g., a membrane onto which DNA/RNA has been transferred from a gel and cross-linked by UV exposure, can be used for search with more than one probe after removing the earlier probe by high temperature washing or some other denaturing procedure; in other words, the nylon membranes are reusable.

### **3.2.2. Northern Hybridization** - In this technique,

RNAs are separated by gel electrophoresis, the RNA bands are transferred onto a suitable membrane, e.g., diazobenzyloxymethyl (DBM) paper or nylon membranes, and immobilized; the bands are hybridized with radioactive single stranded DNA probes, and the bands showing hybridization are detected by autoradiography.

Clearly, northern blotting/hybridization is simply an extension of the Southern blotting technique. The two techniques, however, show the following differences:

- (1) in Southern hybridization, DNA is separated by gel electrophoresis, while in northern blotting RNAs are separated;
- (2) as a result, in Southern hybridization DNA has to be denatured before blotting, while this step is not needed in northern hybridization;
- (3) nitrocellulose membrane is generally not used for northern, while it is often used for Southern hybridization; and finally
- (4) hybridization with the probe produces DNA: DNA hybrid molecules in Southern hybridization but RNA : DNA molecules in northern hybridization.

Initially, specially prepared paper (diazobenzyloxymethyl, DBM, paper prepared by diazotization of aminobenzyloxymethyl paper) was used for northern blotting since RNA did not

bind to nitrocellulose membrane. RNA becomes covalently bound to DBM paper due to which these blot transfers are reusable.

DBM is also equally effective in binding to denatured DNA, and is more efficient than nitrocellulose in binding to small DNA fragments. Recently developed nylon membranes have superseded the use of DBM paper as they are robust, reusable and bind (by cross linking) to RNA on a brief exposure to UV light.

Northern hybridization is useful in the identification and separation of the RNA that is complementary to a specific DNA probe; this is a sensitive test for the detection of transcription of a DNA sequence that is used as probe.

### **3.2.3. Western Blotting** - In western blotting,

proteins are electrophoresed in polyacrylamide gel, transferred onto a nitrocellulose or nylon membrane (to which they bind strongly), and the protein bands are detected by their specific interaction with antibodies, lectins or some other compounds. The various steps of this technique are briefly described below.

1. Protein bands are separated by polyacrylamide gel electrophoresis.
2. The protein bands are transferred onto a nitrocellulose or nylon membrane; initially this was achieved by a capillary movement of buffer similar to Southern blotting (capillary blotting), but nowadays it is usually done by electrophoresis (electrophoretic blotting).

Electrophoresis has been applied for the blotting step in Southern and northern hybridizations as well; in such cases buffer of low ionic strength (to avoid overheating during electrophoresis) and nylon membranes (since nucleic acids bind to nitrocellulose membrane only under conditions of high ionic strength) are used. The electrophoretic blotting, both of proteins and nucleic acids is much faster and more efficient than capillary blotting.

3. The specific protein bands are identified in a variety of ways.

- (i) Antibodies are the most commonly used as probes for detecting specific antigens.
- (ii) Lectins are used as probes for the identification of glycoproteins.
  - o These probes may themselves be radioactive or a radioactive molecule may be tagged to them. Often the identification process is based on a 'sandwich' reaction.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: III

BATCH-2017-2021

- In such an approach, a species specific second antibody or protein A of *Staphylococcus aureus* (protein A binds to certain subclasses of IgG antibodies) or streptavidin (it binds to biotinylated antibodies) is used to bind to the antibodies bound to the protein bands.

These second molecules may be labelled with radioactive, enzyme or fluorescent tags; a single preparation of these labelled molecules can be employed as a general detector for various probes.

UNIT-IV

SYLLABUS

**UNIT-IV**

**Random and site-directed mutagenesis:** Primer extension and PCR based methods of site directed mutagenesis, Random mutagenesis, Gene shuffling, production of chimeric proteins, Protein engineering concepts and examples (any two).

**PCR (Polymerase Chain Reaction)**

- first proposed in the early 1970s by **Ghobind Khorana** and his colleagues
- did not seem practicable at a time when genes had not yet been sequenced, synthesis of oligonucleotide primers and thermostable DNA polymerases had not been described.

After 15 years, **Kary Mullis and coworkers** at Cetus Corporation,

- Performed in vitro amplification of single-copy mammalian genes using the Klenow fragment of *Escherichia coli* DNA polymerase I.
- The use of a thermostable polymerase from *Thermus aquaticus* greatly increased the efficiency of PCR and opened the door to automation of the method.

**By the end of the 1980s,**

- cloning was no longer the only way to isolate genes:
- DNA sequencing had been revolutionized
- and **PCR had become a fundamental step of genetic and molecular analyses.**

**4.1. Essential Components of Polymerase Chain Reactions**

PCRs contain **seven** essential components:

- Thermostable DNA polymerase
- Pair of synthetic oligonucleotides
- Deoxynucleoside triphosphates (dNTPs).
- Divalent cations
- Buffer
- Monovalent cations

## Template DNA

### 1. A thermostable DNA polymerase

to catalyze template-dependent synthesis of DNA.

A wide choice of enzymes is now available that vary in their fidelity, efficiency, and ability to synthesize large DNA products

For routine PCRs, *Taq* polymerase (0.5-2.5 units per standard 25-50  $\mu$ l reaction) remains the enzyme of choice.

The specific activity of most commercial preparations of *Taq* is  $\sim 80,000$  units/mg of protein.

Standard PCRs contain  $2 \times 10^{12}$  to  $10 \times 10^{12}$  molecules of enzyme.

efficiency of primer extension with *Taq* polymerase is generally  $\sim 0.7$ .

### 2. A pair of synthetic oligonucleotides

- to prime DNA synthesis.
- Careful design of primers is required to obtain the desired products in high yield, to suppress amplification of unwanted sequences, and to facilitate subsequent manipulation of the amplified product.
- Standard reactions contain non limiting amounts of primers, typically 0.1-0.5  $\mu$ M of each primer ( $6 \times 10^{12}$  to  $3 \times 10^{13}$  molecules).
- This quantity is enough for at least 30 cycles of amplification of a 1 -kb segment of DNA.
- Higher concentrations of primers favor mispriming, which may lead to nonspecific amplification.
- Oligonucleotide primers synthesized on an automated DNA synthesizer can generally be used in standard PCRs without further purification.

### 3. Deoxynucleoside triphosphates (dNTPs)

- Standard PCRs contain equimolar amounts of dATP, dTTP, dCTP, and dGTP.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: IV

BATCH-2017-2021

- Concentrations of 200-250  $\mu$ M of each dNTP are recommended for *Taq* polymerase in reactions containing 1.5 mM  $MgCl_2$ .
- In a 50- $\mu$ l reaction, these amounts should allow synthesis of ~6-6.5  $\mu$ g of DNA, which should be sufficient even for multiplex reactions in which eight or more primer pairs are used at the same time.
- High concentrations of dNTPs ( $>4$  mM) are inhibitory, perhaps because of sequestering of  $Mg^{2+}$ .
- However, a satisfactory amount of amplified product can be produced with dNTP concentrations as low as 20  $\mu$ M. 0.5- $\mu$ mol of an amplified fragment ~1 kb in length,

#### 4. **Divalent cations.**

- All thermostable DNA polymerases require free divalent cations – usually  $Mg^{2+}$  – for activity.
- Calcium ions are quite ineffective). Because dNTPs and oligonucleotides bind  $Mg^{2+}$ , the molar concentration of the cation must exceed the molar concentration of phosphate groups contributed by dNTPs plus primers.
- It is therefore impossible to recommend a concentration of  $Mg^{2+}$  that is optimal in all circumstances.
- Although a concentration of 1.5 mM  $Mg^{2+}$  is routinely used, increasing the concentration of  $Mg^{2+}$  to 4.5 mM or 6 mM has been reported to decrease nonspecific priming in some cases and to increase it in others.
- The optimal concentration of  $Mg^{2+}$  must therefore be determined empirically for each combination of primers and template.

#### 5. **Buffer to maintain pH**

- Tris-Cl, adjusted to a pH between 7.5 and 8.8 at room temperature, is included in standard PCRs at a concentration of 10 mM.
- Then incubated at 72°C (the temperature commonly used for the extension phase of PCR), the pH of the reaction mixture drops by more than a full unit, producing a buffer whose pH is ~7.2.

#### 6. **Monovalent cations**

- Standard PCR buffer contains 50 mM KCl and works well for amplification of segments of DNA >500 bp in length.
- Raising the KCl concentration to 70-100 mM often improves the yield of shorter DNA segments.

#### 7. Template DNA

- Template DNA containing target sequences can be added to PCR in single- or double-stranded form.
- Closed circular DNA templates are amplified slightly less efficiently than linear DNAs. Although the size of the template DNA is not critical, amplification of sequences embedded in high-molecular-weight DNA (> 10 kb) can be improved by digesting the template with a restriction enzyme that does not cleave within the target sequence.
- In the case of mammalian genomic DNA, up to 1.0 µg of DNA is utilized per reaction, an amount that contains ~3 x 10<sup>5</sup> copies of a single-copy autosomal gene.
- The typical amounts of yeast, bacterial, and plasmid DNAs used per reaction are 10 ng, 1 ng, and 1 pg, respectively.

#### 8. Thermostable DNA Polymerases

- Thermostable DNA polymerases are isolated from two classes of organisms:
- the thermophilic and hyperthermophilic eubacteria Archaeobacteria,
- *T. aquaticus*, an organism from the thermophilic Archaea family *Taq* (*T. aquaticus*) DNA polymerase, the first isolated and best understood of the thermostable DNA polymerases, remains the workhorse of PCR in most laboratories.

### 4.2. Programming Polymerase Chain Reactions

PCR is an iterative process, consisting of three elements:

- denaturation of the template by heat,
- annealing of the oligonucleotide primers to the single-stranded target sequence(s), and
- extension of the annealed primers by a thermostable DNA polymerase.

#### Denaturation

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: IV

BATCH-2017-2021

- Double-stranded DNA templates denature at a temperature that is determined in part by their G+C content.
- The higher the proportion of G+C, the higher the temperature required to separate the strands of template DNA.
- The longer the DNA molecules, the greater the time required at the chosen denaturation temperature to separate the two strands completely.
- If the temperature for denaturation is too low or if the time is too short, only AT-rich regions of the template DNA will be denatured.
- When the temperature is reduced later in the PCR cycle, the template DNA will reanneal into a fully native condition.
- In PCRs catalyzed by *Taq* DNA polymerase, denaturation is carried out at 94-95°C, which is the highest temperature that the enzyme can endure for 30 or more cycles without sustaining excessive damage.
- In the first cycle of PCR, denaturation is sometimes carried out for 5 minutes to increase the probability that long molecules of template DNA are fully denatured.
- We recommend denaturation for 45 seconds at 94-95°C for routine amplification of linear DNA templates whose contents of G+C is 55% or less.
- Higher temperatures may be required to denature template and/or target DNAs that are rich in G+C (>55%).

### ***Annealing of primers to template DNA***

- The temperature used for the annealing step ( $T$ ) is critical.
- If the annealing temperature is too high, the oligonucleotide primers anneal poorly, if at all, to the template and the yield of amplified DNA is very low.
- If the annealing temperature is too low, nonspecific annealing of primers may occur, resulting in the amplification of unwanted segments of DNA.
- Annealing is usually carried out 3-5°C lower than the calculated melting temperature at which the oligonucleotide primers dissociate from their templates.
- Many formulas exist to determine the theoretical melting temperature, but none of them are accurate for oligonucleotide primers of all lengths and sequences.

- It is best to optimize the annealing conditions by performing a series of trial PCRs at temperatures ranging from 20°C to 100°C below the lower of the melting temperatures calculated for the two oligonucleotide primers.
- Alternatively, the thermal cycler can be programmed to use progressively lower annealing temperatures in consecutive pairs or cycles ("touchdown" PCR).

#### **Extension of oligonucleotide primers**

- is carried out at or near the optimal temperature for DNA synthesis catalyzed by the thermostable polymerase, which in the case of *Taq* DNA polymerase is 72-78°C.
- In the first two cycles, extension from one primer proceeds beyond the sequence complementary to the binding site of the other primer.
- In the next cycle, the first molecules are produced whose length is equal to the segment of DNA delimited by the binding sites of the primers.
- From the third cycle onward, this segment of DNA is amplified geometrically, whereas longer amplification products accumulate arithmetically.
- The polymerization rate of *Taq* polymerase is - 2000 nucleotides/minute at the optimal temperature (72-78°C) and as a rule of thumb, extension is carried out for 1 minute for every 1000 bp of product.
- For the last cycle of PCR, many investigators use an extension time that is three times longer than in the previous cycles, to allow completion of all amplified products.

#### **Number of cycles**

- the number of cycles required for amplification depends on the number of copies of template DNA present at the beginning of the reaction and the efficiency of primer extension and amplification.
- Once established in the geometric phase, the reaction proceeds until one of the components becomes limiting.
- At this point, the yield of specific amplification products should be maximal, whereas nonspecific amplification products should be barely detectable, if at all.
- This is generally the case after ~30 cycles in PCRs containing  $10^5$  copies of the target sequence and *Taq* DNA polymerase (efficiency ~0.7).

- At least 25 cycles are required to achieve acceptable levels of amplification of single-copy target sequences in mammalian DNA templates.

### 4.3. Applications of PCR

1. Medical applications
2. Infectious disease applications
3. Forensic applications
4. Research applications
5. Other applications

#### Medical applications

PCR has been applied to a large number of medical procedures:

- The first application of PCR was for **genetic testing**, where a sample of DNA is analyzed for the presence of genetic disease mutations.

Prospective parents can be tested for being genetic carriers, or their children might be tested for actually being affected by a disease.

DNA samples for Prenatal testing can be obtained by amniocentesis, chorionic villus sampling, or even by the analysis of rare fetal cells circulating in the mother's bloodstream.

PCR analysis is also essential to Preimplantation genetic diagnosis, where individual cells of a developing embryo are tested for mutations.

- PCR can also be used as part of a sensitive test for **tissue typing**, vital to organ transplantation. As of 2008, there is even a proposal to replace the traditional antibody-based tests for blood type with PCR-based tests.
- Many forms of cancer involve alterations to **oncogenes**.

By using PCR-based tests to study these mutations, therapy regimens can sometimes be individually customized to a patient.

#### Infectious disease applications

Characterization and detection of infectious disease organisms have been revolutionized by PCR:

- The **Human Immunodeficiency Virus** (or HIV), responsible for **AIDS**, is a difficult target to find and eradicate.

The earliest tests for infection relied on the presence of antibodies to the virus circulating in the bloodstream.

However, antibodies don't appear until many weeks after infection, maternal antibodies mask the infection of a newborn, and therapeutic agents to fight the infection don't affect the antibodies.

PCR tests have been developed that can detect as little as one viral genome among the DNA of over 50,000 host cells.

Infections can be detected earlier, donated blood can be screened directly for the virus, newborns can be immediately tested for infection, and the effects of antiviral treatments can be quantified.

- Some disease organisms, such as that for **Tuberculosis**, are difficult to sample from patients and slow to be grown in the laboratory. PCR-based tests have allowed detection of small numbers of disease organisms (both live or dead), in convenient samples.

Detailed genetic analysis can also be used to detect antibiotic resistance, allowing immediate and effective therapy.

The effects of therapy can also be immediately evaluated.

- The spread of a **disease organism** through populations of domestic or wild animals can be monitored by PCR testing.

In many cases, the appearance of new virulent sub-types can be detected and monitored. The sub-types of an organism that were responsible for earlier epidemics can also be determined by PCR analysis.

### Forensic applications

The development of PCR-based genetic (or DNA) fingerprinting protocols has seen widespread application in forensics:

- In its most discriminating form, **Genetic fingerprinting** can uniquely discriminate any one person from the entire population of the world.

Minute samples of DNA can be isolated from a crime scene, and compared to that from suspects, or from a DNA database of earlier evidence or convicts.

Simpler versions of these tests are often used to rapidly rule out suspects during a criminal investigation.

Evidence from decades-old crimes can be tested, confirming or exonerating the people originally convicted.

- Less discriminating forms of DNA fingerprinting can help in **Parental testing**, where an individual is matched with their close relatives.

DNA from unidentified human remains can be tested, and compared with that from possible parents, siblings, or children.

Similar testing can be used to confirm the biological parents of an adopted (or kidnapped) child. The actual biological father of a newborn can also be confirmed (or ruled out).

### Research applications

PCR has been applied to many areas of research in molecular genetics:

- PCR allows rapid production of short pieces of DNA, even when nothing more than the sequence of the two primers is known.

This ability of PCR augments many methods, such as generating hybridization probes for Southern or northern blot hybridization.

PCR supplies these techniques with large amounts of pure DNA, sometimes as a single strand, enabling analysis even from very small amounts of starting material.

- The task of **DNA sequencing** can also be assisted by PCR. Known segments of DNA can easily be produced from a patient with a genetic disease mutation.

Modifications to the amplification technique can extract segments from a completely unknown genome, or can generate just a single strand of an area of interest.

- PCR has numerous applications to the more traditional process of **DNA cloning**.

It can extract segments for insertion into a vector from a larger genome, which may be only available in small quantities.

Using a single set of 'vector primers', it can also analyze or extract fragments that have already been inserted into vectors.

Some alterations to the PCR protocol can **generate mutations** (general or site-directed) of an inserted fragment.

- **Sequence-tagged sites** is a process where PCR is used as an indicator that a particular segment of a genome is present in a particular clone.

The Human Genome Project found this application vital to mapping the cosmid clones they were sequencing, and to coordinating the results from different laboratories.

- An exciting application of PCR is the phylogenic analysis of DNA from **ancient sources**, such as that found in the recovered bones of Neanderthals, or from frozen tissues of Mammoths.

In some cases the highly degraded DNA from these sources might be reassembled during the early stages of amplification.

- A common application of PCR is the study of patterns of **gene expression**.

Tissues (or even individual cells) can be analyzed at different stages to see which genes have become active, or which have been switched off.

This application can also use Q-PCR to quantitate the actual levels of expression

- The ability of PCR to simultaneously amplify several loci from individual sperm has greatly enhanced the more traditional task of **genetic mapping** by studying chromosomal crossovers after meiosis.

Rare crossover events between very close loci have been directly observed by analyzing thousands of individual sperms.

Similarly, unusual deletions, insertions, translocations, or inversions can be analyzed, all without having to wait (or pay for) the long and laborious processes of fertilization, embryogenesis, etc.

### Others applications

PCR is also important in answering basic scientific questions.

- In the field of evolutionary biology, PCR has been used to establish relationships among species.
- In anthropology, it has used to understand ancient human migration patterns.
- In archaeology, it has been used to help identify ancient human remains.
- Paleontologists have used PCR to amplify DNA from extinct insects preserved in amber for 20 million years.

- The Human Genome Project, which had a goal of determining the sequence of the 3 billion base pairs in the human genome, relied heavily on PCR.
- The genes responsible for a variety of human diseases have been identified using PCR.
- For example, a PCR technique called multiplex PCR identifies a mutation in a gene in boys suffering from Duchenne muscular dystrophy.
- PCR can also be used to search for DNA from foreign organisms such as viruses or bacteria.

## **5. DNA sequencing**

### **Methods of sequencing:**

Two different methods are now routinely used for determination of DNA sequences


1. Chemical degradation method
2. Chain termination method

### **5.1. Maxam and Gilbert's Chemical Degradation Method**

- developed in the late 1970's, by Allan Maxam and Walter Gilbert
- first method to determine the sequence of a DNA molecule of up to 500 bp.
- based on chemical modification of DNA and subsequent cleavage at specific bases.

#### **In this method, following steps are involved:**

- (i) Label the 3'ends of DNA with  $^{32}\text{P}$ .
- (ii) Separate two strands, both labelled at 3'ends.
- (iii) Divide the mixture in four samples, each treated with a different reagent having the property of destroying either only G, or only C, or 'A and G' or 'T and C'. The concentration of reagent is so adjusted that 50% of target base is destroyed, so that fragments of different sizes having  $^{32}\text{P}$  are produced.
- (i) Electrophoreses each of the four samples in four different lanes of the gel.
- (ii) Autoradiograph the gel and determine the sequence from positions of bands in four lanes.
- vi) Read the sequence based on autoradiography.




### Disadvantages of this method:

This technology has the disadvantage of relying on toxic chemicals.

### 5.2. Chain termination method

- Developed by Fred Sanger
- Also called as Dideoxynucleotide Synthetic Method
- utilizing single stranded DNA as template for DNA synthesis, in which 2',3' dideoxynucleotides were incorporated leading to termination of DNA synthesis.
- These dideoxynucleotides are used as triphosphate and can be incorporated in a growing chain, but these dideoxynucleotides terminate synthesis, since they can not form a phosphodiester bond with next incoming deoxynucleotide triphosphate (dNTP).

### Dideoxynucleotides:




### Following steps are involved in Sanger's dideoxy method for DNA sequencing:

(i) Four reaction tubes are set up, each containing single stranded DNA sample to be sequenced, all four dNTPs (radioactively labelled) and an enzyme for DNA synthesis (DNA polymerase I). Each tube also contains a small amount (much smaller amount relative to four dNTPs) of one of the four ddNTP, so that four tubes have each a different ddNTP, bringing about termination at a specific base adenine (A), cytosine (C), guanine (G) and thymine (T).

ii) The fragments, generated by random incorporation of ddNTP leading to termination of reaction, are then separated by electrophoresis

Polyacrylamide

This is done for mixtures on gel.




then separated by on a high resolution gel.

all the four reaction adjoining lanes in the

(iii) The gel is used for autoradiography so that the position of different bands in each lane can be visualized.

(iv) The bands on autoradiogram can be used for getting the DNA sequence.

### Example



Now, variations of the above dideoxy method was developed as automatic sequencers.

#### **In this automatic sequencer approach,**

- a different fluorescent dye is tagged to the oligonucleotide primer in each of the four reaction tubes (blue for: A, red for C, etc.).
- The four reaction mixtures are pooled and electrophoreses together in a single Polyacrylamide of the tube.
- A high sensitivity fluorescence detector, placed near the bottom of the tube, measures the amount, of each fluorophore as a function of time.
- The sequence is determined from the temporal order of peaks corresponding to four different dyes.

The chain termination method of DNA sequencing can only be used for short strands (100 to 1000 base pairs), longer sequences must be subdivided into smaller fragments, and subsequently re-assembled to give the overall sequence.

### Chromosome walking

- The principle of chromosome walking is that overlapping clones will hybridize to each other, allowing them to be assembled into a contiguous sequence.
- This can be used to isolate genes whose function is unknown but whose genetic location is known, a technique known as *positional cloning*.

To begin a chromosome walk, it is necessary to have in hand a genomic clone that is known to lie very close to the suspected location of the target gene.

For example - In humans,

- this could be a restriction fragment length polymorphism that has been genetically mapped to the same region.
- This clone is then used to screen a genomic library by hybridization, which should reveal any overlapping clones.
- These overlapping clones are then isolated, labeled and used in a second round of screening to identify further overlapping clones,
- and the process is repeated to build up a contiguous map.

### 1.2.3. Screening by PCR

The PCR is widely used to isolate specific DNA sequences from uncloned genomic DNA or cDNA, but it also a useful technique for library screening.

As a screening method,

- PCR has the same versatility as hybridization, and the same limitations.
- It is possible to identify any clone by PCR but only if there is sufficient information about its sequence to make suitable primers.

To isolate a specific clone, PCR is carried out with gene-specific primers that flank a unique sequence in the target.

- Instead of plating the library out on agar, as would be necessary for screening by hybridization, pools of clones are maintained in multiwell plates.
- Each well is screened by PCR and positive wells are identified.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: IV

BATCH-2017-2021

- The clones in each positive well are then diluted into a series in a secondary set of plates and
- screened again.
- The process is repeated until wells carrying homogeneous clones corresponding to the gene of interest have been identified.

There are also several applications where the use of *degenerate primers* is favourable.

- A degenerate primer is a mixture of primers, all of similar sequence but with variations at one or more positions.
- This is analogous to the use of degenerate oligonucleotides as hybridization probes, and the primers are synthesized in the same way.
- A common circumstance requiring the use of degenerate primers is when the primer sequences have to be deduced from amino acid sequences (Lee *et al.* 1988).

Degenerate primers may also be employed to search for novel members of a known family of genes or to search for homologous genes between species.

UNIT-V

SYLLABUS

**UNIT-V**

**Applications of Genetic Engineering:** In plants: use of *Agrobacterium tumefaciens* and *A. rhizogenes*, Ti plasmids, Strategies for gene transfer to plant cells, Direct DNA transfer to plants, Gene targeting in plants, Use of plant viruses as episomal expression vectors.

**In animals:** Production and applications of transgenic mice, role of ES cells in gene targeting in mice, Therapeutic products - blood proteins, human hormones, immune modulators and vaccines (one example each). Ethical, legal and social issues.

**Applications of recombinant DNA technology**

**Production of recombinant proteins**

- One potential of recombinant DNA is the production of proteins.
- Many systems have been designed for the expression of recombinant proteins.
- Although *E. coli* are easy to grow in large quantities there are some problems associated with the expression of eukaryotic proteins in prokaryotes.
- Therefore, expression systems in eukaryotic systems have also been devised.

The following are the general considerations in the over production of recombinant proteins.

- strong promoter
- regulatable promoter
- gene dosage
- localization signals
- protease defective hosts
- fusion proteins

Ideally,

- the recombinant protein should be cloned in conjunction with a strong promoter.

Promoters are elements found on the 5'- end of genes and control their expression.

- A strong promoter results in a high level of gene expression.
- In addition to being strong, promoters should be regulatable so that transcription can be turned off and on.

- Recombinant proteins are sometimes toxic for the host cell and this toxicity can be minimized by controlling the expression.
- The use of high copy number plasmids also increases the production of recombinant proteins by increasing the gene dosage.
- it is often convenient to have recombinant proteins that are exported from the host cell. This can be accomplished by engineering localization signals within the recombinant protein so that they are directed to a particular cellular compartment.
- In addition, the recombinant protein can also be expressed as a fusion protein with another protein. This will sometimes stabilize the recombinant protein and/or assist in the purification and characterization of the recombinant protein.
- Proteolysis of recombinant proteins is often a major problem that is partially alleviated by cloning into protease deficient hosts.

### Expression in bacterial cell

The general procedure for expression of cloned genes in *E. coli* involves

- the insertion of the coding region of interest into a vector, usually a plasmid, so the region is efficiently transcribed and translated.
- Since eukaryotic genes do not contain the proper signals for transcription initiation, ribosome recognition, translation initiation, and translation termination, these signals need to be supplied by the vector.

**Common promoters** used in bacterial expression vectors are:

$\lambda$  PL, lac, tac and T7

Of these, the lac and tac promoters are the most widely used.

**Control elements** that regulate expression from these promoters are supplied by either the host or the vector.

The  **$\lambda$  PL promoter** is controlled by a mutant cI repressor protein that is non-functional at 42°C.

- At temperatures less than 42°C the repressor protein (cI) binds to the promoter and prevents expression.
- To induce expression the temperature is raised so that the repressor becomes nonfunctional and expression is now permitted.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: V

BATCH-2017-2021


- One problem with this system is that heat-shock proteins may also be induced.

The **T7 promoter** is from the T7 phage

- is only transcribed by the T7 RNA polymerase (T7 gene 1).
- The host cell must also contain the T7 gene 1 in order to express from this promoter.
- To regulate expression from this promoter it is necessary to be able to regulate the expression of the T7 gene 1.

The **lac** and **tac** promoters

- are controlled by the lac repressor.



- The lac repressor binds to the lac promoter and prevents RNA polymerase from transcribing the gene. Host strains with lacIq gene express the lac repressor at 10-fold higher concentrations than the normal lacI gene.
- IPTG (isopropyl-1-thio-β-D-galactopyranoside, an analog of lactose) binds to the repressor and prevents its interaction with the lac promoter and allowing RNA polymerase to transcribe the regulated gene.
- The tac promoter is a fusion of trp and lac promoters and is also regulated by IPTG.

The production of recombinant **fusion proteins** often stabilizes the expression of foreign proteins in *E. coli*.

Several plasmids that express recombinant proteins as fusion proteins have been developed (Table).

### Fusion Protein

### Affinity Matrix

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: V

BATCH-2017-2021

Glutathione-Transferase	glutathione agarose
Thioredoxin	phenylarsine oxide-agarose
Maltose Binding Protein	amyloseagarose
Six Histidine Residues	Ni-agarose

In addition to stabilizing the recombinant protein, the fusion partner is often exploited for affinity purification or for the analysis of the recombinant protein.

- Recombinant proteins expressed at high levels will sometimes form insoluble aggregates known as **inclusion bodies**.
- In some applications it is possible to take advantage of this phenomenon.
- For example, the inclusion bodies can be isolated by differential centrifugation and solubilized under denaturing conditions (eg., urea). It is sometimes possible to renature the protein and regain activity. In addition, fusions with *E. coli* thioredoxin can circumvent inclusion body formation.

### Expression in eukaryotes

- Although expression of recombinant proteins in *E. coli* is usually fairly straight forward, it is often desirable or necessary to express cloned genes in eukaryotes.
- Eukaryotic expression systems are often needed to insure correct folding and disulfide-bond formation, post-translational modifications, and processing.
- Yeast, such as *Saccharomyces cerevisiae* and *Pichia* species, are useful hosts for the expression of recombinant proteins since they can be grown and manipulated like a bacteria.
- *Pichia* expression systems often allow for high level expression of recombinant proteins using a strong alcohol oxidase promoter which is induced by methanol.

**Shuttle vectors** contain both a bacterial origin of replication and the origin of replication of interest.

This allows for manipulations of the plasmids to be carried out in *E. coli* before transforming yeast or other eukaryotes.

The two major strategies for expression of recombinant sequences in mammalian cells are

- 1) stable or transient expression of transfected DNA and
- 2) the use of viral expression vectors.

Mammalian and other eukaryotic cells are able to take up DNA.

Several different methods for introducing DNA to cells are available

- calcium phosphate
- DEAE-dextran
- electroporation
- liposomes
- protoplast fusion
- ballistics (gene gun)
- microinjection

Several different **eukaryotic viruses** can also be used as cloning vectors for the expression of recombinant proteins.

- Lytic viruses are good for transient expression whereas episomal viruses are better for constitutive expression.
- Retroviruses can become incorporated into the host cell genome and possibly lead to a stable transformation.

**Baculoviruses** are used for the production of recombinant eukaryotic proteins.

- *Autographica californica* is a nuclear polyhedrosis virus (AcNPV) that infects insects.
- Sf9 cells, derived from *Spodoptera frugiperda*, are readily grown in vitro and can be infected with baculovirus.

#### **Insect cells**

- Extremely high levels of recombinant proteins can be expressed and many eukaryotic post-translational modifications are correctly made in the host insect cells.
- This high level of expression is driven by a strong promoter for the polyhedrin gene.

- The polyhedron protein is expressed late in infection as the virus is killing the host cell and is needed for the dissemination of the virus in nature.
- However, the polyhedrin gene is unnecessary for viral growth in tissue culture and can be replaced with a gene of interest.

## **2. Practical Applications of DNA Technology**

### **Medicine and the Pharmaceutical Industry**

#### **Diagnosis of diseases**

- It is possible to clone entire DNA molecules or genes associated with various diseases (e.g., HIV DNA; genes associated with hemophilia, cystic fibrosis, Huntington's disease, and Duchenne muscular dystrophy)
- These can be cloned in vectors or by PCR to obtain large amounts, and then used to produce probes. Probes can be used to detect infection (e.g., HIV) or the presence of a defective allele.
- Thus, it is possible to identify individuals carrying certain disease-related alleles even if they show no symptoms of the disease (e.g., heterozygous for cystic fibrosis)
- Even if a disease-causing allele has not been cloned and its precise locus is unknown, its presence can sometimes be detected by testing for restriction fragment length polymorphism (RFLP) markers that are very close to the gene in question

#### **Human gene therapy**

- For any genetic disorder traceable to a defective allele, it should theoretically be possible to replace or supplement the defective allele with a functional, normal allele using recombinant DNA techniques
- For gene therapy of somatic cells to be permanent, cells that receive the normal allele must be ones that multiply throughout the life of the patient
- Bone marrow stem cells are ideal candidates, because they continue to divide throughout life, and they give rise to cells of the blood and immune systems. If, for example, the patient's cells lacked a particular normal gene, it might be possible to insert the normal allele into a retrovirus

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: V

BATCH-2017-2021

- Let the retrovirus infect bone marrow cells that have been removed from the patient and cultured
- Allow the retrovirus to insert its DNA, along with the normal allele, into the chromosomal DNA of the patient's bone marrow cells
- Inject the engineered cells into the patient

### Pharmaceutical products

- By recombinant DNA technology discussed previously, human genes for various proteins have been engineered into expression vectors and then into bacterial host cells that produce and secrete the protein
- Examples include insulin, used to treat diabetes, and human growth hormone, used to treat hypopituitarism, which causes a form of dwarfism
- Another example is tissue plasminogen activator (TPA), which helps dissolve blood clots and reduces the risk of subsequent heart attacks if administered shortly after an initial attack
- It is also possible to construct desired molecules
- Genetically engineered proteins can block or mimic surface receptors on cell membranes; an example is a molecule designed to mimic a receptor protein that HIV binds to in entering white blood cells (if HIV binds to the drug molecules instead of those on the cell surface, it would fail to enter the blood cell)
- Recombinant DNA techniques can generate large amounts of proteins associated with the immune response against pathogens or be used to modify the genome of a pathogen to attenuate it, and thus lead to more specific and safer vaccines

### Forensic, environmental, and agricultural applications

#### Forensic use of DNA technology

- DNA fingerprinting involving RFLP analysis by Southern blotting
- Due to previous work, many RFLP sequences are well known, and probes to them have been developed
- These select portions of the total DNA serve as markers when the probes are used to tag the corresponding DNA fragment bands and visualize them via autoradiography

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: II BSC BT

COURSE NAME: Recombinant DNA Technology

COURSE CODE: 17BTU402

UNIT: V

BATCH-2017-2021

- The pattern of RFLP bands is unique to each individual; therefore, one can compare DNA from a crime scene (or, perhaps, an item associated with a crime, such as clothing) to DNA from a victim and DNA from a suspect
  - DNA fingerprinting involving satellite DNA analysis DNA fingerprinting is also done with satellite DNA consisting of tandemly repeated base sequences that vary in length from person to person
  - The most useful are microsatellites, which are roughly 10 to 100 base pairs long, have repeating units of only a few base pairs, and are highly variable from person to person
- For example, one person may have the sequence A-C-A repeated 10 times at one genome locus, 30 times at a second locus, and so on, while another person is likely to have different numbers of repeats at these loci
- Restriction fragments containing these simple tandem repeats (STRs) vary in size from one individual to another because of differences in STR lengths; therefore, they provide a unique pattern of markers for each individual
  - Actually, it is possible for two individuals who are not identical twins to have identical DNA fingerprints, but the chances range from one in 100,000 to one in a billion, and that is only because in most cases relatively few markers are examined

### **Environmental use of DNA technology**

- Considerable work is aimed at engineering organisms to cope with various environmental problems based on the ability of microorganisms to transform chemicals
- For example, the ability to extract heavy metals such as copper, lead, and nickel from the environment, which already exists in many bacteria, might lead to the development of ultra efficient genetically engineered microbes that could be used for mining minerals or cleaning up toxic mining wastes
- Likewise, genetic engineering is leading to the development of microbes that can degrade toxic compounds into nontoxic forms during the manufacturing process, at waste treatment facilities, at dump sites, and at sites of environmental disasters such as oil spills

### **Agricultural use of DNA technology**

### **Animal husbandry and “pharm” animals**

- Farm animals are regularly treated with engineered products such as vaccines, antibodies, and hormones
- Transgenic organisms (organisms that contain genes from another species) have been developed, such as fish containing a foreign growth hormone gene, which makes them grow faster
- Transgenic animals can also be engineered to produce large amounts of an otherwise rare biological substance for medical use; this can be done such that the gene's product is secreted in the animal's milk.

### **Genetic engineering in plants**

- It is possible to develop transgenic plants that contain genes for such things as resistance to herbicides, pathogens, pest insects, and freeze damage
- DNA containing the gene of interest is inserted into the T DNA area of the Ti plasmid from *Agrobacterium tumefaciens*
- The recombinant Ti plasmid is introduced into cultured plant cells where the T DNA carrying the gene of interest is inserted into the plant chromosomal DNA
- As the plant cell divides, each descendant receives a copy of the T DNA and the foreign gene
- Individual cells can then be induced to differentiate and regenerate an entire plant, all of the cells of which contain the foreign gene

### **3.Ethical issues involved in cloning**

- The success rate in cloning is quite low.
- Even if we can increase the odds of success, problems can arise during the clone's development, both before and after pregnancy.
- What are the possible implications of cloning to society?
- All of us - researchers, policymakers and the public - have a responsibility to explore the potential effects of cloning technologies on our lives so that we can make informed decisions.

For each new application of cloning technologies, we must consider:

- What are the benefits?
- What are the risks?
- Whom will the technology help? Does it have the potential to hurt anyone?
- What does this mean for me? For my family? For others around me?
- Why might others not share my view?

### **1.1. Ethical, legal and social issues**

There are several types of issues to consider as we think about cloning

- Ethical issues are those that ask us to consider the potential moral outcomes of cloning technologies
- Legal issues require researchers and the public to help policymakers decide whether and how cloning technologies should be regulated by the government.
- Social issues involve the impact of cloning technologies on society as a whole.

#### **Pros of Cloning**

- Cloning finds applications in genetic fingerprinting,
- amplification of DNA and alteration of the genetic makeup of organisms.
- It can be used to bring about desired changes in the genetic makeup of individuals thereby introducing positive traits in them, as also for elimination of negative traits.
- Cloning can also be applied to plants to remove or alter defective genes, thereby making them resistant to diseases.
- Cloning may find applications in development of human organs, thus making human life safer. Here we look at some of the potential advantages of cloning.

**Organ Replacement:** If the vital organs of the human body can be cloned, they can serve as backup systems for human beings. Cloning body parts can serve as a lifesaver. When a body organ such as a kidney or heart fails to function, it may be possible to replace it with the cloned body organ.

**Substitute for Natural Reproduction:** Cloning in human beings can prove to be a solution to

infertility. Cloning can serve as an option for producing children. With cloning, it would be possible to produce certain desired traits in human beings. We might be able to produce children with certain qualities.

**Help in Genetic Research:** Cloning technologies can prove helpful to researchers in genetics. They might be able to understand the composition of genes and the effects of genetic constituents on human traits, in a better manner. They will be able to alter genetic constituents in cloned human beings, thus simplifying their analysis of genes. Cloning may also help us combat a wide range of genetic diseases.

**Obtain Specific Traits in Organisms:** Cloning can make it possible for us to obtain customized organisms and harness them for the benefit of society. Cloning can serve as the best means to replicate animals that can be used for research purposes. Cloning can enable the genetic alteration of plants and animals. If positive changes can be brought about in living beings with the help of cloning, it will indeed be a boon to mankind.

### Cons of Cloning

Like every coin has two sides, cloning has its flip side too.

- Though cloning may work wonders in genetics, it has potential disadvantages.
- Cloning, as you know, is copying or replicating biological traits in organisms.
- Thus it might reduce the diversity in nature.
- it is not clear whether we will be able to bring all the potential uses of cloning into reality.

**Detrimental to Genetic Diversity:** Cloning creates identical genes. It is a process of replicating a genetic constitution, thus hampering the diversity in genes. While lessening the diversity in genes, we weaken our ability of adaptation. Cloning is also detrimental to the beauty that lies in diversity.

**Invitation to Malpractices:** While cloning allows man to tamper with genetics in human beings, it also makes deliberate reproduction of undesirable traits, a probability. Cloning of body organs might invite malpractices in society.

***Will this Technology Reach the Common Man?:*** In cloning human organs and using them for transplant, or in cloning human beings themselves, technical and economic barriers will have to be considered. Will cloned organs be cost-effective? Will cloning techniques really reach the common man?

***Man, a Man-made Being?:*** Moreover, cloning will put human and animal rights at stake. Will cloning fit into our ethical and moral principles? Cloning will make man just another man-made being. Won't it devalue mankind? Won't it demean the value of human life?

Questions	opt1	opt2	opt3	opt4	Answer
<b>Unit I</b>					
Enzymes used to degrade nucleic acid is called as	Nucleases	Ligases	Polymerases	Modifying enzymes	Nucleases
The enzyme used to add a methyl group to the DNA is called	Amylase	Protease	Methylase	Ligases	Methylase
The enzyme used to joining nucleic acid molecules is	Topoisomerases	Exonuclease	Endonuclease	Ligases	Ligases
Which enzyme used to make copies of nucleic acid molecules	Nucleases	Ligases	Polymerases	Modifying enzymes	Polymerases
How many types of nucleases are present?	One	Two	Three	Four	Two
Which enzyme used to remove or add chemical groups?	Nucleases	Ligases	Polymerases	Modifying enzymes	Modifying enzymes
Which enzyme used to remove nucleotides from the ends of the DNA molecule?	Endonuclease	Exonuclease	Topoisomerases	Ligases	Exonuclease
The enzyme that introduce or remove supercoils from covalently closed circular DNA is	Topoisomerases	Exonuclease	Endonuclease	Ligases	Topoisomerases
The functions of Endonuclease is	Cleaving the DNA molecule	Add a methyl group to the DNA	Remove or add chemical groups to DNA	Introduce or remove supercoils from	Cleaving the DNA molecule
How many types of Restriction endonucleases are present?	One	Two	Three	Four	Three
Which types of Restriction endonucleases require Mg <sup>2+</sup> ions for cleavage?	Type I	Type II	Type III	Type IV	Type II
How many types of DNA ligases are available?	One	Two	Three	Four	Four
Which enzyme helps to catalyse the polymerization of deoxyribonucleotides into a DNA strand ?	Restriction endonucleases	DNA polymerase	RNA polymerase	Exonuclease	DNA polymerase
DNA polymerase III enzyme synthesizes at a rate of _____ nucleotides per second	1000	2000	3000	5000	1000
The enzyme that produces RNA is called	Restriction endonucleases	DNA polymerase	RNA polymerase	Exonuclease	RNA polymerase
How many types in DNA polymerases are present in prokaryote?	One	Two	Three	Four	Two
How many types in DNA polymerases are present in eukaryote?	One	Two	Three	Four	Three
Which organism produces Taq DNA polymerase	Thermus	E.coli	Bacillus spp	Pseudomonas spp	Thermus aquaticus
Recognition sequence for Eco RI	AATTCCTTAA	AG/CT	TC/GA	GA/TC	GA/TC
What is the product produced by reverse transcriptase	single strand DNA	double strand DNA	single strand RNA	double strand RNA	single strand DNA
Restriction enzymes cleave DNA at-----	Specific nucleotide	Interior part of nucleotide sequence	Ends of nucleotide sequence	middle part of sequence	Specific nucleotide sequence
Endonucleases are enzymes that cleaves DNA at	Defined sequence	3' end of nucleotide	Internal position in random manner	external position	Internal position in random manner
Term endonucleases was coined by-----	Lederberg and Meselson	Lederberg and tatum	Lederberg and yaun	Smith and Nathans	Lederberg and Meselson
Restriction enzyme from Escherichia coli K 12 was first isolated from	Meselson and	Lederberg and	yaun	Meselson	Meselson and yaun
Enzymes were classified by	Smith and	arber	rodriguez	Boliver	Smith and Nathans
Type I restriction enzymes cleave DNA at	The point of recognition	1000 to 5000 nucleotides away	25 base pairs away	30 base pairs away	1000 to 5000 nucleotides away
co factor needed for type II restriction enzyme were	ATP and Mg <sup>2+</sup>	S- Adenosyl	Fe	DNP+	ATP and Mg <sup>2+</sup>
Type I restriction enzyme have	3 subunit	5 subunit	2 subunit	1 subunit	3 subunit
Type I restriction enzymes are isolated from the organisms	<i>Escherichia coli</i> B	Virus	pig	<i>Escherichia coli</i> D am	<i>Escherichia coli</i> B
Type II restriction enzymes cleaves DNA at	Defined recognition site	Random sites	25 bases away	1000 basepairs away	Defined recognition site
Smith and Nathans received nobel prize during	1973	1978	1980	1981	1978
Type II restriction enzymes cleaves DNA at	tetra nucleotide sequence	Penta nucleotide sequence	Hexa nucleotide sequence	tri nucleotide sequence	Penta nucleotide sequence
Ligases are called	Molecular	Molecular glue	Molecular tool	None of the above	Molecular glue
Ligases form phosphodiester bond between	3" hydroxyl and 5" phosphate	3" hydroxyl	5" hydroxyl	3" hydroxyl and 5" triphosphate	3" hydroxyl and 5" phosphate
Sources of Ligases are	Bacillus	Bacteriophage	Bacteriophage infected bacterium	retrovirus	Bacteriophage
Escherichia coli DNA ligase has the molecular weight of	73kDa	72 kDa	50kDa	35kDa	73kDa

Bacteriophage ligases have molecular weight of	68 kDa	70 kDa	60 kDa	95 kDa	68 kDa
Linkers used for producing sticky end have	Modified blunt end	A short nucleotide sequence with restriction site	Modified sticky end	Both a and c	A short nucleotide sequence with restriction site
Escherichia coli ligase seal nick at	3TC	4°C	SO-C	40-C	4°C
Adaptors have a .	3' OH and 5' P end	a short nucleotide sequence with a restriction site	Linker end	Homopolymer tails	3' OH and 5' P end
T4 bacteriophage ligase need	Mg <sup>2+</sup> as cofactor	ATP as cofactor	SAM as cofactor	NAD <sup>+</sup> as cofactor	ATP as cofactor
Alkaline phosphatase catalyze	Removal of phosphate at 5'terminal	Removal of 3" terminal hydroxyl group	Removal of 3 'terminal phosphate	None of the above	Removal of phosphate at 5'terminal
Alkaline phosphatase is a	Monomeric	HomoDimeric protein	Polymeric protein	Single glycoprotein	HomoDimeric protein
Alkaline phosphatase contain	Zn <sup>2+</sup>	Mg <sup>2+</sup>	Fe	Energy containing enzyme	Zn <sup>2+</sup>
The role of alkaline phosphatase is to	Prevent recircularisation	Induce ligation	Cleavage	Induction of recircularisation	Prevent recircularisation
Bacterial Alkaline Phosphatase action is inhibited by	Lysozyme	CTAB	EDTA	MgSO <sub>4</sub>	EDTA
Methylases methylate nucleotides at	C5 cytosine	Both C5cytosine and N6 adenine	Two adenine at N6 position	Two cytosine at C5 position	Both C5 cytosine and N6 adenine
Dam methylase used in	recombinational repair	Mismatch repair	SOS repair	Excision repair	Mismatch repair
Dam methylase are single nucleotide independent which under go	Type I modification/Restriction system	Type II modification/Restriction system	Type III modification/Restriction system	None	Type II modification/Restriction system
Dam methylasetransferase methylate at	.. .5" CCAGG 3"	.. .5" CCTGG 3"	randomly	.5 GATC .... 3	.. .5" CCAGG 3"
Source of methionine for methylation obtained from	S- Adenosyl Methionine	Methionine	Protein synthesis	None	S- Adenosyl Methionine
Polynucleotide kinase involves	Transfer of Hydroxyl group to the terminal end of DNA	Transfer of terminal phosphate group from ATP	catalyst of biochemical reaction	Prevents the transfer	Transfer of Hydroxyl group to the terminal end of DNA
DNA Polymerase I Holoenzyme are	Single functional enzyme	Bi functional enzyme	Multifunctional enzyme	None	Bi functional enzyme
A Polymerase Klenow fragment	Poses Polymerase	Poses 5" exonuclease activity	Lack exonuclease activity	have no activity	Poses Polymerase activity
T4 DNA Polymerase optimum temperature is	37 degree	45 degree	94 degree	50 degree	94 degree
Taq DNA Polymerase requires ..... Cofactor for activity	Mg <sub>2</sub> <sup>+</sup>	Na	Fe	Cu	Mg <sub>2</sub> <sup>+</sup>
Reverse transcriptase isolated from	Retrovirus	Adenovirus	Baculovirus	Gemini virus	Retrovirus
T4 bacteriophage ligase need	Mg <sup>2+</sup> as cofactor	ATP as cofactor	SAM as cofactor	NAD <sup>+</sup> as cofactor	ATP as cofactor
Which organism produces Taq DNA polymerase	Thermus	E.coli	Bacillus spp	Pseudomonas spp	Thermus aquaticus
EcoRI is isolated from	Bacillus	Pseudomonas	<i>Escherichia coli</i>	vibrio	<i>Escherichia coli</i>

<b>unit II</b>					
plasmids are extra chromosomal DNA that vary in size from	1Kb to More than 200kb	2Kb to More than 100kb	5Kb to More than 200kb	More than 300kb	1Kb to More than 200kb
Plasmid was coined by	Maccarthy	Avery	Lederberg	Smith	Lederberg
Plasmid carries	PMBl replicon	PI5A replicon	ColEi replicon	PkN402	PMBl replicon
The copy number of plasmid was controlled by	Host chromosome	Regulatory genes	Motar genes	Replicons	Replicons
Plasmids are commonly found in	Algae	Virus	Protozoa	Prokaryotes	Prokaryotes
The term episome was coined by	A.Jacob and Wollman	Lederberg	Watson	Collins	A.Jacob and Wollman
PBR322 plasmid vectors have	Amp resistance gene	kan resistance gene	Lac z gene	X-gal	Amp resistance gene
Runaway vectors replicate upto	34°C	40°C	94°C	noc	34°C
Eukaryotes that harbour a plasmid	Fusarium	Saccharomyces cerevisiae	Pencillium	Aspergillus	Saccharomyces cerevisiae
There are two types of naturally occurring plasmids	Col plasmid and F plasmid	E plasmid	E.coli	PBR322	Col plasmid and F plasmid
Conjugation is biological phenomena observed in the bacterial genetics directly by	Functional genes	Transfer genes and Mobilizing genes	Nick gene	Basis of mobilizing genes	Transfer genes and Mobilizing genes
Transfer genes have cluster of	12 different genes	11 different genes	10 different genes	20 different genes	12 different genes
Mob function defined by	3 regions of plasmid DNA	5 regions of plasmid DNA	7 regions of plasmid DNA	2 regions of plasmid DNA	2 regions of plasmid DNA
The bacteria associated with the discovery of R plasmids	Bacillus	Pseudomonas	Shigella	Salmonella	Shigella
Non-conjugative plasmids have	tra+/ mob+ genes	tra-/ mob- genes	tra-/ mob+ genes	Mob+/ mob+ genes	tra-/ mob+ genes
Plasmids carrying genes, which is responsible for antibiotic resistance known as	R plasmids	Cancer inducing	Col E 1 plasmid	F plasmid	R plasmids
Plasmid contain autonomous replication (rep) includes	Origin of	chemicals	Copy number	genes	Origin of replication
The plasmid that maintains low copy number in a cell	Stringent plasmid	Relaxed plasmid	Conjugative plasmids	None of the above	Stringent plasmid
The genes for drug resistance is initially coded on	PBR322	Transposons	Yeast plasmid	None of the ooove	PBR322
PBR322 composed of	3 section	2 section	1 section	5 section	3 section
Tn3 from transposon for PBR322 have	Tetracyclin resistant	Streptomycin resistant	Ampicillin resistant	Chloramphenicol resistant	Ampicillin resistant
The best-studied Col plasmid is	ColA	ColB	ColE	ColD	ColE
The replication of DNA is initiated at	2534 position of PBR322	2000 position of PBR322	2435 position of PBR322 .	1000 position of PBR322	2534 position of PBR322
One gene important for ampicillin resistant in PBR322 is	D galactose gene	P -galactose gene	B lactamase gene	B- Lactase gene	B- lactamase gene
PBR 322 vectors are restricted to	Gram positive bacteria	Gram negative bacteria	Archae bacteria	Eubacteria.	Gram negative bacteria
Lambda DNA has a genetic material as	ds DNA	Ss DNA	ds RNA	ssRNA	Ss DNA
Lambda genome has a length of	48502bp	33402bp	40000bp	50,435bp	48502bp
The non-essential region of lambda phage is	int,xis,mob	int,xis,att	tra,mob,xis	B2 region	B2 region
DNA replication during lytic growth required for the lysis of cellular membrane	Sand R	Rand T	Sand Q	Q and U .	Sand R
Vectors, which contains a unique site (x) for the insertion of foreign DNA have been designated	Expression vector	Replacement vector	Insertional vector	Shuttle vector	Insertional vector
λgt10 carry up to	8 Kb of new DNA	10kb new DNA	15kb new DNA	20 kb new DNA	10kb new DNA
λgt10 E COR1 site is located in	R genes	C 1 genes	b 2 genes	D genes	C 1 genes
λEMBL4 carries up to	10 Kb	20kb	15kb	25 kb	10 Kb
λEMBL4 selected on the basis of	Size /	Size / X gal Phenotype	Size / Non recombinants	None .	Size / Spiphenotype
The expression vector for somatostatin is	Psom II 3	P som II	P BH 20	P som I	Psom II 3
The viral genome of SV40 is	5243 bp	5842bp	3483bp	2800bp	5243 bp
Sv40 DNA is isolated from virus particles in the form of	Super helical DNA	Double Stranded Super helix DNA	RNA	Inner DNA	Super helical DNA

Early region codes for two partially overlapping genes, which direct the synthesis of	T -Antigen	t-antigen	R -antigen	r -antigen	T -Antigen
The function of T antigen	Initiation DNA replication	Control of its own transcription	translation control	Control of its own translation	Initiation DNA replication
Which organism is called as "Super Bug"	Escherichia coli	Bacillus spp	Pseudomonas putida	Streptococcus	Pseudomonas putida
Vir E codes for which protein that binds to T-Dna during transfer	Competence	SSB proteins	DSB proteins	Tyrosine	SSB proteins
PH V 33 Multiply both in	E.Coli and Bacillus	Bacillus and Streptomyces	Bacillus and Yeast	Bacillus and Pseudomonas	E.Coli and Bacillus
Agrobacterium tumefaciens infects	Monocotyledon Palnts	Dicotyledon Plants	trees	creepers	Dicotyledon Plants
Ti Plasmids are	Transgenic plasmids	Tumour inducing plasmids	Inhibiting plasmids	Haemorrhagic plasmids	Tumour inducing plasmids
In retrovirus gag codes for	Reverse transcriptase	Polymerase	Synthase	Envelop proteins	Reverse transcriptase
In retrovirus pol codes for	Reverse transcriptase	Polymerase	Synthase	Envelop proteins	Polymerase
In retrovirus env codes for	Reverse transcriptase	Polymerase	Synthase	Envelope proteins	Envelope proteins
Vectors, which contains two restriction sites for the insertion of foreign DNA have been	Expression vector	Replacement vector	Insertional vector	Shuttle vector	Replacement vector
The first step I to create recombinant AcMNPV is to generate	Transfer vector	Helper vector	Modified vector	Initiator sequence	Transfer vector
Copy number are either strictly controlled by correlating with the number of	Plasmid DNA	Chromosomal DNA	Insertional DNA	None of the above	Chromosomal DNA
During infection ----- forms of virions produced	One form	Two form	Three form	Fiveform	Two form
In second form the protein of matrix is called as -----	Monohedrin	Polyhedrin	Dihedrin	Trihedrin	Polyhedrin
The specific baculoviruss that has been used extensively as as expression vector is	Autographa califonia	Autographa mixa	Autographa indica	Autographa esculenta	Autographa califonia
pUC18 vector is ----- selection vector	Amp selection vector	Kan selection vector	Lac selection vector	Spi selection vector	Lac selection vector
PBR322 the word BR stands for	Berger	Bunsen rodergh	Beep run	Boliver and	Boliver and Rodriguez
pBR322 was developed in	1960	1980	1970	1990	1970
EM vector have	SP6 promoter	T7 Promoter	Both SP6 and T7	T8 Promoter	Both SP6 and T7
_____ is produced from E.coli	F plasmid and COL E plasmid	Ti plasmid	degradative plasmid	None of the above	F plasmid and COL E plasmid
T4 DNA polymerase optimum temperature is	37	45	55	94	94
Reverse transcriptase is isolated from	Retrovirus	Bacteria	Fungus	Aedinovirus	Retrovirus

<b>Unit III</b>					
Microinjection in mice was performed at	Male pronuclei	Female pronuclei	Blastocyst	Pronucleus	Male pronuclei
The protoplast fusion is done by	PEG method	Electrofusion	Sodium nitrate method	agrobactrium	PEG method
Microinjection technique proves direct delivery in	Intranuclear	Intracytoplasmic region	Intro cellular region	Intra species region	Intranuclear region
Macroinjection technique involved injection of DNA in to	Male pronuclei	Female pronuclei	Wound site within tissue-	Cytoplasm of cell	Wound site within tissue-
Liposome are defined by	Bongham	Winston	Hogness	Grunstein	Bongham
Liposome were first described for delivering	Polio virus RNA	Adenovirus RNA	Baculovirus RNA	Adenovirus RNA	Polio virus RNA
The plasma membrane of pro top lasts are dissolved in the presence of	Mg ions	Fe ions	Ca ions	None of the above	None of the above
The biolistic process currently in widest use employs gunpowder charge to drive	Plastic cylindrical macro projectile	Plastic cylindrical microprojectile	DNA-tungsten suspension	Blank cartridge	Plastic cylindrical microprojectile
The microprojectile are commonly DNA coated with	Gold or Tungsten	Silver or Molybdenum	Copper or ~inc	Lithium or Molybdenum	Gold or Tungsten
The nick translation was performed by	Making ss cuts in ds DNA molecule.	Making ds cuts in ds DNA molecule.	Making ss cuts in ss DNA molecule.	Making no cleavage in DNA molecule	Making ss cuts in ds DNA molecule.
The probes can be Non-radiolabelled by	Biotin	Toluene	Epinephrin	Globulins	Biotin
Northern hybridization is used to measure	The amount of RNA	Size of protein	Amount of DNA	Amount of micro satellites	The amount of RNA
The first description of northern blotting was published by	Alwine et al	EM Southern	Karry mullois	Hogness	Alwine et al
The most effective denaturing agent ofRNA is	Glyonol/	Formaldehyde	Dimethyl sulphide	Guanidine	Glyonol/ Formamide
The commonly used membrane in northern blotting are	Polyethylene and nylon	Nylon and nitrocellulose	Terylene and nitrocellulose	Nylon and terylene	Nylon and nitrocellulose
The original name of nylon is	Nylon	Polystyrene	Artificial wool	Filter 66	Polystyrene
Which of the following is tracking dye	Methylene blue	Malachite green	Bromophenol blue	Xylene cyanol	Methylene blue
The crucial step in northern blotting is	Isolation of m RNA	Transfer of denatured RNA from gel to membrane	Denaturation of mRNA	Seperation through AGE	Isolation of m RNA
Southern blotting technique was identified by	Southern (1975)	Kary Mullis (1985)	Lederberg (1994)	Guray (1998)	Southern (1975)
DNA blotting technique is used to study about	DNA	RNA	Proteins	Lipids	DNA
The upward and backward transfer of DNA into membrane in southern blotting was performed by	Upward capillary transfer	Downward capillary transfer	Bi-directional transfer	Vacuum transfer	Bi-directional transfer
The method used when the target DNA fragment when present in higher concentration is	Capillary method	Vacuum blotting	Bi-directional blotting	Transfer in Alkaline buffer	Vacuum blotting
The polysaccharide contained in agarose is	Sucrose	Galactose	Fructose	Maltose	Galactose
The glycosidic linkage in Agarose is	a (1,2) ~ (1,4)	a (1,2) ~ (1,3)	a (1,3) ~ (1, 4)	a (1,4) f3 (1, 6)	a (1,4) f3 (1, 6)
The rate of the movement of DNA is ----- as the size of molecule increases	Slow	Fast	No change	Scattering	Slow
The mostly preferred buffer in AGE	TAE	TBE	TPE	TE	TAE
The DNA should move from	Negative to	Positive to negative	Negative to neutral	Positive to neutral	Negative to Positive
Selection by complementation is the existence of	Auxotroph	Prototroph	Heterotroph	Lethal genes	Auxotroph
Nonsense suppression is a special case of complementation operating only in the instance of	Frame Shift mutation	Transition mutation	Transversion	Nonsense mutation	Nonsense mutation
Electroporation can be other words called as	Electropermeabilization	Induction	Dittusion	Localization	Electropermeabilization
Electropermeabilization, is a significant increase in	electrical conductivity and permeability of the cell plasma membrane	Damage to membrane	Destroys cell structure	None of the above	electrical conductivity and permeability of the cell plasma membrane
Pores are formed when the voltage across a plasma membrane	Exceeds its dielectric strength.	Exceeds its trielectric value	Falls below the dielectric value	Maintains at the dielectric value	Exceeds its dielectric strength.

Excessive exposure of live cells to electrical fields can cause	apoptosis and/or necrosis	Lineage	Pores	Condense mass	apoptosis and/or necrosis
In molecular biology, the process of electroporation is often used for the	transformation	recombination	congigation	transduction	transformation
Electroporation is done with	Electroporators	Magnetic stirrers	Sonoicators	Hot plates	Electroporators
Method (biological ballistics) is the method of transfecting cells by'	Bombarding them with Microprojectiles coated with DNA.	By mixing with DNA of interest	By encapsulation	By Precipitation	Bombarding them with Microprojectiles coated with DNA.
RNA can be generated from DNA by a process known as	Reverse transcription	Forward reversion	Mutation	Ligation	Reverse transcription
Biolistics has been a useful technique for	Testing expression of genes particularly	Testing expression of genes particularly in earthworm	Testing expression of genes particularly in drosophila	Testing expression of genes particularly in plant	Testing expression of genes particularly in plant
-----is the method used to 'burst of a gas,	Water flush	A capacitive electric discharge through a wire electrode	A burst of compressed carbon dioxide	A burst of compressed oxygen	A capacitive electric discharge through a wire electrode
Probes are short nucleotide of ----- in length	15 - 20 base pair	20 - 40 base pair	80 base pair	100 base pair	15 - 20 base pair
Which one of the following is not used in Non radiolabelling	Biotin	Horse radish peroxidase	Digitoxigenin	Cyanocobalamine	Cyanocobalamine
In ----- method the DNA should be pure,	Biolistic	Calcium phosphate	Lipofection	Electroporation	Biolistic
The marker gene present in PBR322 is	Ampr	Kamr	Camr	Metr	Ampr
The southern hybridization results was read by	UV	Autoradiography	Stainiing	Visible light	Autoradiography
Biolistic operation cost	351akhs	50lakhs	60lakhs	10lakhs	351akhs
The DNA should be mixed with -----in biolistic process	Calcium	Biotin	Digitoxigenin	Spermidine	Spermidine
----- is the disadvantage 'in ETBR attaining	Limit to sensitivity less	Limit to sensitivity less than 25ng	Limit to sensitivity less than 35ng	Limit to sensitivity less than 45ng	Limit to sensitivity less than 25ng
Marker inactivation of antibiotics can be studied by	Activity of antibiotics	Resistance of antibiotics	Both resistance and activity	None of the above	Resistance of antibiotics
----- is the sample used for nuclec acid blotting	DNA	RNA	Protein	Both DNA and RNA	Both DNA and RNA
For nucleic acid hybridization nucleic acid is digested with----- restriction enzyme	One	Two	More than two	Other than restriction enzymes	More than two
----- refers to blotting of electrophoresed protein bands from SDS PAGE to membrane	Southern blotting	Northern blotting	Western blotting	Dot blotting	Western blotting
Which technique purified nucleic acid sample directlyapplied to nitrocellulose filter	Southern blotting	Northern blotting	Western blotting	Dot blotting	Dot blotting
Joining of donor DNA fragments and vector DNA fragments with the help of DNA ligase enzyme is known as	Gene cloning	Splicing	Gene manipulation	Moleccular cloning	Gene cloning
Suitable host in genetic engineering to introduce DNA fragments of donor is	Bacillus subtilis	Escherechia coli	Bacteriophages	virus	Bacillus subtilis
A southern blot is a technique used to detect	proteins	DNA	RNA	None	DNA
Type of restriction endonucleases most useful in molecular cloning is:	Type I	Type II	Type III	None	Type II
Northern blots probes are	DNA and RNA	DNA	RNA and protein	RNA	RNA
Which is the most specific recombinant library?	Genomic	protein	cDNA	cRNA	cDNA
Recombinant DNA research is dependent on.....	Cloning	Hosts	Vectros	carrier	Vectros
The polymerase chain reaction is a technique that selectively replicate	RNA	DNA	Protein	None	DNA

<b>Unit IV</b>					
Gene knockout is a genetically engineered organism that carries	One or more genes in its	One or more genes in its cell	One gene of assortment	None of the above	One or more genes in its chromosomes
Knockout is a route to learning about a gene	That is active	That has been	That are suppressed	That are ordered	That are suppressed
Knockout is accomplished through	a combination of techniques	a single technique	Bifunctional technique	None of the above	a combination of techniques
Knockout require	a plasmid	bacterial artificial chromosome	gene	tissues	a plasmid
Gene therapy is the insertion of genes into an individual's	Bones	cells and tissues	Ligaments	Skull	cells and tissues
----- to treat a disease, and hereditary diseases	Gene therapy	Genetic instability	Genome stability	Transplantation	Gene therapy
A carrier called a ----- must be used to deliver the therapeutic gene to the patient's	Fusion agent	Trancription initiator	Vector	Illucitor	Vector
The most common type of vector are ----- that have been genetically altered to carry normal human DNA	PBR322	Cosmids	Viruses	Yeast	Viruses
All gene therapy to date on humans has been directed	somatic cells	Gene level	RNA level	Plasmid level	somatic cells
Somatic gene therapy can be broadly split in to	one category	Five category	Six category	Two categories	Two categories
<i>ex vivo, which means</i>	Cells are modified outside the body and then transplanted back	genes are changed in cells still in the body	recombination with a very low probability	Recombination approach	Cells are modified outside the body and then transplanted back in again
<i>vivo, which means</i>	Cells are modified outside the body and then transplanted back	genes are changed in cells still in the body	recombinatiorr with a very low probability	Recombination approach	genes are changed in cells still in the body
----- are the methods to replace or repair the genes targeted in gene therapy	A normal gene may be inserted into a nonspecific location within the genome to replace a nonfunctional gene. This approach is most	An abnormal gene could be swapped for a normal gene through homologous recombination	The abnormal gene could be repaired through selective reverse mutation, which returns the gene to its normal function	insertion of artificial gene	The abnormal gene could be repaired through selective reverse mutation, which returns the gene to its normal function
The genetic material in retroviruses is in the form	DNA molecule	RNA molecules	Proteims	Lipids	RNA molecules
Antisense refers to short	DNA sequences	RNA sequences	Both DNA and RNA sequences	Any chemical substances	Both DNA and RNA sequences
----- are designed to be complementary to a specific gene sequence to inhibit activity	Oligonucleotides	Gene fragments	Similar sequences	Antisense oligonucleotide	Antisense oligonucleotide
In principle, antisense technology is supposed to prevent	Protein production from a targeted gene	Amino acid synthesis	Mutate cells	Elongate the cell cycle	Protein production from a targeted gene
RNA interference is a mechanism	DNA -guide regulation of gene expression	RNA-guided regulation of gene expression	Reverse trancriptase - guided regulation mechanism	Conserved pathway modification	RNA-guided regulation of gene expression
In RNAi _____ inhibits the expression of genes	Viruses	Double stranded DNA	double-stranded ribonucleic acid	Bacterial DNA	double-stranded ribonucleic acid
RNAi pathway is thought to have evolved as a form of	innate immunityagainst viruses	Innate immunity against protozao	Innate immunity against bacteria	Innate immunity against fungi	innate immunityagainst viruses
The RNA interference pathway is often exploited in experimental biology	To study the activation of genes	To study the biology of gene	to study the function of genes in cell culture and in vivo in model organisms.	To study the history of genes	to study the function of genes in cell culture and in vivo in model

Vaccines that use components of a pathogenic organism rather than whole organism is called	Peptide vaccines	Triplet vaccines	MAB	Subunit vaccines	Subunit vaccines
Since RNAi may not totally abolish expression of the gene, this technique is sometimes referred	knock out	knock Down	knock up	Knock in	knock Down
In subunit vaccine for Herpes simplex virus are cloned into	Chines hamster ovary	Lung cell lines	Mamalian cell lines	Hela cell lines	Chines hamster ovary
The genome of FMDV consist of	Single stranded DNA	Single stranded RNA	Double stranded DNA	Double stranded RNA	Single stranded RNA
Hexadimeric enterotoxin of Vibrio cholerae made of	Single A and 5 identical B	Single B and identical a subunit	Single alpha subunit	Sinale beta subunit	Single A and 5 identical B subunit
There are ----- of interferons	Two major classes	Three major classes	Five major classes	Six major classes	Three major classes
Golden rice is a variety	<i>Oryza glaberrima</i>	<i>Oryza sativa</i>	<i>Oryza triticum</i>	<i>Oryza annuum</i>	<i>Oryza sativa</i>
( <i>Oryza sativa</i> . produced through genetic engineering to biosynthesize the precursors of	beta-carotene pro-vitamin A	Alpha - Pro-vitamin A	beta-carotene pro-Vitamin B	beta-carotene pro-vitamin c	beta-carotene pro-vitamin A
Beta carotein is present at ----- in the goldenrice	Capsule	Husk	Endospore	Stem	Endospore
Golden rice was created by transforming rice with	three beta-	Two beta - carotene	Eive beta - carotene	SiX beta carotene	three beta-carotene'
<i>Psy codes for ----- in golden rice</i>	Phosphoribosyl transferase	phytoene synthase	Phythose synthase	Phosphatase	phytoene synthase
In golden rice crtI gene is obtained from the soil bacterium	<i>Erwinia caratovora</i>	<i>Erwinia ungalculata</i>	<i>Erwinia uredova</i>	<i>Erwinia rhiza</i>	<i>Erwinia uredova</i>
----- causes second generation seeds to be sterile	Antisense technology	Mutational technology	Terminator seed technology	Knock out technology	Terminator seed technology
The coagulation is an important part of	Thrombosis	Emolbism	Hemostatis	Hemorrhage	Hemostatis
Eventually, all blood clots are reorganised and resorbed by a process termed	Plastolysis	Fibrynolysis	Rectolysis	Clustering	Fibrynolysis
Fibrynogen forms	Channels	Blood clots	Blood platelets	Permiations	Blood clots
Dicer is	chemical	Cofactor	Ribonulease enzyme	Ligase enzyme	Ribonulease enzyme
The RNAi pathway is initiated by the	Protein P	REC.	Dicer enzyme	None ofthe above	Dicer enzyme
Galactose dehydrogenase is isolated from	<i>Pseudomanas</i>	<i>E.Coli</i>	Bacillus	Corynebacteriumi	<i>Pseudomanas</i>
L aspariginase have----- property	Antitumour	Antialleric property	Allergic property	Dehydration	Antitumour property
PCR was first proposed by	Kary Mullis	Hargobind khurrana	James Watson	Barbara mcclintock	Kary Mullis
PCR is a technique of	DNA degradation	DNA amplification	DNA sequencing	DNA ligation	DNA amplification
The basic requirements of PCR reaction excludes	DNA	primers	heat stable DNA	restriction enzymes	restriction enzymes
The process of binding of primer to the denatured DNA strand is called as	Denaturation	Annealing	Renaturation	None of these	Annealing
The technique used to identify specific DNA sequence in bacterial colonies is	Colony	In situ hybridization	Dot blot	Western blotting	Colony hybridization
Asymmetric PCR is to generate	single stranded copies	double stranded copies	multiple copies	half copy	single stranded copies
Reverse transcriptase PCR uses	mRNA as template to form	protein as template to form DNA	DNA as tempelate to form protein	restriction enzymes	mRNA as template to form cDNA
Reverse transcriptase enzyme is	DNA dependent DNA polymerase	RNA dependent DNA polymerase	RNA dependent RNA polymerase	DNA dependent RNA polymerase	RNA dependent DNA polymerase
Nucleic acid hybridization is the most commonly used method because	Rapid	screen cDNA libraries	identifies clones	restriction enzymes	Rapid
First step of PCR is	Annealing	renaturation	denaturation	ligation	denaturation
Which of the following technique is used in DNA fingerprinting	Western blotting	Southern blotting	Variable blotting	Eastern blotting	Southern blotting
Labeled antibody are used to detect	DNA	RNA	protein	lipid	protein
Which of the following technique doesn't involve electrophoresis for the separation of	Dot blotting	Southern blotting	Northern blotting	Western blotting	Dot blotting
Probe is a	Protein	Short piece of labeled DNA	Short piece of labeled protein	enzyme	Short piece of labeled DNA
Nitrocellulose membrane is used for	Eastern blotting	Southern blotting	Northern blotting	Both B and C	Southern blotting
Reverse transcription PCR is used for analysis in	Forensic analysis of DNA	genomic DNA	mRNA	binding	mRNA
The molecular target for recognition in immunological screening is generally	Paratope	Epitope	ab	antigen	Paratope

In the Sanger method, what causes the termination of chain elongation	incorporation of DNA	dideoxy nucleotide	Denaturation	amplification	dideoxy nucleotide
Which of the following is a chemical nucleotide sequencing method	Sanger	Maxam-Gilbert's	Edmans	Automated	Maxam-Gilbert's

<b>Unit V</b>					
Which of the following statements are true regarding rDNA technology	obtain large number copies of specific DNA	physics technology	computer technology	chemical technology	obtain large number copies of specific DNA fragments
The first successful transformation of rDNA molecule into a bacterium was carried out by	Nathan, Arber and Smith	Watson, Crick and Wilkins	Boyer and Cohen	Paul Berg	Boyer and Cohen
Which are responsible for over production of recombinant protein	Strong promoter	Gene dosage	weak promoter	small promoter	Strong promoter
Promoters used in bacterial expression vector are	lac	BAX	TIC	LOCK	lac
Which of the following is used yeast as host	<i>Saccharomyces cerevisiae</i>	<i>E. Coli</i>	<i>Bacillus subtilis</i>	<i>Aspergillus niger</i>	<i>Saccharomyces cerevisiae</i>
Which of the following would be useful selectable marker	Antibiotic	DNA	protein	RNA	DNA
Recombinant proteins expressed at high levels will sometimes form insoluble aggregates known	Inclusion bodies	Exclusion bodies	partition bodies	microbodies	Inclusion bodies
The tac promoter is a fusion of	Trp and Hfr	Trp and lac	T7 and lac	Lac and y <sub>pl</sub>	T7 and lac
Shuttle vectors contain	Origin of replication of bacteria and	Origin of replication of eukaryote	marker	operon	Origin of replication of bacteria and eukaryotes
Recombinant insulin was expressed in	Bacterial expression system	Plant expression system	Insect expression system	Mammalian expression system	Mammalian expression system
Molecular markers include	RAPD	Satellite	SMP	GLP	RAPD
The variation in the restriction DNA fragment lengths between individuals of a species is called	Restriction fragment length	Random amplified polymorphic DNA	Amplified fragment length polymorphisms	Simple sequence repeats	Restriction fragment length polymorphism
Specific biomolecules which shows easily detectable differences among different strains of a species is termed as	DNA fingerprinting	Molecular marker	Molecular scissors	RFLP	Molecular marker
Locations of quantitative genes on chromosomes are called	Qualitative trait	Quantitative trait loci	allels	recon	Qualitative trait loci
DNA of eukaryotic organisms has several repeating units of short sequences called	Random repeats	Tandem repeats	Mini satellites	mega satellite	Tandem repeats
_____ issues are those that ask to consider the potential moral outcomes	Ethical	Legal	social	scientific	Ethical
_____ issues are those that help for policy making	Ethical	Legal	social	scientific	Legal
_____ issues are those that involves the impact of cloning technologies	Ethical	Legal	social	scientific	social
The part of Ti plasmid which get transferred to plant chromosome is	A DNA	B DNA	Z DNA	T DNA	T DNA
Ti plasmids are isolated from	<i>Staphylococcus aureus</i>	<i>Agrobacterium tumefaciens</i>	<i>Sacchromyces cerviciae</i>	<i>Escherichia coli</i>	<i>Agrobacterium tumefaciens</i>
Treatment of genetic diseases by introducing proper genes into patients cell is called	gene therapy	enzyme therapy	genetic engineering	protein engineering	gene therapy
Treatment of genetic diseases by introducing a remedial gene into sperm,egg or zygote is known as	somatic cell gene therapy	embryo therapy	germline gene therapy	antisense therapy	germline gene therapy
Somatic cell gene therapy involves	introduction of remedial gene into zygote	introduction of remedial gene into blood cells	introduction of remedial gene into sperm	introduction of remedial gene into egg	introduction of remedial gene into blood cells
Treatment of genetic diseases by introducing a remedial gene that prevents the expression of the specific defective gene is called	exvivo gene therapy	invivo gene therapy	germline therapy	antisense gene therapy	antisense gene therapy
The functional gene is introduced in addition to defective gene endogenous to the cells is known as	somatic cell gene therapy	augmentation gene therapy	targeted gene therapy	germline gene therapy	augmentation gene therapy
Transfer of cloned genes into cells grown in culture and then introduction of these cells into	invivo gene	embryo therapy	somatic cell gene therapy	exvivo therapy	exvivo therapy
Synthetic vesicles composed of lipid bilayer used for transfer of genes is	lipoproteins	liposomes	lysosomes	lysozymes	liposomes
Binding of gene specific oligonucleotides to double stranded DNA to inhibit transcription of a gene is	triple helix therapeutics	antisense therapy	ribozyme therapeutics	antisense oligonucleotide	triple helix therapeutics
Self splicing RNA molecules are known as	exons	introns	ribozymes	ribosomes	ribozymes
<i>Agrobacterium tumefaciens</i> infect _____ family	roseaceae	solanaceae	liliaceae	malvaceae	roseaceae
The portion of Ti plasmid transferred to the chromosomal DNA of infected cells is called as	vir gene	T-DNA	Ori c	Tra gene	T-DNA
Substance released by plants in response to wounding and can induce vir genes of Ti plasmid is	opines	acetosyringone	nopaline	auxin	acetosyringone
The genes which help in transfer of T-DNA	Tra genes	vir genes	Occ genes	Noc gene	vir genes

Genes of Ti plasmid that help in auxin biosynthesis is	iaaH	iaaM	ipt	both a and b	both a and b
The vir gene responsible for topoisomerase activity	vir D	vir E	vir B	vir G	vir D
The genes which produces toxic crystals in Bacillus thuringiensis is	Vir genes	Cry genes	Nif genes	Tra genes	Cry genes
The toxic crystals in B.thuringiensis kills the insect by	causing pores in epithelial cells	by blocking respiration	by inhibition of metabolic enzymes	by inhibiting production of eggs	causing pores in epithelial cells
Genes coding for ..... are integrated into plants for preventing fruit ripening and softening during transport	poly galacturonase	pectinase	ethylene	auxin	poly galacturonase
The gene coding for..... is used to produce phosphinothricin tolerant plants	phosphinothricin	phosphinothricin acetyl transferase	phosphinothricin butyl transferase	phosphino acetyl transferase	phosphinothricin acetyl transferase
T7 promoter is derived from the .....phage	λ phage	T7 phage	M13 Phage	pUC phage	T7 phage
The insoluble recombinant protein aggregates are known as	Inclusion bodies	Fusion proteins	Affinity proteins	Protein clot	Inclusion bodies
A strong promoter results ..... level of gene expression	Low	High	Medium	very low	High
Which of the following is used yeast as host	<i>Saccharomyces cerevisiae</i>	<i>E. Coli</i>	<i>Bacillus subtilis</i>	<i>Aspergillus niger</i>	<i>Saccharomyces cerevisiae</i>
Selectable marker in this is	Antibiotic	DNA	protein	RNA	DNA
During high levels of expression recombinant proteins will sometimes form insoluble aggregates	Inclusion bodies	perfusion bodies	foreign bodies	macro bodies	Inclusion bodies
The fusion combination of Tac promoter is	Trp and Hfr	Trp and lac	T7 and lac	Lac and ypl	T7 and lac
The Shuttle vectors contain	Origin of replication of	Origin of replication of eukaryote	Origin of replication of bacteria and eukaryote	promoter	Origin of replication of bacteria and eukaryote
Expression system for recombinant insulin is	Bacterial expression system	Plant expression system	Insect expression system	Mammalian expression system	Bacterial expression system
Pick the molecular markers in this	RFLP	RFPD	ALLP	SMP	RFLP
Restriction DNA fragment lengths variation between individuals of a species is called	Restriction fragment length	Random amplified polymorphic DNA	Amplified fragment length polymorphisms	Simple sequence repeats	Restriction fragment length polymorphism
Easily detectable differences showed by specific biomolecules among different strains of a species is termed as	DNA fingerprinting	Molecular marker	Molecular scissors	RFLP	Molecular marker
Quantitative genes location on chromosomes are called	Qualitative trait	Quantitative trait loci	quantam trait loci	parallel gene	Qualitative trait loci
First step of PCR is	Annealing	renaturation	denaturation	ligation	denaturation
Technique used in DNA fingerprinting	Western blotting	Southern blotting	Variable blotting	Eastern blotting	Southern blotting
In blotting Labeled antibody are used to detect	DNA	RNA	protein	lipid	protein
The technique doesn't involve electrophoresis for the separation of biomolecules	Dot blotting	Southern blotting	Northern blotting	Western blotting	Dot blotting
Molecular Probe is a	Protein	Short piece of labeled DNA	Short piece of labeled protein	enzyme	Short piece of labeled DNA
The process somatic cell gene therapy involves	introduction of remedial gene into zygote	introduction of remedial gene into blood cells	introduction of remedial gene into sperm	introduction of remedial gene into egg	introduction of remedial gene into blood cells
Promoters used in bacterial expression vector are	lac	BAX	TIC	LOCK	lac
The vir gene responsible for topoisomerase activity	vir D	vir E	vir B	vir G	vir D