

Instruction hours/week: L:0 T:0 P:3 Marks: Internal: 40 External: 60 Total: 100**End Semester Exam: 3 Hours****Course Objectives**

- To know the sources of enzymes and study the extraction and partial purification of enzyme acid phosphatase
- To standardize the optimum pH, optimum substrate concentration required for the maximum activity of acid phosphatase
- To analyse the inhibition pattern by various competitive inhibitors for the enzyme acid phosphatase purified from germinated mung bean
- To assay the activity of Lactate dehydrogenase and glucose – 6 – phosphate dehydrogenase enzymes

Course outcome (CO's)

1. The students will be expertise in extraction of protein from germinated mung bean.
 2. Understand the purification process of enzyme.
 3. Analyze the optimum pH, substrate concentration for the maximum enzyme activity.
 4. Gain hands on training in assay of enzymes acid phosphatase, Lactate dehydrogenase and Glucose – 6- phosphate dehydrogenase.
-
1. Partial purification of acid phosphatase from germinating mung bean.
 2. Assay of enzyme activity and specific activity, e.g. acid phosphatase.
 3. Effect of pH on enzyme activity
 4. Determination of K_m and V_{max} using Lineweaver-Burk graph.
 5. Enzyme inhibition - calculation of K_i for competitive inhibition.
 6. Continuous assay of lactate dehydrogenase.
 7. Assay of glucose-6-phosphate dehydrogenase.

SUGGESTED READING

1. Nelson, D.L. and Cox, M.M., (2013). Lehninger: Principles of Biochemistry 6th ed., W.H. Freeman and Company (New York), ISBN:13: 978-1-4641-0962-1 / ISBN:10:1-4292-3414-8.
2. Donald, V., and Judith G.V., (2011). Biochemistry 4th ed., John Wiley & Sons Asia Pvt. Ltd. (New Jersey), ISBN:978-1180-25024.



KARPAGAM ACADEMY OF HIGHER EDUCATION

(Deemed to be University Established Under Section 3 of UGC Act 1956)

Coimbatore – 641 021.

LECTURE PLAN DEPARTMENT OF BIOCHEMISTRY

STAFF NAME: Ms.P.LOGANAYAKI

SUBJECT NAME: PRACTICAL - ENZYMES

SUB.CODE: 18BCU212

CLASS: I B.Sc.(BC)

SEMESTER: II

S.No	Duration of period	Topics covered	Support material
1	2	Partial purification of acid phosphatase from germinating mung bean	T1:16777- 16782
2	2	Assay of enzyme activity and specific activity, e.g. acid phosphatase.	J3:743- 756
3	1	Effect of pH on enzyme activity	
4	1	Determination of K_m and V_{max} using Lineweaver-Burk graph	
5	1	Enzyme inhibition - calculation of K_i for competitive inhibition	
6	1	Continuous assay of lactate dehydrogenase.	T1: 331- 334
7	1	Assay of glucose-6-phosphate dehydrogenase.	J2:313-315
8	1	Model practical Examination	-

REFERENCES:

T1:Dr.S.Rajan (2012) manual for medical laboratory technology- 1st Edition, Anjanna book house, Chennai.

J1: Surchandra et.al.,(2012), African journal of biotechnology:11(103) 16777- 16782

J2: Balinsky.D and Bernstern .R.E.1963: Biochemistry, Biophysicsaeta 67:313-315.

J3: A.K.M. Asaduxaman et al, (2011) Arch.bio Bdgrade 63(3).

SYLLABUS

1. Partial purification of acid phosphatase from germinating mung bean.
2. Assay of enzyme activity and specific activity, e.g. acid phosphatase.
3. Effect of pH on enzyme activity
4. Determination of K_m and V_{max} using Lineweaver-Burk graph.
5. Enzyme inhibition - calculation of K_i for competitive inhibition.
6. Continuous assay of lactate dehydrogenase.
7. Assay of glucose-6-phosphate dehydrogenase.

1. PARTIAL PURIFICATION OF ACID PHOSPHATASE FROM GERMINATED MUNG BEAN

1.A. AMMONIUM SULPHATE PRECIPITATION

AIM

To precipitate the protein content by ammonium sulphate precipitation method using various concentrations of ammonium sulphate.

THEORY

Precipitation is the method of concentrating proteins prior to analysis for a subsequent purification step. Protein precipitate is formed by the aggregation of protein molecules induced by changing of pH or ionic strength or by addition of organic solvents or other inert solutes or polymers. The solubility of a protein molecule in an aqueous solvent is determined by the distribution of charged hydrophilic and hydrophobic groups.

PRINCIPLE

Ammonium sulphate is particularly a useful salt in the fractional precipitation of proteins which precipitates the protein by changing its ionic strength of the medium as the ionic strength increase protein solubility.

This is referred to as 'salting in' however at a certain point the solubility starts to decrease, and this is known as 'salting out'.

Salting out depends on the hydrophobic nature of the surface of the proteins, hydrophobic groups predominate in the interior of the protein but some are located at the surface often in patches. When the protein is in solution, water molecules surround the protein forming hydrogen bonds with the protein. When salts are added to the system, water solubilises the ions and as the salt concentration increases, water is removed around the protein eventually exposed in the hydrophobic patches. Hydrophobic patches on one protein molecule can interact with those on another resulting in aggregation. Thus protein with large or more hydrophobic patches will aggregate and precipitate before with smaller or fewer hydrophobic patches

resulting in fractionation, hydrophobic proteins precipitate out first. But the hydrophilic proteins precipitate with much higher salt concentration. Advantages of ammonium sulphate are high solubility, low cost and easy availability. It is also non-toxic to most enzymes and it has stabilizing effect to all enzymes.

MATERIALS REQUIRED

1. Reagents

Extract preparation

Mung bean seeds were collected from local market and 10g of seeds are weighed and washed using distilled water and soaked with 20ml of distilled water in petridish and incubated at 20°C for 24 hours to induce germination. After 24 hours the seeds were crushed using mortar and pestle with appropriate solutions and made upto 100 ml with the same buffer solution and filtered. This filtrate is used as crude extract.

2. Ammonium sulphate

3. Citrate buffer

Solution A:

Dissolve 21.01g of citric acid in 1 litre of distilled water.

Solution B:

Dissolved 21.41g of sodium citrate in 1 litre of distilled water.

PROCEDURE

1. The prepared crude extract is transferred to the beaker placed in a tray.
2. Check the ammonium sulphate chart and weigh the appropriate amount of ammonium sulphate to get 30% solution fraction of protein.
3. Slowly add ammonium sulphate crystal with crude extract collected in beaker and dissolved it completely. Vigorous stirring will lead to denaturation of protein.
4. Keep the saturated solution undisturbed at 4°C for 30-60 minutes.
5. Then transferred the solution to centrifuge tubes and spin at 1000rpm for 20 minutes at 4°C.
6. Carefully transferred a supernatant to beaker placed in an ice tray.

7. Dissolved the pellet in 1ml of buffer.
8. Put the supernatant again and slowly add ammonium sulphate up to the final concentration of 60% mixing continuously until its dissolve completely
9. Repeat the steps 4,5,6 and 7
10. Again slowly add ammonium sulphate to the supernatant to a final concentrations of 90% repeat the steps 4,5,6 and 7
11. All the three pellets dissolved in minimal volume of extraction buffer and dialysed against the same buffer with constant stirring for overnight
12. It was used for acid phosphatase activity assay and the protein concentration is measured throughout the purification process by the method of process given by Lowry's method 1951 using BSA as standard protein.

1.b. ESTIMATION OF PROTEIN

Lowry *et al.*, 1957

AIM

To estimate the amount of protein present in the given sample.

PRINCIPLE

The blue color developed by the reduction of the phosphomolybdic phosphotungstic components in the Folin-Ciocalteu reagent by the amino acids tyrosine and tryptophan present in the protein plus the color developed by the biuret reaction of the protein with the alkaline cupric tartarate are measured at 660nm.

Reagents

1. 2% Sodium carbonate in 0.1N NaOH (Reagent A)
2. 0.5% Copper sulphate in 1% potassium sodium tartarate (Reagent B)
3. Alkaline copper reagent:

Mixed 50ml of A and 0.1ml of B prior to use.

4. Folin-Ciocalteu reagent:

Mixed 1 part of reagent with 2 parts of water.

5. Stock standard:

Weighed 50mg of bovine serum albumin and made up to 50ml in a standard flask with saline.

6. Working standard:

Diluted 10ml of the stock with 50ml of distilled water. 1.0ml of this solution contains 200µg of protein.

PROCEDURE

Pipetted out 0.2 to 1.0ml working standard solution, 0.1ml of the sample was taken. The volume in all the tubes was made up to 1.0ml with distilled water. Added 5.0ml of alkaline copper reagent to each tube. Mixed well and allowed to stand for 15 minutes. Then added 0.5ml of Folin-Ciocalteu reagent. Mixed well and incubated at room temperature for 30 minutes. A reagent blank was also prepared. After 30 minutes, the blue color developed was read at 660nm. A standard graph was drawn by plotting the concentration on x axis and the optical density on y axis. From this, the concentration of protein in the given sample solution was calculated.

RESULT

The amount of protein present in the given solution was found to be ----- g/dl

2. ASSAY OF ENZYME ACTIVITY AND SPECIFIC ACTIVITY OF ACID PHOSPHATASE

King, 1965

AIM

To estimate the amount of acid phosphatase present in the given sample.

PRINCIPLE

The method used was that of King and Armstrong in which disodium phenylphosphate is hydrolyzed with the liberation of phenol and inorganic phosphate. The liberated phenol is measured at 700nm with Folin-Ciocalteu reagent.

REAGENTS

1. Citrate buffer: 0.1M, pH 5.

A: Citric acid (21.01g in 100ml)

B: Sodium citrate (29.41g in 100ml)

2. Disodium phenyl phosphate, 100mmol/L:

Dissolved 2.18g of disodium phenylphosphate in distilled water and heated to boil, cooled and made to a litre. Added 1.0ml of chloroform and stored in the refrigerator.

3. Buffered substrate:

Prepared by mixing equal volume of the above two solutions. This has a pH of 5.0.

4. Folin-Ciocalteu reagent:

Prepared by mixing one volume of reagent and two volumes of water. Sodium carbonate solution, 15%: Dissolved 15g of anhydrous sodium carbonate in 100ml water.

5. Standard phenol solution, 1g/L:

Dissolved 1g pure crystalline phenol in 100mmol/L HCl and made to a litre with acid.

6. Working standard solution:

Diluted 10ml of stock standard to 100ml with water. This contains 100 μ g of phenol/ml.

PROCEDURE

Pipetted out 4.0ml of buffered substrate into a test tube and incubated at 37°C for 5 minutes. Added 0.2ml of sample and incubated further for exact 60 minutes. Removed and immediately added 1.8ml of diluted phenol reagent. At the same time, set up a control containing 4.0ml buffered substrate and 0.2ml of sample to which 1.8ml of phenol reagent was added immediately. Mixed and centrifuged. To 4.0ml of supernatant added 2.0ml of 15% sodium carbonate. Took 4.0 ml of working standard solution and for blank taken 3.2 ml water and 0.8ml of phenol reagent. Then added 2.0 ml of sodium carbonate. Incubated all the tubes at 37°C for 5 minutes. Read the color developed at 700nm.

A standard graph was drawn by plotting the concentration on x axis and the optical density on y axis. From this, the concentration of acid phosphatase in the given sample solution was calculated.

RESULT

The activity of acid phosphatase was found to be ----- μ moles of phenol liberated per litre.

3. EFFECT OF PH ON ENZYME ACTIVITY

AIM:

To find the effect of pH on acid phosphatase activity.

Principle:

This enzyme catalyses the hydrolysis of phosphate ester to phosphoric acid and alcohol. The amount of phosphoric acid produced during hydrolysis is the measure of enzyme activity. The liberated phosphoric acid containing the inorganic phosphorous is estimated by fiske and subbarow method.

Reagents

- Citrate buffer (0.2M)
- Sodium β glycerophosphate (substrate) (0.1M)
- 15% Trichloro acetic acid
- Ammonium molybdate solution
- ANSA reagent
- Magnesium acetate (0.2M)

PREPARATION OF BUFFER

pH	Solution A: x ml of 0.2 M citric acid (ml)	Solution B: y ml of 0.2 M sodium citrate	H ₂ O
4.6	12.75	12.25	25
5.0	10.25	14.75	25
5.6	6.85	18.15	25
6.0	4.75	20.25	25
6.6	2.5	22.5	25

Procedure:

Citrate buffer of various pH (4.6-6.4) is pipetted out into a clean dry test tubes. Then added 0.5ml of magnesium acetate solution and 1ml of the substrate to each followed by the addition of 2.0ml of the enzyme. Incubated at 37°C for 1 hour. At the end of incubation period added 1.0ml of 15% trichloroacetic acid to stop the reaction. Controls were also conducted along with these to which the enzyme was added at the end of incubation period.

Standard phosphate solution in the range 1 µg to 8 µg were pipetted out into clean dry test tubes. This serves as the standard to produce the standard graph.

All the test tubes are centrifuged and the supernatant of 1ml is transferred to other tubes. Added 1ml of molybdate solution followed by 0.4ml of ANSA and made up to 10ml with distilled water. The colour developed was read at 660nm after 20 minutes. The enzyme activity at various pH is found out from the standard graph.

Result:

The activity of the enzyme acid phosphatase was found to be maximum at pH ____ for an incubation period of 1 hour at 37°C.

EFFECT OF PH ON ACID PHOSPHATASE ACTIVITY

Ph	4.6	5.0	5.6	6.0	6.6
Volume of citrate buffer (ml)	5.0	5.0	5.0	5.0	5.0
Volume of magnesium acetate (ml)	0.5	0.5	0.5	0.5	0.5
Volume of substrate (ml)	1.0	1.0	1.0	1.0	1.0
Volume of enzyme (ml)	2.0	2.0	2.0	2.0	2.0
Incubate at 37°C for one hour					
Volume of 10% TCA	1.0	1.0	1.0	1.0	1.0
Add enzyme to the control and centrifuge at 5000 rpm for 10 minutes					
Volume of supernatant (ml)	1.0	1.0	1.0	1.0	1.0
Volume of ammonium molybdate (ml)	1.0	1.0	1.0	1.0	1.0
Volume of ANSA (ml)	0.4	0.4	0.4	0.4	0.4
Volume of water (ml)	7.6	7.6	7.6	7.6	7.6

Colorimeter reading	Control					
	Experiment					
	Difference					

4. DETERMINATION OF K_M AND V_{MAX} USING LINEWEAVER-BURK GRAPH.

Aim:

To find the effect of substrate concentration on acid phosphatase activity.

Principle:

This enzyme catalyses the hydrolysis of phosphate ester to phosphoric acid and alcohol. The amount of phosphoric acid produced during hydrolysis is the measure of enzyme activity. The liberated phosphoric acid containing the inorganic phosphorous is estimated by Fiske and Subbarow method.

Procedure:

Into clean dry test tubes pipetted out 0.5ml of magnesium acetate solution and varying volume of substrates (0.5, 1.0, 1.5, 2.0, 2.5 and 3.0ml) and made up to 7.0ml with 0.2M citrate buffer of pH 5.6 (test) followed by the addition of 1.0 ml of the enzyme. Incubated at 37 °C temperature for one hour, stop the reaction by the addition of 2.0ml of 10% TCA. Along with this control is also conducted to which the enzyme was added at the end of the incubation period. Centrifuged and transferred 1ml of the supernatant of each tubes to other tubes. Added 1ml of the ammonium molybdate solution followed by 0.4 ml of ANSA reagent. Made up the volume of each tube to 10 ml with distilled water. Read the colour developed at 660nm in a colorimeter after 20 minutes. A Michaelis Menton plot graph was drawn by plotting the substrate variation on x axis and colorimeter reading on y axis. From this the V_{max} , $1/2V_{max}$ and k_m values were calculated.

A Line Weaver burk plot was also prepared by plotting the reciprocal of the substrate concentration on x axis and the reciprocal of the optical density on y axis. The graph was plotted to get the value of K_m .

KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I BSC BC
COURSE CODE: 18BCU212

COURSE NAME: ENZYMES-PRACTICALS
BATCH-2018-2021

Result:

The value of K_m from Michaelis Menton plot = $1/2 V_{max}$ =

The value of K_m from Line Weaver burk plot = $1/2 V_{max}$ =

EFFECT OF SUBSTRATE CONCENTRATION ON ACID PHOSPHATASE ACTIVITY

S.NO	1	2	3	4	5	6	7
Volume of substrate (ml)	0.5	1.0	1.5	2.0	2.5	3.0	3.5
Volume of citrate buffer (ml)	6.5	6.0	5.5	5.0	4.5	4.0	3.5
Volume of enzyme (ml)	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Volume of magnesium acetate (ml)	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Incubate at 37°C for one hour							
Volume of 10% TCA	2.0	2.0	2.0	2.0	2.0	2.0	2.0
Add enzyme to the control and centrifuge at 5000 rpm for 10 minutes							
Volume of supernatant (ml)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Volume of ammonium molybdate (ml)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Volume of ANSA (ml)	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Volume of water (ml)	7.6	7.6	7.6	7.6	7.6	7.6	7.6
Colorimeter reading	Control						
	Experiment						
	Difference						

Calculation

Molecular weight of sodium β – glycerol phosphate = 315.4 g

31.54 g of sodium β – glycerol phosphate was dissolved in a litre of distilled water

The molarity of sodium β – glycerol phosphate = $\frac{31.54}{315.4} = 0.1\text{M}$

Dilution factor and substrate concentration in each tube

$$\frac{\text{molarity}}{\text{dilution factor}} = \text{substrate concentration}$$

0.5 ml of substrate was diluted to 10 ml = $\frac{10}{0.5} = 20$, $\frac{0.1}{20} = 0.5 \times 10^{-2} \text{ M}$

1.0 ml of substrate was diluted to 10 ml = $\frac{10}{1.0} = 10$, $\frac{0.1}{10} = 1.0 \times 10^{-2} \text{ M}$

1.5 ml of substrate was diluted to 10 ml = $\frac{10}{1.5} = 6.66$, $\frac{0.1}{6.66} = 1.5 \times 10^{-2} \text{ M}$

2.0 ml of substrate was diluted to 10 ml = $\frac{10}{2.0} = 5$, $\frac{0.1}{5} = 2.0 \times 10^{-2} \text{ M}$

2.5 ml of substrate was diluted to 10 ml = $\frac{10}{2.5} = 4$, $\frac{0.1}{4} = 2.5 \times 10^{-2} \text{ M}$

3.0 ml of substrate was diluted to 10 ml = $\frac{10}{3.0} = 3.33$, $\frac{0.1}{3.33} = 3.0 \times 10^{-2} \text{ M}$

3.5 ml of substrate was diluted to 10 ml = $\frac{10}{3.5} = 2.85$, $\frac{0.1}{2.85} = 3.5 \times 10^{-2} \text{ M}$

Line Weaver Burk plot

[S] X 10^{-2}m	V	1/[S] X 10^{-2}m	1/[V]
0.5		2.0	
1.0		1.0	
1.5		0.6	
2.0		0.5	
2.5		0.4	

3.0		0.3	
3.5		0.2	

5. ENZYME INHIBITION - CALCULATION OF K_i FOR COMPETITIVE INHIBITION

AIM

To study the effect of metal ions in inhibiting enzyme activity.

PROCEDURE

The mixture, consisting of 100 μ l of the 0.1M cation solution under test, 600 μ l of 1M acetate buffer pH 5.5 and 100 μ l of enzyme solution, was pre-incubated for 10 min at 37°C. After preincubation, 200 μ l of 20 mM p-nitrophenyl phosphate was added to determine the activity as usual. Simultaneously control and blank experiments were run in which the cation and enzyme solutions were replaced by water, respectively, in the pre-incubation mixtures. Similarly, the effect of some compounds reacting with SH-groups of the enzyme on the enzyme activity at pH values varying from 3 to 9 was determined as described above. Kinetic parameters: The K_m , V_{max} and K_i values were determined using p-nitrophenyl phosphate as the substrate in concentrations of 0.06 - 4 mM in the absence or presence of two or three fixed concentrations of inhibitors. These kinetic parameters were determined from Line-weaver-Burk plots. Straight lines were drawn by applying least square rule. The pH dependence studies of K_m , V_{max} and specificity constants were carried out.

6. CONTINUOUS ASSAY OF LACTATE DEHYDROGENASE.

AIM

To assay the activity of lactate dehydrogenase in the given sample.

PRINCIPLE

The lactate is acted upon by lactate dehydrogenase to form pyruvate in the presence of NAD. The pyruvate forms pyruvate phenyl hydrazone with 2,4 dinitrophenyl hydrazine. The color developed is read in a spectrophotometer at 440nm.

REAGENTS

1. Glycine buffer, 0.1M, pH 10: 7.505 g of glycine and 5.85 g of sodium chloride were dissolved in 1 litre of water.
2. Buffered substrate: 125 ml of glycine buffer and 75 ml of 0.1 N NaOH were added to 4 g of lithium lactate and mixed well.
3. Nicotinamide Adenine Dinucleotide: 10 mg of NAD was dissolved in 2.0 ml of water.
4. 2, 4 Dinitrophenyl hydrazine: 20 mg of DNPH was dissolved in 100 ml of 1 N HCl.
5. 0.4 N NaOH.
6. Standard, 1 μ mol/ml: 11 mg of sodium pyruvate was dissolved in 100 ml of buffered substrate (1 μ mole of pyruvate/ml).
7. NADH solution, 1 μ mol/ml: 8.5 mg/10 ml buffered substrate.

PROCEDURE

Placed 1.0 ml of buffered substrate and 0.1 ml of sample into each of two tubes. Added 0.2 ml of water to the blank. Then to the test added 0.2 ml of NAD. Mixed and incubated at 37°C for 15 min. Exactly after 15 min, 1.0 ml of dinitrophenyl hydrazine was added to each (test

and control). Left for further 15 min. Then added 10 ml of 0.4N sodium hydroxide and the color developed was read immediately at 440 nm. A standard curve with sodium pyruvate solution with the concentration range 0.02 - 0.10 μ mole was taken.

LDH activity in serum was expressed as μ moles of pyruvate liberated/L and in tissue homogenate as nmoles of pyruvate liberated/min/mg protein.

7. ASSAY OF GLUCOSE-6-PHOSPHATE DEHYDROGENASE

AIM

To assay the activity of glucose-6-phosphate dehydrogenase in the given sample.

PRINCIPLE

Glucose-6-phosphate dehydrogenase is assayed by measuring the increase in absorbance, which occurs at 340 nm when NADP reduces to glucose-6-phosphate to NADP in the reaction catalyzed by glucose-6-phosphate dehydrogenase.

REAGENTS

- ❖ 0.1M Tris-HCl buffer, pH 8.2
 - A: 0.1M solutions of Tris (12.1g/1000 ml water)
 - B: 0.1 M HCl
- ❖ Mixed 50 ml of solution A and 21.9 ml of B and diluted to a total of 200 ml
- ❖ 0.2 mM NADP
- ❖ 0.1M magnesium chloride
- ❖ 6 mM glucose-6-phosphate

PROCEDURE

0.4 ml of Tris-HCl buffer, 0.2 ml of NADP, 0.2 ml of magnesium chloride, 1.0 ml water and 0.2 ml of enzyme were taken in a cuvette. The reaction was started by the addition of 0.2 ml of glucose-6-phosphate and the increase in OD was measured at 340 nm. The activity was expressed in terms of units/mg protein, in which one unit is equal to the amount of enzyme that brought about a change in OD of 0.01/min.