Semester – IV



# KARPAGAM ACADEMY OF HIGHER EDUCATION

(Deemed to be University Established Under Section 3 of UGC Act 1956)

**Coimbatore – 641 021.** 

**SYLLABUS** 

		L	T	P	C	•
17MMU402	PDE AND SYSTEMS OF ODE	6	2	0	6	

**Scope:** This Course enables the students to learn the various methods of solving Differential equations which is very much used in the field of Engineering.

**Objectives:** To enable the students to learn and gain knowledge about forming the differential equations method of separation of Variables, Initial Boundary Value Problems and method of successive approximations

## UNIT I

Partial Differential Equations – Basic concepts and Definitions -Mathematical Problems. First Order Equations: Classification - Construction and Geometrical Interpretation- Method of characteristics for obtaining General Solution of Quasi Linear Equations- Canonical Forms of First-order Linear Equations.

## UNIT II

Method of Separation of Variables for solving first order partial differential equations. Derivation of Heat equation -Wave equation and Laplace equation. - Classification of second order - linear equations as hyperbolic, parabolic or elliptic.

## UNIT III

Reduction of second order Linear Equations to canonical forms- The Cauchy problem- The Cauchy-Kowaleewskaya theorem -Cauchy problem of an infinite string - Initial Boundary Value Problems -Semi-Infinite String with a fixed end - Semi-Infinite String with a Free end-Equations with non-homogeneous boundary conditions -Non- Homogeneous Wave Equation. **UNIT I V** 

Method of separation of variables - Solving the Vibrating String - Problems- Solving the Heat Conduction problem - Systems of linear differential equations - Types of linear systems differential operators - an operator method for linear systems with constant coefficients.

## UNIT V

Basic Theory of linear systems in normal form : Homogeneous linear systems with constant coefficients -Two Equations in two unknown functions -The method of successive

approximations -The Euler method-The modified – Euler method -The Runge-Kutta method. **SUGGESTED READINGS** 

# TEXT BOOK

1. Tyn Myint-U and Lokenath Debnath., (2006). Linear Partial Differential Equations for Scientists and Engineers, 4th edition, Springer, Indian reprint.

# REFERENCES

- 1. Ross S.L., (2004). Differential equations, Third Edition, John Wiley and Sons, India.
- 2. Martha L Abell., James P Braselton, (2004). Differential equations with MATHEMATICA, Third Edition Elsevier Academic Press.

CLASS: II B.Sc COURSE CODE: 17MMU402

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: I BATCH-2017-2020

UNIT I Partial Differential Equations SYLLABUS

Partial Differential Equations – Basic concepts and Definitions -Mathematical Problems. First Order Equations: Classification - Construction and Geometrical Interpretation- Method of characteristics for obtaining General Solution of Quasi Linear Equations- Canonical Forms of First-order Linear Equations.

# **Basic Concepts and Definitions**

A differential equation that contains, in addition to the dependent variable and the independent variables, one or more partial derivatives of the dependent variable is called a *partial differential equation*. In general, it may be written in the form

$$f(x, y, \dots, u, u_x, u_y, \dots, u_{xx}, u_{xy}, \dots) = 0, \qquad (1.2.1)$$

involving several independent variables  $x, y, \ldots$ , an unknown function u of these variables, and the partial derivatives  $u_x, u_y, \ldots, u_{xx}, u_{xy}, \ldots$ , of the function. Subscripts on dependent variables denote differentiations, e.g.,

 $u_x = \partial u / \partial x, \qquad u_{xy} = \partial^2 / \partial y \, \partial x.$ 

Here equation (1.2.1) is considered in a suitable domain D of the *n*dimensional space  $\mathbb{R}^n$  in the independent variables  $x, y, \ldots$ . We seek functions  $u = u(x, y, \ldots)$  which satisfy equation (1.2.1) identically in D. Such functions, if they exist, are called *solutions* of equation (1.2.1). From these many possible solutions we attempt to select a particular one by introducing suitable additional conditions.

For instance,

$$uu_{xy} + u_x = y,$$
  

$$u_{xx} + 2yu_{xy} + 3xu_{yy} = 4\sin x,$$
  

$$(u_x)^2 + (u_y)^2 = 1,$$
  

$$u_{xx} - u_{yy} = 0,$$
  
(1.2.2)

### CLASS: II B.Sc COURSE CODE: 17MMU402

### COURSE NAME: PDE AND SYSTEM OF ODE UNIT: I BATCH-2017-2020

are partial differential equations. The functions

$$u(x, y) = (x + y)^{3},$$
  
 $u(x, y) = \sin(x - y),$ 

are solutions of the last equation of (1.2.2), as can easily be verified.

The *order* of a partial differential equation is the order of the highestordered partial derivative appearing in the equation. For example

 $u_{xx} + 2xu_{xy} + u_{yy} = e^y$ 

is a second-order partial differential equation, and

$$u_{xxy} + xu_{yy} + 8u = 7y$$

is a third-order partial differential equation.

A partial differential equation is said to be *linear* if it is linear in the unknown function and all its derivatives with coefficients depending only on the independent variables; it is said to be *quasi-linear* if it is linear in the highest-ordered derivative of the unknown function. For example, the equation

$$yu_{xx} + 2xyu_{yy} + u = 1$$

is a second-order linear partial differential equation, whereas

$$u_x u_{xx} + x u u_y = \sin y$$

is a second-order quasi-linear partial differential equation. The equation which is not linear is called a *nonlinear* equation.

We shall be primarily concerned with linear second-order partial differential equations, which frequently arise in problems of mathematical physics. The most general second-order linear partial differential equation in n independent variables has the form

$$\sum_{i,j=1}^{n} A_{ij} u_{x_i x_j} + \sum_{i=1}^{n} B_i u_{x_i} + F u = G, \qquad (1.2.3)$$

CLASS: II B.Sc	COURSE NAME: PDE AN	<b>ND SYSTEM OF ODE</b>
COURSE CODE: 17MMU402	UNIT: I	BATCH-2017-2020

where we assume without loss of generality that  $A_{ij} = A_{ji}$ . We also assume that  $B_i$ , F, and G are functions of the n independent variables  $x_i$ .

If G is identically zero, the equation is said to be *homogeneous*; otherwise it is *nonhomogeneous*.

The general solution of a linear ordinary differential equation of nth order is a family of functions depending on n independent arbitrary constants. In the case of partial differential equations, the general solution depends on arbitrary functions rather than on arbitrary constants. To illustrate this, consider the equation

 $u_{xy} = 0.$ 

If we integrate this equation with respect to y, we obtain

$$u_x\left(x,y\right) = f\left(x\right).$$

A second integration with respect to x yields

$$u(x,y) = g(x) + h(y),$$

where g(x) and h(y) are arbitrary functions.

Suppose u is a function of three variables, x, y, and z. Then, for the equation

$$u_{yy} = 2,$$

one finds the general solution

$$u(x, y, z) = y^{2} + yf(x, z) + g(x, z),$$

where f and g are arbitrary functions of two variables x and z.

# Mathematical Problems

A problem consists of finding an unknown function of a partial differential equation satisfying appropriate supplementary conditions. These conditions may be *initial conditions* (I.C.) and/or *boundary conditions* (B.C.). For example, the partial differential equation (PDE)

CLASS: II B.Sc	<b>COURSE NAME: PDE AND</b>	SYSTEM OF ODE
COURSE CODE: 17MMU402	UNIT: I	BATCH-2017-2020

 $u_t - u_{xx} = 0, \qquad 0 < x < l, \quad t > 0,$ with *I.C.*  $u(x,0) = \sin x, \quad 0 \le x \le l, \quad t > 0,$ *B.C.*  $u(0,t) = 0, \qquad t \ge 0,$ *B.C.*  $u(l,t) = 0, \qquad t \ge 0,$ 

constitutes a problem which consists of a partial differential equation and three supplementary conditions. The equation describes the heat conduction in a rod of length l. The last two conditions are called the *boundary conditions* which describe the function at two prescribed boundary points. The first condition is known as the *initial condition* which prescribes the unknown function u(x,t) throughout the given region at some initial time t, in this case t = 0. This problem is known as the *initial boundary-value problem*. Mathematically speaking, the time and the space coordinates are regarded as independent variables. In this respect, the initial condition is merely a point prescribed on the t-axis and the boundary conditions are prescribed, in this case, as two points on the x-axis. Initial conditions are usually prescribed at a certain time  $t = t_0$  or t = 0, but it is not customary to consider the other end point of a given time interval.

In considering the problem of unbounded domain, the solution can be determined uniquely by prescribing initial conditions only. The corresponding problem is called the *initial-value problem* or the *Cauchy problem*. The

A mathematical problem is said to be *well-posed* if it satisfies the following requirements:

- 1. Existence: There is at least one solution.
- 2. Uniqueness: There is at most one solution.
- 3. Continuity: The solution depends continuously on the data.

# Classification of First-Order Equations

The most general, first-order, partial differential equation in two independent variables x and y is of the form

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

$$F(x, y, u, u_x, u_y) = 0,$$
  $(x, y) \in D \subset \mathbb{R}^2,$  (2.2.1)

where F is a given function of its arguments, and u = u(x, y) is an unknown function of the independent variables x and y which lie in some given domain D in  $R^2$ ,  $u_x = \frac{\partial u}{\partial x}$  and  $u_y = \frac{\partial u}{\partial y}$ . Equation (2.2.1) is often written in terms of standard notation  $p = u_x$  and  $q = u_y$  so that (2.2.1) takes the form

$$F(x, y, u, p, q) = 0.$$
 (2.2.2)

Similarly, the most general, first-order, partial differential equation in three independent variables x, y, z can be written as

$$F(x, y, z, u, u_x, u_y, u_z) = 0.$$
(2.2.3)

Equation (2.2.1) or (2.2.2) is called a *quasi-linear partial differential* equation if it is linear in first-partial derivatives of the unknown function u(x, y). So, the most general quasi-linear equation must be of the form

$$a(x, y, u) u_x + b(x, y, u) u_y = c(x, y, u), \qquad (2.2.4)$$

where its coefficients a, b, and c are functions of x, y, and u.

The following are examples of quasi-linear equations:

$$x(y^{2}+u)u_{x} - y(x^{2}+u)u_{y} = (x^{2}-y^{2})u, \qquad (2.2.5)$$

$$uu_x + u_t + nu^2 = 0, (2.2.6)$$

$$(y^2 - u^2) u_x - xy u_y = xu. (2.2.7)$$

Equation (2.2.4) is called a *semilinear partial differential equation* if its coefficients a and b are independent of u, and hence, the semilinear equation can be expressed in the form

$$a(x, y) u_x + b(x, y) u_y = c(x, y, u).$$
(2.2.8)

CLASS: II B.Sc	<b>COURSE NAME: I</b>	PDE AND SYSTEM OF ODE
COURSE CODE: 17MMU402	UNIT: I	BATCH-2017-2020

Examples of semilinear equations are

$$xu_x + yu_y = u^2 + x^2, (2.2.9)$$

$$(x+1)^2 u_x + (y-1)^2 u_y = (x+y) u^2, \qquad (2.2.10)$$

$$u_t + au_x + u^2 = 0, (2.2.11)$$

where a is a constant.

Equation (2.2.1) is said to be *linear* if F is linear in each of the variables  $u, u_x$ , and  $u_y$ , and the coefficients of these variables are functions only of the independent variables x and y. The most general, first-order, *linear* partial differential equation has the form

$$a(x, y) u_x + b(x, y) u_y + c(x, y) u = d(x, y), \qquad (2.2.12)$$

where the coefficients a, b, and c, in general, are functions of x and y and d(x, y) is a given function. Unless stated otherwise, these functions are assumed to be continuously differentiable. Equations of the form (2.2.12) are called *homogeneous* if  $d(x, y) \equiv 0$  or *nonhomogeneous* if  $d(x, y) \neq 0$ .

Obviously, linear equations are a special kind of the quasi-linear equation (2.2.4) if a, b are independent of u and c is a linear function in u. Similarly, semilinear equation (2.2.8) reduces to a linear equation if c is linear in u.

Examples of linear equations are

$$xu_x + yu_y - nu = 0, (2.2.13)$$

$$nu_x + (x+y)u_y - u = e^x, \qquad (2.2.14)$$

$$yu_x + xu_y = xy, \tag{2.2.15}$$

$$(y-z)u_x + (z-x)u_y + (x-y)u_z = 0. (2.2.16)$$

An equation which is *not* linear is often called a *nonlinear equation*. So, first-order equations are often classified as linear and nonlinear.

CLASS: II B.Sc	COURSE NAME: 1	PDE AND SYSTEM OF ODE
COURSE CODE: 17MMU402	UNIT: I	BATCH-2017-2020

# Construction of a First-Order Equation

We consider a system of geometrical surfaces described by the equation

$$f(x, y, z, a, b) = 0, (2.3.1)$$

where a and b are arbitrary parameters. We differentiate (2.3.1) with respect to x and y to obtain

$$f_x + p f_z = 0, \qquad f_y + q f_z = 0,$$
 (2.3.2)

where  $p = \frac{\partial z}{\partial x}$  and  $q = \frac{\partial z}{\partial y}$ .

The set of three equations (2.3.1) and (2.3.2) involves two arbitrary parameters a and b. In general, these two parameters can be eliminated from this set to obtain a first-order equation of the form

F(x, y, z, p, q) = 0.(2.3.3)

Thus the system of surfaces (2.3.1) gives rise to a first-order partial differential equation (2.3.3). In other words, an equation of the form (2.3.1)containing two arbitrary parameters is called a *complete solution* or a *complete integral* of equation (2.3.3). Its role is somewhat similar to that of a general solution for the case of an ordinary differential equation.

On the other hand, any relationship of the form

$$f(\phi, \psi) = 0,$$
 (2.3.4)

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

First, we prescribe the second parameter b as an arbitrary function of the first parameter a in the complete solution (2.3.1) of (2.3.3), that is, b = b(a). We then consider the envelope of the one-parameter family of solutions so defined. This envelope is represented by the two simultaneous equations

$$f(x, y, z, a, b(a)) = 0,$$
 (2.3.5)

$$f_a(x, y, z, a, b(a)) + f_b(x, y, z, b(a)) b'(a) = 0, \qquad (2.3.6)$$

where the second equation (2.3.6) is obtained from the first equation (2.3.5) by partial differentiation with respect to a. In principle, equation (2.3.5) can be solved for a = a(x, y, z) as a function of x, y, and z. We substitute this result back in (2.3.5) to obtain

$$f\{x, y, z, a(x, y, z), b(a(x, y, z))\} = 0,$$
(2.3.7)

where b is an arbitrary function. Indeed, the two equations (2.3.5) and (2.3.6) together define the general solution of (2.3.3). When a definite b(a) is prescribed, we obtain a *particular solution* from the general solution. Since the general solution depends on an arbitrary function, there are infinitely many solutions. In practice, only one solution satisfying prescribed conditions is required for a physical problem. Such a solution may be called a *particular solution*.

# Geometrical Interpretation

To investigate the geometrical content of a first-order, partial differential equation, we begin with a general, quasi-linear equation

$$a(x, y, u) u_x + b(x, y, u) u_y - c(x, y, u) = 0.$$
(2.4.1)

We assume that the possible solution of (2.4.1) in the form u = u(x, y)or in an implicit form

$$f(x, y, u) \equiv u(x, y) - u = 0 \tag{2.4.2}$$

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

represents a possible solution surface in (x, y, u) space. This is often called an *integral surface* of the equation (2.4.1). At any point (x, y, u) on the solution surface, the gradient vector  $\nabla f = (f_x, f_y, f_u) = (u_x, u_y, -1)$  is normal to the solution surface. Clearly, equation (2.4.1) can be written as the dot product of two vectors

$$a u_x + b u_y - c = (a, b, c) \cdot (u_x, u_y - 1) = 0.$$
 (2.4.3)

This clearly shows that the vector (a, b, c) must be a tangent vector of the integral surface (2.4.2) at the point (x, y, u), and hence, it determines a direction field called the the *characteristic direction* or *Monge axis*. This direction is of fundamental importance in determining a solution of equation (2.4.1). To summarize, we have shown that f(x, y, u) = u(x, y) - u = 0, as a surface in the (x, y, u)-space, is a solution of (2.4.1) if and only if the direction vector field (a, b, c) lies in the tangent plane of the integral surface f(x, y, u) = 0 at each point (x, y, u), where  $\nabla f \neq 0$ , as shown in Figure 2.4.1.

A curve in (x, y, u)-space, whose tangent at every point coincides with the characteristic direction field (a, b, c), is called a *characteristic curve*. If the parametric equations of this characteristic curve are

$$x = x(t), \quad y = y(t), \quad u = u(t),$$
 (2.4.4)

then the tangent vector to this curve is  $\left(\frac{dx}{dt}, \frac{dy}{dt}, \frac{du}{dt}\right)$  which must be equal to (a, b, c). Therefore, the system of ordinary differential equations of the characteristic curve is given by

$$\frac{dx}{dt} = a\left(x, y, u\right), \quad \frac{dy}{dt} = b\left(x, y, u\right), \quad \frac{du}{dt} = c\left(x, y, u\right). \quad (2.4.5)$$

These are called the *characteristic equations* of the quasi-linear equation (2.4.1).

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

# Method of Characteristics and General Solutions

Theorem 2.5.1. The general solution of a first-order, quasi-linear partial differential equation

$$a(x, y, u) u_x + b(x, y, u) u_y = c(x, y, u)$$
(2.5.1)

is

$$f(\phi,\psi) = 0,$$
 (2.5.2)

where f is an arbitrary function of  $\phi(x, y, u)$  and  $\psi(x, y, u)$ , and  $\phi = \text{constant} = c_1$  and  $\psi = \text{constant} = c_2$  are solution curves of the characteristic equations

$$\frac{dx}{a} = \frac{dy}{b} = \frac{du}{c}.$$
(2.5.3)

The solution curves defined by  $\phi(x, y, u) = c_1$  and  $\psi(x, y, u) = c_2$  are called the families of *characteristic curves* of equation (2.5.1).

*Proof.* Since  $\phi(x, y, u) = c_1$  and  $\psi(x, y, u) = c_2$  satisfy equations (2.5.3), these equations must be compatible with the equation

$$d\phi = \phi_x dx + \phi_y dy + \phi_u du = 0. \tag{2.5.4}$$

This is equivalent to the equation

$$a\,\phi_x + b\,\phi_y + c\,\phi_u = 0. \tag{2.5.5}$$

Similarly, equation (2.5.3) is also compatible with

$$a\,\psi_x + b\,\psi_y + c\,\psi_u = 0. \tag{2.5.6}$$

We now solve (2.5.5), (2.5.6) for a, b, and c to obtain

$$\frac{a}{\frac{\partial(\phi,\psi)}{\partial(y,u)}} = \frac{b}{\frac{\partial(\phi,\psi)}{\partial(u,x)}} = \frac{c}{\frac{\partial(\phi,\psi)}{\partial(x,y)}}.$$
(2.5.7)

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

It has been shown earlier that  $f(\phi, \psi) = 0$  satisfies an equation similar to (2.3.14), that is,

$$p \frac{\partial(\phi, \psi)}{\partial(y, u)} + q \frac{\partial(\phi, \psi)}{\partial(u, x)} = \frac{\partial(\phi, \psi)}{\partial(x, y)}.$$
(2.5.8)

Substituting, (2.5.7) in (2.5.8), we find that  $f(\phi, \psi) = 0$  is a solution of (2.5.1). This completes the proof.

**Theorem 2.5.2.** (*The Cauchy Problem for a First-Order Partial Differential Equation*). Suppose that C is a given curve in the (x, y)-plane with its parametric equations

$$x = x_0(t), \quad y = y_0(t),$$
 (2.5.9)

where t belongs to an interval  $I \subset R$ , and the derivatives  $x'_0(t)$  and  $y'_0(t)$  are piecewise continuous functions, such that  $(x'_0)^2 + (y'_0)^2 \neq 0$ . Also, suppose that  $u = u_0(t)$  is a given function on the curve C. Then, there exists a solution u = u(x, y) of the equation

$$F(x, y, u, u_x, u_y) = 0$$
 (2.5.10)

in a domain D of  $\mathbb{R}^2$  containing the curve C for all  $t \in I$ , and the solution u(x, y) satisfies the given initial data, that is,

$$u(x_0(t), y_0(t)) = u_0(t)$$
 (2.5.11)

for all values of  $t \in I$ .

Theorem 2.5.3. (The Cauchy Problem for a Quasi-linear Equation). Suppose that  $x_0(t)$ ,  $y_0(t)$ , and  $u_0(t)$  are continuously differentiable functions of t in a closed interval,  $0 \le t \le 1$ , and that a, b, and c are functions of x, y, and u with continuous first-order partial derivatives with respect to their arguments in some domain D of (x, y, u)-space containing the initial curve

$$\Gamma: x = x_0(t), \quad y = y_0(t), \quad u = u_0(t), \quad (2.5.12)$$

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

where  $0 \le t \le 1$ , and satisfying the condition

$$y'_{0}(t) a(x_{0}(t), y_{0}(t), u_{0}(t)) - x'_{0}(t) b(x_{0}(t), y_{0}(t), u_{0}(t)) \neq 0.$$
 (2.5.13)

Then there exists a unique solution u = u(x, y) of the quasi-linear equation (2.5.1) in the neighborhood of C:  $x = x_0(t)$ ,  $y = y_0(t)$ , and the solution satisfies the initial condition

$$u_0(t) = u(x_0(t), y_0(t)), \text{ for } 0 \le t \le 1.$$
 (2.5.14)

# Canonical Forms of First-Order Linear Equations

It is often convenient to transform the more general first-order linear partial differential equation (2.2.12)

$$a(x, y) u_x + b(x, y) u_y + c(x, y) u = d(x, y),$$
 (2.6.1)

into a *canonical* (or *standard*) form which can be easily integrated to find the general solution of (2.6.1). We use the characteristics of this equation (2.6.1) to introduce the new transformation by equations

$$\xi = \xi (x, y), \qquad \eta = \eta (x, y), \qquad (2.6.2)$$

where  $\xi$  and  $\eta$  are once continuously differentiable and their Jacobian  $J(x, y) \equiv \xi_x \eta_y - \xi_y \eta_x$  is nonzero in a domain of interest so that x and y can be determined uniquely from the system of equations (2.6.2). Thus, by chain rule,

$$u_x = u_\xi \xi_x + u_\eta \eta_x, \qquad u_y = u_\xi \xi_y + u_\eta \eta_y, \qquad (2.6.3)$$

we substitute these partial derivatives (2.6.3) into (2.6.1) to obtain the equation

$$A \, u_{\xi} + B \, u_{\eta} + cu = d, \tag{2.6.4}$$

where

$$A = u\xi_x + b\xi_y, \qquad B = a\eta_x + b\eta_y. \tag{2.6.5}$$

From (2.6.5) we see that B = 0 if  $\eta$  is a solution of the first-order equation This equation has infinitely many solutions.

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IBATCH-2017-2020

 $\ln u\left(\xi,\eta\right) = -\eta + \ln f\left(\xi\right),$ 

where  $f(\xi)$  is an arbitrary function of  $\xi$  only. Equivalently,

$$u\left(\xi,\eta\right) = f\left(\xi\right)e^{-\eta}.$$

In terms of the original variables x and y, the general solution of equation (2.6.8) is

$$u(x,y) = f(x+y)e^{-y}, \qquad (2.6.10)$$

where f is an arbitrary function.

Example 2.6.1. Reduce each of the following equations

$$u_x - u_y = u,$$
 (2.6.8)

$$yu_x + u_y = x, \tag{2.6.9}$$

to canonical form, and obtain the general solution.

In (2.6.8), a = 1, b = -1, c = -1 and d = 0. The characteristic equations are

$$\frac{dx}{1} = \frac{dy}{-1} = \frac{du}{u}.$$

The characteristic curves are  $\xi = x + y = c_1$ , and we choose  $\eta = y = c_2$ where  $c_1$  and  $c_2$  are constants. Consequently,  $u_x = u_{\xi}$  and  $u_y = u_{\xi} + u_{\eta}$ , and hence, equation (2.6.8) becomes

$$u_{\eta} = u.$$

Integrating this equation gives

### CLASS: II B.Sc COURSE CODE: 17MMU402

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: II BATCH-2017-2020

## UNIT II SYLLABUS

Method of Separation of Variables for solving first order partial differential equations.Derivation of Heat equation -Wave equation and Laplace equation. -Classification of second order - linear equations as hyperbolic, parabolic or elliptic.

# Method of Separation of Variables

Example 2.7.1. Solve the initial-value problem

$$u_x + 2u_y = 0,$$
  $u(0, y) = 4e^{-2y}.$  (2.7.1ab)

We seek a separable solution  $u(x, y) = X(x) Y(y) \neq 0$  and substitute into the equation to obtain

$$X'(x) Y(y) + 2X(x) Y'(y) = 0.$$

This can also be expressed in the form

$$\frac{X'(x)}{2X(x)} = -\frac{Y'(y)}{Y(y)}.$$
(2.7.2)

Since the left-hand side of this equation is a function of x only and the right-hand is a function of y only, it follows that (2.7.2) can be true if both sides are equal to the same constant value  $\lambda$  which is called an arbitrary separation constant. Consequently, (2.7.2) gives two ordinary differential equations

$$X'(x) - 2\lambda X(x) = 0, \qquad Y'(y) + \lambda Y(y) = 0.$$
 (2.7.3)

These equations have solutions given, respectively, by

$$X(x) = A e^{2\lambda x}$$
 and  $Y(y) = B e^{-\lambda y}$ , (2.7.4)

where  ${\cal A}$  and  ${\cal B}$  are arbitrary integrating constants.

Consequently, the general solution is given by

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIBATCH-2017-2020

$$u(x,y) = AB \exp(2\lambda x - \lambda y) = C \exp(2\lambda x - \lambda y), \qquad (2.7.5)$$

where C = AB is an arbitrary constant.

Using the condition (2.7.1b), we find

$$4 e^{-2y} = u(0, y) = C e^{-\lambda y},$$

and hence, we deduce that C=4 and  $\lambda=2.$  Therefore, the final solution is

$$u(x,y) = 4\exp(4x - 2y).$$
(2.7.6)

# Classical Equations

The three basic types of second-order partial differential equations are: (a) The wave equation

$$u_{tt} - c^2 \left( u_{xx} + u_{yy} + u_{zz} \right) = 0. \tag{3.1.1}$$

(b) The heat equation

$$u_t - k (u_{xx} + u_{yy} + u_{zz}) = 0.$$
 (3.1.2)

(c) The Laplace equation

$$u_{xx} + u_{yy} + u_{zz} = 0. ag{3.1.3}$$

# Classification of Second-Order Linear Equations

# (A) Hyperbolic Type

If  $B^2 - 4AC > 0$ , then integration of equations (4.2.5) and (4.2.6) yield two real and distinct families of characteristics. Equation (4.1.11) reduces to

$$u_{\xi\eta} = H_1,$$
 (4.2.7)

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIBATCH-2017-2020

where  $H_1 = H^*/B^*$ . It can be easily shown that  $B^* \neq 0$ . This form is called the first canonical form of the hyperbolic equation.

Now if new independent variables

$$\alpha = \xi + \eta, \qquad \beta = \xi - \eta, \qquad (4.2.8)$$

are introduced, then equation (4.2.7) is transformed into

$$u_{\alpha\alpha} - u_{\beta\beta} = H_2(\alpha, \beta, u, u_\alpha, u_\beta).$$
 (4.2.9)

This form is called the second canonical form of the hyperbolic equation.

#### (B) Parabolic Type

In this case, we have  $B^2 - 4AC = 0$ , and equations (4.2.5) and (4.2.6) coincide. Thus, there exists one real family of characteristics, and we obtain only a single integral  $\xi = \text{constant}$  (or  $\eta = \text{constant}$ ).

Since  $B^2 = 4AC$  and  $A^* = 0$ , we find that

$$A^* = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = \left(\sqrt{A}\,\xi_x + \sqrt{C}\,\xi_y\right)^2 = 0.$$

From this it follows that

$$A^* = 2A\xi_x\eta_x + B\left(\xi_x\eta_y + \xi_y\eta_x\right) + 2C\xi_y\eta_y$$
  
=  $2\left(\sqrt{A}\xi_x + \sqrt{C}\xi_y\right)\left(\sqrt{A}\eta_x + \sqrt{C}\eta_y\right) = 0,$ 

for arbitrary values of  $\eta(x, y)$  which is functionally independent of  $\xi(x, y)$ ; for instance, if  $\eta = y$ , the Jacobian does not vanish in the domain of parabolicity.

Division of equation (4.1.11) by  $C^*$  yields

$$u_{\eta\eta} = H_3(\xi, \eta, u, u_{\xi}, u_{\eta}), \quad C^* \neq 0.$$
 (4.2.10)

This is called the *canonical form of the parabolic equation*.

Equation (4.1.11) may also assume the form

$$u_{\xi\xi} = H_3^* \left(\xi, \eta, u, u_{\xi}, u_{\eta}\right), \tag{4.2.11}$$

if we choose  $\eta = \text{constant}$  as the integral of equation (4.2.5).

## CLASS: II B.Sc COURSE CODE: 17MMU402

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: II BATCH-2017-2020

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIBATCH-2017-2020

## (C) Elliptic Type

For an equation of elliptic type, we have  $B^2 - 4AC < 0$ . Consequently, the quadratic equation (4.2.4) has no real solutions, but it has two complex conjugate solutions which are continuous complex-valued functions of the real variables x and y. Thus, in this case, there are no real characteristic curves. However, if the coefficients A, B, and C are analytic functions of x and y, then one can consider equation (4.2.4) for complex x and y. A function of two real variables x and y is said to be analytic in a certain domain if in some neighborhood of every point  $(x_0, y_0)$  of this domain, the

function can be represented as a Taylor series in the variables  $(x - x_0)$  and  $(y - y_0)$ .

Since  $\xi$  and  $\eta$  are complex, we introduce new real variables

$$\alpha = \frac{1}{2} \left( \xi + \eta \right), \qquad \beta = \frac{1}{2i} \left( \xi - \eta \right), \qquad (4.2.12)$$

so that

$$\xi = \alpha + i\beta, \quad \eta = \alpha - i\beta.$$
 (4.2.13)

First, we transform equations (4.1.10). We then have

$$A^{**}(\alpha,\beta) u_{\alpha\alpha} + B^{**}(\alpha,\beta) u_{\alpha\beta} + C^{**}(\alpha,\beta) u_{\beta\beta} = H_4(\alpha,\beta,u,u_\alpha,u_\beta),$$
(4.2.14)

in which the coefficients assume the same form as the coefficients in equation (4.1.11). With the use of (4.2.13), the equations  $A^* = C^* = 0$  become

$$(A\alpha_x^2 + B\alpha_x\alpha_y + C\alpha_y^2) - (A\beta_x^2 + B\beta_x\beta_y + C\beta_y^2) + i [2A\alpha_x\beta_x + B(\alpha_x\beta_y + \alpha_y\beta_x) + 2C\alpha_y\beta_y] = 0,$$

### CLASS: II B.Sc COURSE CODE: 17MMU402

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: II BATCH-2017-2020

$$(A\alpha_x^2 + B\alpha_x\alpha_y + C\alpha_y^2) - (A\beta_x^2 + B\beta_x\beta_y + C\beta_y^2) -i [2A\alpha_x\beta_x + B(\alpha_x\beta_y + \alpha_y\beta_x) + 2C\alpha_y\beta_y] = 0,$$

or,

$$(A^{**} - C^{**}) + iB^{**} = 0, \qquad (A^{**} - C^{**}) - iB^{**} = 0.$$

These equations are satisfied if and only if

$$A^{**} = C^{**}$$
 and  $B^{**} = 0$ .

Hence, equation (4.2.14) transforms into the form

$$A^{**}u_{\alpha\alpha} + A^{**}u_{\beta\beta} = H_4(\alpha, \beta, u, u_\alpha, u_\beta).$$

Dividing through by  $A^{**}$ , we obtain

$$u_{\alpha\alpha} + u_{\beta\beta} = H_5(\alpha, \beta, u, u_{\alpha}, u_{\beta}),$$
 (4.2.15)

where  $H_5 = (H_4/A^{**})$ . This is called the *canonical form of the elliptic equation*.

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIBATCH-2017-2020

Example 4.2.1. Consider the equation

$$y^2 u_{xx} - x^2 u_{yy} = 0.$$

Here

$$A = y^2$$
,  $B = 0$ ,  $C = -x^2$ .

Thus,

$$B^2 - 4AC = 4x^2y^2 > 0.$$

The equation is hyperbolic everywhere except on the coordinate axes x = 0and y = 0. From the characteristic equations (4.2.5) and (4.2.6), we have

$$\frac{dy}{dx} = \frac{x}{y}, \qquad \frac{dy}{dx} = -\frac{x}{y}.$$

After integration of these equations, we obtain

$$\frac{1}{2}y^2 - \frac{1}{2}x^2 = c_1, \qquad \frac{1}{2}y^2 + \frac{1}{2}x^2 = c_2.$$

The first of these curves is a family of hyperbolas

$$\frac{1}{2}y^2 - \frac{1}{2}x^2 = c_1,$$

and the second is a family of circles

$$\frac{1}{2}y^2 + \frac{1}{2}x^2 = c_2.$$

To transform the given equation to canonical form, we consider

$$\xi = \frac{1}{2}y^2 - \frac{1}{2}x^2, \qquad \eta = \frac{1}{2}y^2 + \frac{1}{2}x^2.$$

From the relations (4.1.6), we have

$$u_{x} = u_{\xi}\xi_{x} + u_{\eta}\eta_{x} = -xu_{\xi} + xu_{\eta},$$
  

$$u_{y} = u_{\xi}\xi_{y} + u_{\eta}\eta_{y} = yu_{\xi} + yu_{\eta},$$
  

$$u_{xx} = u_{\xi\xi}\xi_{x}^{2} + 2u_{\xi\eta}\xi_{x}\eta_{x} + u_{\eta\eta}\eta_{x}^{2} + u_{\xi}\xi_{xx} + u_{\eta}\eta_{xx}$$
  

$$= x^{2}u_{\xi\xi} - 2x^{2}u_{\xi\eta} + x^{2}u_{\eta\eta} - u_{\xi} + u_{\eta}.$$

Prepared by K.Aarthiya, Asst Prof, Department of Mathematics, KAHE

Page 7/9

CLASS: II B.Sc COURSE CODE: 17MMU402

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: II BATCH-2017-2020

$$u_{yy} = u_{\xi\xi}\xi_y^2 + 2u_{\xi\eta}\xi_y\eta_y + u_{\eta\eta}\eta_y^2 + u_{\xi}\xi_{yy} + u_{\eta}\eta_{yy}$$
  
=  $y^2 u_{\xi\xi} + 2y^2 u_{\xi\eta} + y^2 u_{\eta\eta} + u_{\xi} + u_{\eta}.$ 

Thus, the given equation assumes the canonical form

$$u_{\xi\eta} = \frac{\eta}{2(\xi^2 - \eta^2)} u_{\xi} - \frac{\xi}{2(\xi^2 - \eta^2)} u_{\eta}.$$

Example 4.2.2. Consider the partial differential equation

$$x^2 u_{xx} + 2xy \, u_{xy} + y^2 u_{yy} = 0.$$

In this case, the discriminant is

$$B^2 - 4AC = 4x^2y^2 - 4x^2y^2 = 0.$$

The equation is therefore parabolic everywhere. The characteristic equation is

$$\frac{dy}{dx} = \frac{y}{x},$$

and hence, the characteristics are

$$\frac{y}{x} = c,$$

which is the equation of a family of straight lines.

Consider the transformation

$$\xi = \frac{y}{x}, \quad \eta = y$$

where  $\eta$  is chosen arbitrarily. The given equation is then reduced to the canonical form

$$y^2 u_{\eta\eta} = 0$$

Thus,

$$u_{\eta\eta} = 0$$
 for  $y \neq 0$ .

Prepared by K.Aarthiya, Asst Prof, Department of Mathematics, KAHE

Page 8/9

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIBATCH-2017-2020

Example 4.2.3. The equation

$$u_{xx} + x^2 u_{yy} = 0,$$

is elliptic everywhere except on the coordinate axis x = 0 because

$$B^2 - 4AC = -4x^2 < 0, \quad x \neq 0.$$

The characteristic equations are

$$\frac{dy}{dx} = ix, \qquad \frac{dy}{dx} = -ix.$$

Integration yields

$$2y - ix^2 = c_1, \qquad 2y + ix^2 = c_2.$$

Thus, if we write

$$\xi = 2y - ix^2, \qquad \eta = 2y + ix^2,$$

and hence,

$$\alpha = \frac{1}{2}(\xi + \eta) = 2y, \qquad \beta = \frac{1}{2i}(\xi - \eta) = -x^2,$$

we obtain the canonical form

$$u_{\alpha\alpha} + u_{\beta\beta} = -\frac{1}{2\beta} \, u_{\beta}.$$

It should be remarked here that a given partial differential equation may be of a different type in a different domain. Thus, for example, *Tricomi's equation* 

$$u_{xx} + xu_{yy} = 0, (4.2.16)$$

is elliptic for x > 0 and hyperbolic for x < 0, since  $B^2 - 4AC = -4x$ . For a detailed treatment, see Hellwig (1964).

QUESTION	OPT 1	OPT 2	OPT 3	OPT 4	ANSWER
A differential equation involving of one or more variables	Ordinary	ordinary	partial	partial variales	Ordinary
with respect to a single independent variable is called ODE	derivatives	variales	derivatives		derivatives
An equator which is is called non linear	not linear	dependent	linear	independent	not linear
Initial value prolem is also called	one point	two point	simply	wave equation	one point
initial value protein is also caned	boundary	boundary	boundary	wave equation	boundary
	value prolem	value problem	value problem		value prolem
Simply boundary value problem is also called	initial value	one point	two point	transport	two point
simply boundary value problem is also caned	prolem	boundary	boundary	equation	boundary
	protein	value prolem	value problem	equation	value problem
An equator involving derivatives of one or more dependent	homogeneous	non	quasi linear	linear	homogeneous
variables with respect to one or more independent variable is called	nomogeneous	homogeneous	quasi-inteat	inical	nomogeneous
· · · · · · · · · · · · · · · · · · ·					
Non-linear ODE is an ODE that is	linear	non-linear	homogeneous	non-	non-linear
ODE of order n is the dependent variable y and the	linear	non-linear	homogeneous	non-	linear
independent variable x is an equation				homogeneous	
An equation involving of one or more dependent variables	equation	derivative	euler	linear	derivatives
with respect to one or more independent variables is called					
A differential equation involving partial equation of one or more	one or more	two or more	more than one	more than two	more than one
variables with respect to independent variable is called PDE					
The order of the highest ordered derivatives involves in a	order of	dagraa of	product of	all the chore	order of
differential equation is called	differential	differential	differential	an the above	differential
unterential equation is called	equation	equation	equation		equation
Madhamadian I much law in an id da ha mull man diff id addiefing dha				-11 (h h	
whether the problem is said to be well posed if it satisfies the	atteast	annost	more man	an the above	annost
Initial value prolom is also called	houndary	anahy	all the chore	none of these	annahu
initial value protein is also caneu	value problem	problem	an the above	none of these	problem
A partial differential equation is the equation involving partial	one most one	atleast one	more than two	more than one	more than one
A partial unreferring equation is the equation involving partial derivatives of one or more dependent variables with respect to	one most one	alleast olle	more mail two	more man one	more than one
independent variable					
A differential equation involving ordinary derivatives of one or	1	2	3	4	1
more dependent variables with respect to a			-		
independent variable is called an ordinary differential equation.					
· · · · · · · · · · · · · · · · · · ·					
The equation $u_x u_{xx} + x u u_y = \sin y$ is	quasi linear	non linear	Cauchy	boundary	quasi linear
The order of the equation $u_{xx} - u_{yy} = 0$ is	1	2	3	4	2
The partial differential equation $u_t - u_{xx} = 0$ with $u(x, 0) = \sin x$ .	initial	boundary	Cauchy	linear	Cauchy
The given condition is called					
$F(x, y, u, u_{xx}, u_{yy}) = 0$ is order PDE	first	second	third	fourth	second
The equation $F(x, y, u, u_x, u_y) = 0$ is said to be	linear	non linear	quasi-linear	non quasi	linear
Which one of the following is the homogeneous equation?	xy + x = e	ax + by = 0	nx + my =	yx = y + 3x	ax + by = 0
The existence of the mathematical problem have	one most one	more than two	atleast one	more than one	atleast one
solution.					
L + M = M + L & LM = ML are called property.	associative	commutative	distributive	closed	commutative
Which of an operator does not satisfies the linear condition	L(cu) = c L(u)	$L(u_1 + u_2) =$	$L(c_1u_1 + c_2u_2)$	L(c1u1 +	L(c1u1 +
		$L(u_1) + L(u_2)$	$= c_1 L(u_1) +$	c2u2)≠	c2u2)≠
$(\mathbf{I} + \mathbf{M}) + \mathbf{N} - \mathbf{I} + (\mathbf{M} + \mathbf{N})$ is called property	acconistiva	distriutive	commutative	closed	associative
$(\mathbf{L} + \mathbf{M}) + \mathbf{M} = \mathbf{L} + (\mathbf{M} + \mathbf{M})$ is called property.	associative	uisuiuuive	commutative	cioseu	associative
A non-linear ordinary differential equation is an ordinary	linear	non linear	differential	integral	linear
A non mear ordinary unreferitiar equation is an ordinary					
differential equation that is not					
differential equation that is not Linear ordinary differential equations are further classified	single	dependent	independent	constant	dependent
Linear ordinary differential equation is an ordinary Linear ordinary differential equations are further classified according to the nature of the coefficients of the	single	dependent	independent	constant	dependent
A non-inteal ordinary differential equation is an ordinary differential equation that is not Linear ordinary differential equations are further classified according to the nature of the coefficients of the variables and its derivatives	single	dependent	independent	constant	dependent
A non-inical ordinary differential equation is an ordinary differential equation that is not Linear ordinary differential equations are further classified according to the nature of the coefficients of the variables and its derivatives The order of derivatives involved in the differential	single zero	dependent lowest	independent highest	constant	dependent highest

The equation $f(x, y, z, a, b) = 0$ containing two arbitrary parameters is called of an equation.	linear	non linear	complete solution	partial solution	complete solution
A solution which is not everywhere differentiable is called a	strong	weak	low	high	weak
A curve in (x, y, u)-space, whose tangent at every point coincides with the characteristic direction field (a, b, c), is called a	tangent curve	normal curve	characteristic curve	uniform curve	characteristic curve
The characteristic direction is also called as	monge axis	monge curve	monge line	monge line	monge axis
In the equation $u(x_0(t), y_0(t)) = u_0(t)$ , $u_0(t)$ is called the	final data	initial data	curve value	null value	initial data
The equation $f(x, y, z, a, b) = \_$ containing two arbitrary parameters is called linear of an equation.	0	1	2	3	0
The characteristic direction is also called as monge	axis	line	curve value	curve	axis
Linear ordinary differential equations are further classified according to the nature of the coefficients of the dependent	constant	variable	derivatives	equations	variable
The equation $f(x, y, z, a, b) = 0$ containing arbitrary parameters is called linear of an equation.	one	two	three	fourth	two
The equation $f(x, y, z, a, b) = 0$ containing two arbitrary parameters is called linear of an	line	equation	point	graph	equation
value prolem is also called cauchy problem	boundary value problem	initial	all the above	none of these	initial
The order of highest derivatives involved in the differential equations is called of the differential equation	order	variable	constant	linear	order
The order of highest derivatives involved in the equations is called order of the differential equation	differential	linear	non linear	quasi linear	differential

### CLASS: II B.Sc COURSE CODE: 17MMU402

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: III BATCH-2017-2020

### UNIT III SYLLABUS

Reduction of second order Linear Equations to canonical forms- The Cauchy problem- The Cauchy-Kowaleewskaya theorem -Cauchy problem of an infinite string - Initial Boundary Value Problems -Semi-Infinite String with a fixed end -Semi-Infinite String with a Free end- Equations with non-homogeneous boundary conditions -Non- Homogeneous Wave Equation.

## **Reduction of second order Linear Equations to canonical forms:**

# The Cauchy Problem

In the theory of ordinary differential equations, by the initial-value problem we mean the problem of finding the solutions of a given differential equation with the appropriate number of initial conditions prescribed at an initial point. For example, the second-order ordinary differential equation

$$\frac{d^2u}{dt^2} = f\left(t, u, \frac{du}{dt}\right)$$

and the initial conditions

$$u(t_0) = \alpha, \qquad \left(\frac{du}{dt}\right)(t_0) = \beta,$$

constitute an initial-value problem.

An analogous problem can be defined in the case of partial differential equations. Here we shall state the problem involving second-order partial differential equations in two independent variables.

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

We consider a second-order partial differential equation for the function u in the independent variables x and y, and suppose that this equation can be solved explicitly for  $u_{yy}$ , and hence, can be represented in the from

$$u_{yy} = F(x, y, u, u_x, u_y, u_{xx}, u_{xy}).$$
(5.1.1)

For some value  $y = y_0$ , we prescribe the initial values of the unknown function and of the derivative with respect to y

$$u(x, y_0) = f(x), \qquad u_y(x, y_0) = g(x).$$
 (5.1.2)

The problem of determining the solution of equation (5.1.1) satisfying the initial conditions (5.1.2) is known as the *initial-value problem*. For instance, the initial-value problem of a vibrating string is the problem of finding the solution of the wave equation

$$u_{tt} = c^2 u_{xx},$$

satisfying the initial conditions

$$u(x,t_0) = u_0(x), \qquad u_t(x,t_0) = v_0(x),$$

where  $u_0(x)$  is the initial displacement and  $v_0(x)$  is the initial velocity.

In initial-value problems, the initial values usually refer to the data assigned at  $y = y_0$ . It is not essential that these values be given along the line  $y = y_0$ ; they may very well be prescribed along some curve  $L_0$  in the xy plane. In such a context, the problem is called the *Cauchy problem* instead of the initial-value problem, although the two names are actually synonymous.

We consider the Euler equation

$$Au_{xx} + Bu_{xy} + Cu_{yy} = F(x, y, u, u_x, u_y), \qquad (5.1.3)$$

where A, B, C are functions of x and y. Let  $(x_0, y_0)$  denote points on a smooth curve  $L_0$  in the xy plane. Also let the parametric equations of this curve  $L_0$  be

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

where A, B, C are functions of x and y. Let  $(x_0, y_0)$  denote points on a smooth curve  $L_0$  in the xy plane. Also let the parametric equations of this curve  $L_0$  be

$$x_0 = x_0(\lambda), \qquad y_0 = y_0(\lambda), \qquad (5.1.4)$$

where  $\lambda$  is a parameter.

We suppose that two functions  $f(\lambda)$  and  $g(\lambda)$  are prescribed along the curve  $L_0$ . The Cauchy problem is now one of determining the solution u(x, y) of equation (5.1.3) in the neighborhood of the curve  $L_0$  satisfying the Cauchy conditions

$$u = f(\lambda), \qquad (5.1.5a)$$

$$\frac{\partial u}{\partial n} = g\left(\lambda\right),$$
 (5.1.5b)

on the curve  $L_0$  where *n* is the direction of the normal to  $L_0$  which lies to the left of  $L_0$  in the counterclockwise direction of increasing arc length. The function  $f(\lambda)$  and  $g(\lambda)$  are called the *Cauchy data*.

For every point on  $L_0$ , the value of u is specified by equation (5.1.5a). Thus, the curve  $L_0$  represented by equation (5.1.4) with the condition (5.1.5a) yields a twisted curve L in (x, y, u) space whose projection on the xy plane is the curve  $L_0$ . Thus, the solution of the Cauchy problem is a surface, called an *integral surface*, in the (x, y, u) space passing through Land satisfying the condition (5.1.5b), which represents a tangent plane to the integral surface along L.

If the function  $f(\lambda)$  is differentiable, then along the curve  $L_0$ , we have

$$\frac{du}{d\lambda} = \frac{\partial u}{\partial x}\frac{dx}{d\lambda} + \frac{\partial u}{\partial y}\frac{dy}{d\lambda} = \frac{df}{d\lambda},$$
(5.1.6)

and

$$\frac{\partial u}{\partial n} = \frac{\partial u}{\partial x}\frac{dx}{dn} + \frac{\partial u}{\partial y}\frac{dy}{dn} = g,$$
(5.1.7)

but

$$\frac{dx}{dn} = -\frac{dy}{ds}$$
 and  $\frac{dy}{dn} = \frac{dx}{ds}$ . (5.1.8)

Prepared by K.Aarthiya, Asst Prof, Department of Mathematics, KAHE

Page 3/14

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

Equation (5.1.7) may be written as

$$\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial x}\frac{dy}{ds} + \frac{\partial u}{\partial y}\frac{dx}{ds} = g.$$
(5.1.9)

Since

$$\begin{vmatrix} \frac{dx}{d\lambda} & \frac{dy}{d\lambda} \\ \frac{-dy}{ds} & \frac{dx}{ds} \end{vmatrix} = \frac{(dx)^2 + (dy)^2}{ds \, d\lambda} \neq 0,$$
(5.1.10)

it is possible to find  $u_x$  and  $u_y$  on  $L_0$  from the system of equations (5.1.6) and (5.1.9). Since  $u_x$  and  $u_y$  are known on  $L_0$ , we find the higher derivatives by first differentiating  $u_x$  and  $u_y$  with respect to  $\lambda$ . Thus, we have

$$\frac{\partial^2 u}{\partial x^2} \frac{dx}{d\lambda} + \frac{\partial^2 u}{\partial x \partial y} \frac{dy}{d\lambda} = \frac{d}{d\lambda} \left(\frac{\partial u}{\partial x}\right), \qquad (5.1.11)$$

$$\frac{\partial^2 u}{\partial x \,\partial y} \frac{dx}{d\lambda} + \frac{\partial^2 u}{\partial y^2} \frac{dy}{d\lambda} = \frac{d}{d\lambda} \left(\frac{\partial u}{\partial y}\right). \tag{5.1.12}$$

From equation (5.1.3), we have

$$A\frac{\partial^2 u}{\partial x^2} + B\frac{\partial^2 u}{\partial x \partial y} + C\frac{\partial^2 u}{\partial y^2} = F,$$
(5.1.13)

where F is known since  $u_x$  and  $u_y$  have been found. The system of equations can be solved for  $u_{xx}$ ,  $u_{xy}$ , and  $u_{yy}$ , if

$$\begin{vmatrix} \frac{dx}{d\lambda} & \frac{dy}{d\lambda} & 0 \\ 0 & \frac{dx}{d\lambda} & \frac{dy}{d\lambda} \\ A & B & C \end{vmatrix} = C \left(\frac{dx}{d\lambda}\right)^2 - B\left(\frac{dx}{d\lambda}\right) \left(\frac{dy}{d\lambda}\right) + A\left(\frac{dy}{d\lambda}\right)^2 \neq 0. \quad (5.1.14)$$

The equation

$$A\left(\frac{dy}{dx}\right)^2 - B\left(\frac{dy}{dx}\right) + C = 0, \qquad (5.1.15)$$

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

is called the *characteristic equation*. It is then evident that the necessary condition for obtaining the second derivatives is that the curve  $L_0$  must not be a characteristic curve.

If the coefficients of equation (5.1.3) and the function (5.1.5) are analytic, then all the derivatives of higher orders can be computed by the above process. The solution can then be represented in the form of a Taylor series:

$$u(x,y) = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{1}{k! (n-k)!} \frac{\partial^n u_0}{\partial x_0^k \partial y_0^{n-k}} (x-x_0)^k (y-y_0)^{n-k}, (5.1.16)$$

which can be shown to converge in the neighborhood of the curve  $L_0$ . Thus, we may state the famous Cauchy–Kowalewskaya theorem.

# The Cauchy–Kowalewskaya Theorem

Let the partial differential equation be given in the form

$$u_{yy} = F(y, x_1, x_2, \dots, x_n, u, u_y, u_{x_1}, u_{x_2}, \dots, u_{x_n}, u_{x_1y}, u_{x_2y}, \dots, u_{x_ny}, u_{x_1x_1}, u_{x_2x_2}, \dots, u_{x_nx_n}), (5.2.1)$$

and let the initial conditions

$$u = f(x_1, x_2, \dots, x_n),$$
 (5.2.2)

$$u_y = g(x_1, x_2, \dots, x_n), \qquad (5.2.3)$$

be given on the noncharacteristic manifold  $y = y_0$ .

1

If the function F is analytic in some neighborhood of the point  $(y^0, x_1^0, x_2^0, \ldots, x_n^0, u^0, u_y^0, \ldots)$  and if the functions f and g are analytic in some neighborhood of the point  $(x_1^0, x_2^0, \ldots, x_n^0)$ , then the Cauchy problem has a unique analytic solution in some neighborhood of the point  $(y^0, x_1^0, x_2^0, \ldots, x_n^0)$ .

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

For the proof, see Petrovsky (1954).

The preceding statement seems equally applicable to hyperbolic, parabolic, or elliptic equations. However, we shall see that difficulties arise in formulating the Cauchy problem for nonhyperbolic equations. Consider, for instance, the famous Hadamard (1952) example.

The problem consists of the elliptic (or Laplace) equation

$$u_{xx} + u_{yy} = 0,$$

and the initial conditions on y = 0

u(x,0) = 0,  $u_u(x,0) = n^{-1} \sin nx.$ 

The solution of this problem is

$$u(x,y) = n^{-2} \sinh ny \sin nx,$$

which can be easily verified.

It can be seen that, when n tends to infinity, the function  $n^{-1} \sin nx$  tends uniformly to zero. But the solution  $n^{-2} \sinh ny \sin nx$  does not become small, as n increases for any nonzero y. Physically, the solution represents an oscillation with unbounded amplitude  $(n^{-2} \sinh ny)$  as  $y \to \infty$  for any fixed x. Even if n is a fixed number, this solution is unstable in the sense that  $u \to \infty$  as  $y \to \infty$  for any fixed x for which  $\sin nx \neq 0$ . It is obvious then that the solution does not depend continuously on the data. Thus, it is not a properly posed problem.

# Initial Boundary-Value Problems

### (A) Semi-infinite String with a Fixed End

Let us first consider a semi-infinite vibrating string with a fixed end, that is,

$$u_{tt} = c^2 u_{xx}, \qquad 0 < x < \infty, \qquad t > 0, u(x,0) = f(x), \qquad 0 \le x < \infty, \qquad (5.4.1)$$

CLASS: II B.Sc	<b>COURSE NAME: PDE AND</b>	SYSTEM OF ODE
COURSE CODE: 17MMU402	UNIT: III	BATCH-2017-2020

$u_{t}\left( x,0\right) =g\left( x\right) ,$	$0 \le x < \infty,$
u(0,t) = 0,	$0 \le t \le \infty$ .

It is evident here that the boundary condition at x = 0 produces a wave moving to the right with the velocity c. Thus, for x > ct, the solution is the same as that of the infinite string, and the displacement is influenced only by the initial data on the interval [x - ct, x + ct], as shown in Figure 5.4.1.

When x < ct, the interval [x - ct, x + ct] extends onto the negative x-axis where f and g are not prescribed.

But from the d'Alembert formula

$$u(x,t) = \phi(x+ct) + \psi(x-ct),$$
 (5.4.2)

where



Figure 5.4.1 Displacement influenced by the initial data on [x - ct, x + ct].

CLASS: II B.Sc COURSE CODE: 17MMU402	COURSE NAME: P UNIT: III	DE AND SYSTEM OF ODE BATCH-2017-2020
$\phi\left(\xi ight)$ :	$= \frac{1}{2}f(\xi) + \frac{1}{2c}\int_{0}^{\xi} g(\tau)  d\tau$	$+\frac{K}{2},$ (5.4.3)
$\psi\left(\eta ight)$ :	$=\frac{1}{2}f\left(\eta\right)-\frac{1}{2c}\int_{0}^{\eta}g\left(\tau\right)d\tau$	$-\frac{K}{2}$ , (5.4.4)
we see that		
u	$(0,t) = \phi(ct) + \psi(-ct) =$	: 0.
Hence,		
	$\psi\left(-ct\right) = -\phi\left(ct\right).$	
If we let $\alpha = -ct$ , then		
	$\psi\left(\alpha\right) = -\phi\left(-\alpha\right).$	
Replacing $\alpha$ by $x - ct$ , w	we obtain for $x < ct$ ,	
	$\psi\left(x-ct\right) = -\phi\left(ct-x\right)$	,
and hence,		
$\psi\left(x-ct\right) =$	$-\frac{1}{2}f(ct-x) - \frac{1}{2c}\int_{0}^{ct-x}$	$g\left(\tau\right)d\tau - \frac{K}{2}.$
The solution of the initia	al boundary-value proble	m, therefore, is given by
$u(x,t) = \frac{1}{2} [f(x+ct) + f(x+ct)]$	$f(x-ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} g(t) dt$	$(\tau) d\tau$ for $x > ct$ , (5.4.5)

$$u(x,t) = \frac{1}{2} \left[ f(x+ct) - f(ct-x) \right] + \frac{1}{2c} \int_{ct-x}^{x+ct} g(\tau) \, d\tau \quad \text{for } x < ct. \ (5.4.6)$$

In order for this solution to exist, f must be twice continuously differentiable and g must be continuously differentiable, and in addition

$$f(0) = f''(0) = g(0) = 0.$$

Solution (5.4.6) has an interesting physical interpretation. If we draw the characteristics through the point  $(x_0, t_0)$  in the region x > ct, we see, as pointed out earlier, that the displacement at  $(x_0, t_0)$  is determined by the initial values on  $[x_0 - ct_0, x_0 + ct_0]$ .

Prepared by K.Aarthiya, Asst Prof, Department of Mathematics, KAHE

Page 8/14

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

 $Example \ 5.4.1.$  Determine the solution of the initial boundary-value problem

$$u_{tt} = 4 u_{xx}, \qquad x > 0, \qquad t > 0,$$
  
$$u(x,0) = |\sin x|, \qquad x > 0,$$
  
$$u_t(x,0) = 0, \qquad x \ge 0,$$
  
$$u(x,0) = 0, \qquad t \ge 0.$$

For x > 2t,

$$u(x,t) = \frac{1}{2} \left[ f(x+2t) + f(x-2t) \right]$$
  
=  $\frac{1}{2} \left[ |\sin(x+2t)| - |\sin(x-2t)| \right],$ 

and for x < 2t,

$$u(x,t) = \frac{1}{2} \left[ f(x+2t) - f(2t-x) \right]$$
  
=  $\frac{1}{2} \left[ |\sin(x+2t)| - |\sin(2t-x)| \right]$ 

Notice that u(0,t) = 0 is satisfied by u(x,t) for x < 2t (that is, t > 0).

### (B) Semi-infinite String with a Free End

We consider a semi-infinite string with a free end at x = 0. We will determine the solution of

 $u_{tt} = c^2 u_{xx}, \qquad 0 < x < \infty, \qquad t > 0,$   $u(x,0) = f(x), \qquad 0 \le x < \infty,$   $u_t(x,0) = g(x), \qquad 0 \le x < \infty,$   $u_x(0,t) = 0, \qquad 0 \le t < \infty.$ (5.4.7)

As in the case of the fixed end, for x > ct the solution is the same as that of the infinite string. For x < ct, from the d'Alembert solution (5.4.2)

$$u(x,t) = \phi(x+ct) + \psi(x-ct),$$

# CLASS: II B.ScCOURSE NAME: PCOURSE CODE: 17MMU402UNIT: III

## COURSE NAME: PDE AND SYSTEM OF ODE UNIT: III BATCH-2017-2020

we have

$$u_x(x,t) = \phi'(x+ct) + \psi'(x-ct).$$

Thus,

$$u_x(0,t) = \phi'(ct) + \psi'(-ct) = 0.$$

Integration yields

$$\phi\left(ct\right) - \psi\left(-ct\right) = K,$$

where K is a constant. Now, if we let  $\alpha = -ct$ , we obtain

 $\psi\left(\alpha\right) = \phi\left(-\alpha\right) - K.$ 

Replacing  $\alpha$  by x - ct, we have

$$\psi\left(x - ct\right) = \phi\left(ct - x\right) - K,$$

and hence,

$$\psi(x - ct) = \frac{1}{2} f(ct - x) + \frac{1}{2c} \int_0^{ct - x} g(\tau) \, d\tau - \frac{K}{2}.$$

The solution of the initial boundary-value problem, therefore, is given by

$$\begin{aligned} u(x,t) &= \frac{1}{2} \left[ f\left(x+ct\right) + f\left(x-ct\right) \right] + \frac{1}{2c} \int_{x-ct}^{x+ct} g\left(\tau\right) d\tau \quad \text{for } x > ct. \ (5.4.8) \\ u(x,t) &= \frac{1}{2} \left[ f\left(x+ct\right) + f\left(ct-x\right) \right] + \frac{1}{2c} \left[ \int_{0}^{x+ct} g\left(\tau\right) d\tau + \int_{0}^{ct-x} g\left(\tau\right) d\tau \right] \\ \text{for } x < ct. \ (5.4.9) \end{aligned}$$

We note that for this solution to exist, f must be twice continuously differentiable and g must be continuously differentiable, and in addition,

$$f'(0) = g'(0) = 0.$$

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

Example 5.4.2. Find the solution of the initial boundary-value problem

$$u_{tt} = u_{xx}, \qquad 0 < x < \infty, \qquad t > 0,$$
  
$$u(x,0) = \cos\left(\frac{\pi x}{2}\right), \qquad 0 \le x < \infty,$$
  
$$u_t(x,0) = 0, \qquad 0 \le x < \infty,$$
  
$$u_x(x,0) = 0, \qquad t \ge 0.$$

For x > t

$$u(x,t) = \frac{1}{2} \left[ \cos \frac{\pi}{2} \left( x+t \right) + \cos \frac{\pi}{2} \left( x-t \right) \right]$$
$$= \cos \left( \frac{\pi}{2} x \right) \cos \left( \frac{\pi}{2} t \right),$$

and for x < t

$$u(x,t) = \frac{1}{2} \left[ \cos \frac{\pi}{2} \left( x+t \right) + \cos \frac{\pi}{2} \left( t-x \right) \right]$$
$$= \cos \left( \frac{\pi}{2} x \right) \cos \left( \frac{\pi}{2} t \right).$$

# Equations with Nonhomogeneous Boundary Conditions

In the case of the initial boundary-value problems with nonhomogeneous boundary conditions, such as

$$u_{tt} = c^2 u_{xx}, \qquad x > 0, \qquad t > 0,$$
  

$$u(x,0) = f(x), \qquad x \ge 0,$$
  

$$u_t(x,0) = g(x), \qquad x \ge 0,$$
  

$$u(0,t) = p(t), \qquad t \ge 0,$$
  
(5.5.1)
# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

we proceed in a manner similar to the case of homogeneous boundary conditions. Using equation (5.4.2), we apply the boundary condition to obtain

$$u(0,t) = \phi(ct) + \psi(-ct) = p(t)$$
.

If we let  $\alpha = -ct$ , we have

$$\psi(\alpha) = p\left(-\frac{\alpha}{c}\right) - \phi(-\alpha).$$

Replacing  $\alpha$  by x - ct, the preceding relation becomes

$$\psi(x - ct) = p\left(t - \frac{x}{c}\right) - \phi(ct - x)$$

Thus, for  $0 \le x < ct$ ,

$$u(x,t) = p\left(t - \frac{x}{c}\right) + \frac{1}{2}\left[f\left(x + ct\right) - f\left(ct - x\right)\right] + \frac{1}{2c}\int_{ct - x}^{x + ct} g\left(\tau\right)d\tau$$
  
=  $p\left(t - \frac{x}{c}\right) + \phi\left(x + ct\right) - \psi\left(ct - x\right),$  (5.5.2)

where  $\phi(x + ct = \xi)$  is given by (5.3.11), and  $\psi(\eta)$  is given by

$$\psi(\eta) = \frac{1}{2}f(\eta) + \frac{1}{2c}\int_0^{\eta} g(\tau) \,d\tau.$$
 (5.5.3)

The solution for x > ct is given by the solution (5.4.5) of the infinite string.

In this case, in addition to the differentiability conditions satisfied by f and g, as in the case of the problem with the homogeneous boundary conditions, p must be twice continuously differentiable in t and

$$p(0) = f(0), \qquad p'(0) = g(0), \qquad p''(0) = c^2 f''(0).$$

We next consider the initial boundary-value problem

$$u_{tt} = c^2 u_{xx}, \qquad x > 0, \qquad t > 0,$$
  

$$u(x,0) = f(x), \qquad x \ge 0,$$
  

$$u_t(x,0) = g(x), \qquad x \ge 0,$$
  

$$u_x(0,t) = q(t), \qquad t \ge 0.$$

# CLASS: II B.ScCOURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IIIBATCH-2017-2020

Using (5.4.2), we apply the boundary condition to obtain

$$u_x(0,t) = \phi'(ct) + \psi'(-ct) = q(t).$$

Then, integrating yields

$$\phi(ct) - \psi(-ct) = c \int_0^t q(\tau) \, d\tau + K.$$

If we let  $\alpha = -ct$ , then

$$\psi(\alpha) = \phi(-\alpha) - c \int_0^{-\alpha/c} q(\tau) \, d\tau - K.$$

Replacing  $\alpha$  by x - ct, we obtain

$$\psi(x - ct) = \phi(ct - x) - c \int_0^{t - x/c} q(\tau) d\tau - K.$$

The solution of the initial boundary-value problem for x < ct, therefore, is given by

$$u(x,t) = \frac{1}{2} \left[ f(x+ct) + f(ct-x) \right] + \frac{1}{2c} \left[ \int_0^{x+ct} g(\tau) \, d\tau + \int_0^{ct-x} g(\tau) \, d\tau \right] -c \int_0^{t-x/c} q(\tau) \, d\tau. \quad (5.5.4)$$

Here f and g must satisfy the differentiability conditions, as in the case of the problem with the homogeneous boundary conditions. In addition

$$f'(0) = q(0), \qquad g'(0) = q'(0).$$

The solution for the initial boundary-value problem involving the boundary condition

$$u_x(0,t) + h u(0,t) = 0, \qquad h = \text{constant}$$

can also be constructed in a similar manner from the d'Alembert solution.

CLASS: II B.Sc COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: III BATCH-2017-2020

# Nonhomogeneous Wave Equations

We shall consider next the Cauchy problem for the nonhomogeneous wave equation

$$u_{tt} = c^2 u_{xx} + h^* (x, t), \qquad (5.7.1)$$

with the initial conditions

$$u(x,0) = f(x), \qquad u_t(x,0) = g^*(x).$$
 (5.7.2)

By the coordinate transformation

$$y = ct$$
, (5.7.3)

the problem is reduced to

$$u_{xx} - u_{yy} = h(x, y), \qquad (5.7.4)$$

$$u(x,0) = f(x),$$
 (5.7.5)

$$u_y(x,0) = g(x),$$
 (5.7.6)

where  $h(x, y) = -h^*/c^2$  and  $g(x) = g^*/c$ .

Let  $P_0(x_0, y_0)$  be a point of the plane, and let  $Q_0$  be the point  $(x_0, 0)$ on the initial line y = 0. Then the characteristics,  $x \pm y = \text{constant}$ , of equation (5.7.4) are two straight lines drawn through the point  $P_0$  with slopes  $\pm 1$ . Obviously, they intersect the x-axis at the points  $P_1(x_0 - y_0, 0)$ and  $P_2(x_0 + y_0, 0)$ , as shown in Figure 5.7.1. Let the sides of the triangle  $P_0P_1P_2$  be designated by  $B_0$ ,  $B_1$ , and  $B_2$ , and let D be the region representing the interior of the triangle and its boundaries B. Integrating both sides of equation (5.7.4), we obtain

$$\iint_{R} \left( u_{xx} - u_{yy} \right) dR = \iint_{R} h\left( x, y \right) dR.$$
 (5.7.7)

Now we apply Green's theorem to obtain

QUESTION	OPT 1	OPT 2
The method of separation of variables in perhaps the worldest method for soving PDE	systematic	transformation
The required solution of the PDE is then expossed as a product not equal to 0	u(x,t)=X(x)+T(t)	u(x,t)=X(x).T(t)
Complementry function is law the tension is constant	Ae <sup>m</sup> 1x+Be <sup>m</sup> 2x translation	u(x,t)=Y(x).Y(y) Hooke's law
The tension in the string is in the direction of the tangent to the	existing profile	elongation
Classification of second order PDE is said to be hyperbolic . Then the equation is	negative	equal
If alpha and beta are the angles made by the a and b respectively.	perpendicular	parallel
The discriminent $B^2$ -4AC=4 $y^2 x^2$ is	parabolic	eliptic
The string is and elastic	flexible	segment
The complete solution of PDE is u(x,y)=X(x)Y(y) (i.e)u=	xy	x+y
A solution to the partial differential equation $(\partial^2 u)/(\partial x^2)=9(\partial^2 u)/(\partial y^2)$ is	$\cos(3x - y)$	$x^2 + y^2$
The partial differential equation 5 $(\partial^2 z)/(\partial x^2)+6(\partial^2 z)/(\partial y^2)=xy$ is classified as	elliptic	hyperbolic
The partial differential equation xy $\partial z/\partial x=5(\partial^2 z)/(\partial y^2)$ ) is classified as	elliptic	hyperbolic
The partial differential equation $(\partial^2 z)/(\partial x^2)$ -5 $(\partial^2 z)/(\partial y^2)=0$ is classified as	elliptic	hyperbolic
A partial differential equation is one which involves derivatives	single	ordinary
The of PDE satisfies for all values of n	unity	existence
Z=X(x)Y(y) is called of variables	integration	seperation
The separation principle can readily be extended to number of variables	smaller	unique
The solution of PDE satisfies for all values of	n	1
If $f(x,p) = g(y,q)$ is calledequation	Clairaut	Charpit
L(z)+f(x,y,z,p,q)=0 where L is the	laplace	differential
Z=X(x)Y(y)T(z) is the extension of variables	integration	differentiation
The use of the theory of integral transforms is the of PDE	unity	existance
The method of separation of variables applied to diffusion equation is similar totheory	potential	grad
The order of PDE to be the order of the derivative of order occurring in it.	lowest	highest
Z=X(x)Y(y) is separable in the variables	x&y	x&z
In the method of integral transforms L denotesoperator.	nonlinear	linear
The separation principle can readily be extended to larger number of	constant	variable
The variational approach to value problem is useful in the derivation of approximating solution	Euclid	kernal

The use of the theory of integral transforms is the solution of	ode	pde
The use of the of integral transforms is the solution of pde	theory	integral
The use of the theory of integralis the solution of pde	transforms	integral
The use of the theory of integral transforms is the of pde	solution	integral
The use of the theory of transforms is the solution of pde	integral	transforms
The principle can readily be extended to larger number of variable	larger	variable
The separation principle can readily be extended to number of variable	larger	variable
The separation principle can readily be to larger number of variable	extended	variable
The separation principle can readily be extended to larger of variable	number	variable
The separation can readily be extended to larger number of variable	larger	variable
In the method of transforms L denotes linear operator.	nonlinear	integral

OPT 3	OPT 4	ANSWER
wave equation	heat equation	systamatic
X(x).Y(y)	u(x,t)=Y(x).Y(y)	u(x,t)=X(x).T(t)
Ae^m1x Newton's law	Be^m2x Faraday's law	Ae^m1x+Be^m2x Hooke's law
tension	center	existing profile
equal to zero	positive	positive
tangents	center	tangents
hyperbolic	all the aove	hyperbolic
elongation	none of these	flexible
x/y	y/x	xy
$\sin(3x - y)$	$e^{-3\pi x}\sin(\pi y)$	$e^{-3\pi x}\sin\left(\pi y\right)$
parabolic	circle	elliptic
parabolic	sphere	paraolic
parabolic	sphere	hyperbolic
partial	linear	partial
solution	formal	solution
differention	indution	seperation
larger	contrary	larger
2	3	n
Crout	separable	separable
lagrange	longdivision	differential
induction	separation	seperation
solution	formal	solution
calculus	electrostatic	potential
first	second	highest
y&z	x+y	x&y
constant	variable	linear
coefficient	sequence	variable
eigen	node	eigen

C.I	P.I	pde
differentiation	ode	theory
differentiation	ode	transforms
differentiation	ode	solution
differentiation	ode	integral
principle	separation	separation
principle	separation	larger
principle	separation	extended
principle	separation	number
principle	separation	principle
constant	variable	integral

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

# UNIT IV SYLLABUS

Method of separation of variables - Solving the Vibrating String - Problems-Solving the Heat Conduction problem - Systems of linear differential equations -Types of linear systems differential operators - an operator method for linear systems with constant coefficients.

# METHOD OF SEPARATION OF VARIABLES

# The Vibrating String

One of the most important problems in mathematical physics is the vibration of a stretched string. Simplicity and frequent occurrence in many branches of mathematical physics make it a classic example in the theory of partial differential equations.

Let us consider a stretched string of length l fixed at the end points. The problem here is to determine the equation of motion which characterizes the position u(x,t) of the string at time t after an initial disturbance is given.

In order to obtain a simple equation, we make the following assumptions:

- 1. The string is flexible and elastic, that is the string cannot resist bending moment and thus the tension in the string is always in the direction of the tangent to the existing profile of the string.
- 2. There is no elongation of a single segment of the string and hence, by Hooke's law, the tension is constant.
- 3. The weight of the string is small compared with the tension in the string.
- 4. The deflection is small compared with the length of the string.
- 5. The slope of the displaced string at any point is small compared with unity.

Page 1/12

# CLASS: II B.Sc.COURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IVBATCH-2017-2020

6. There is only pure transverse vibration.

We consider a differential element of the string. Let T be the tension at the end points as shown in Figure 3.2.1. The forces acting on the element of the string in the vertical direction are

$$T\sin\beta - T\sin\alpha$$
.

By Newton's second law of motion, the resultant force is equal to the mass times the acceleration. Hence,

$$T\sin\beta - T\sin\alpha = \rho\,\delta s\,u_{tt} \tag{3.2.1}$$

where  $\rho$  is the line density and  $\delta s$  is the smaller arc length of the string. Since the slope of the displaced string is small, we have

$$\delta s \simeq \delta x.$$

Since the angles  $\alpha$  and  $\beta$  are small

$$\sin \alpha \simeq \tan \alpha$$
,  $\sin \beta \simeq \tan \beta$ .

Thus, equation (3.2.1) becomes

$$\tan\beta - \tan\alpha = \frac{\rho\,\delta x}{T}u_{tt}.\tag{3.2.2}$$

But, from calculus we know that  $\tan \alpha$  and  $\tan \beta$  are the slopes of the string at x and  $x + \delta x$ :

$$\tan \alpha = u_x\left(x,t\right)$$

and

$$\tan \beta = u_x \left( x + \delta x, t \right)$$

at time t. Equation (3.2.2) may thus be written as

$$\frac{1}{\delta x}\left[(u_x)_{x+\delta x} - (u_x)_x\right] = \frac{\rho}{T}u_{tt}, \quad \frac{1}{\delta x}\left[u_x\left(x+\delta x,t\right) - u_x\left(x,t\right)\right] = \frac{\rho}{T}u_{tt}.$$

# CLASS: II B.Sc.COURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IVBATCH-2017-2020

In the limit as  $\delta x$  approaches zero, we find

$$u_{tt} = c^2 u_{xx}$$
 (3.2.3)

where  $c^2 = T/\rho$ . This is called the one-dimensional wave equation.

If there is an external force f per unit length acting on the string. Equation (3.2.3) assumes the form

$$u_{tt} = c^2 u_{xx} + F, \quad F = f/\rho,$$
 (3.2.4)

where f may be pressure, gravitation, resistance, and so on.

# Conduction of Heat in Solids

We consider a domain  $D^*$  bounded by a closed surface  $B^*$ . Let u(x, y, z, t) be the temperature at a point (x, y, z) at time t. If the temperature is not constant, heat flows from places of higher temperature to places of lower temperature. Fourier's law states that the rate of flow is proportional to the gradient of the temperature. Thus the velocity of the heat flow in an isotropic body is

$$\mathbf{v} = -K \operatorname{grad} u, \qquad (3.5.1)$$

where K is a constant, called the *thermal conductivity* of the body.

Let D be an arbitrary domain bounded by a closed surface B in  $D^*$ . Then the amount of heat leaving D per unit time is

$$\iint_B v_n ds$$

where  $v_n = \mathbf{v} \cdot \mathbf{n}$  is the component of  $\mathbf{v}$  in the direction of the outer unit normal  $\mathbf{n}$  of B. Thus, by Gauss' theorem (Divergence theorem)

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

$$\iint_{B} v_n ds = \iiint_{D} \operatorname{div} (-K \operatorname{grad} u) \, dx \, dy \, dz$$
$$= -K \iiint_{D} \nabla^2 u \, dx \, dy \, dz. \tag{3.5.2}$$

But the amount of heat in D is given by

$$\iiint_D \sigma \rho u \, dx \, dy \, dz, \tag{3.5.3}$$

where  $\rho$  is the density of the material of the body and  $\sigma$  is its specific heat. Assuming that integration and differentiation are interchangeable, the rate of decrease of heat in D is

$$-\iiint_D \sigma \rho \frac{\partial u}{\partial t} \, dx \, dy \, dz. \tag{3.5.4}$$

Since the rate of decrease of heat in D must be equal to the amount of heat leaving D per unit time, we have

$$-\iiint_D \sigma \rho u_t \, dx \, dy \, dz = -K \iiint_D \nabla^2 u \, dx \, dy \, dz,$$

for an arbitrary D in  $D^*$ . We assume that the integrand is continuous. If we suppose that the integrand is not zero at a point  $(x_0, y_0, z_0)$  in D, then, by continuity, the integrand is not zero in a small region surrounding the point  $(x_0, y_0, z_0)$ . Continuing in this fashion we extend the region encompassing D. Hence the integral must be nonzero. This contradicts (3.5.5). Thus, the integrand is zero everywhere, that is,

$$u_t = \kappa \nabla^2 u, \qquad (3.5.6)$$

where  $\kappa = K/\sigma\rho$ . This is known as the *heat equation*.

# CLASS: II B.Sc.COURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: IVBATCH-2017-2020

# SYSTEMS OF LINEAR DIFFERENTIAL EQUATIONS

# DIFFERENTIAL OPERATORS AND AN OPERATOR METHOD

## A. Types of Linear Systems

We start by introducing the various types of linear systems that we shall consider. The general linear system of two first-order differential equations in two unknown functions x and y is of the form

$$a_{1}(t)\frac{dx}{dt} + a_{2}(t)\frac{dy}{dt} + a_{3}(t)x + a_{4}(t)y = F_{1}(t),$$
  

$$b_{1}(t)\frac{dx}{dt} + b_{2}(t)\frac{dy}{dt} + b_{3}(t)x + b_{4}(t)y = F_{2}(t).$$
(7.1)

We shall be concerned with systems of this type that have constant coefficients. An example of such a system is

$$2\frac{dx}{dt} + 3\frac{dy}{dt} - 2x + y = t^2, \qquad \frac{dx}{dt} - 2\frac{dy}{dt} + 3x + 4y = e^t.$$

We shall say that a solution of system (7.1) is an ordered pair of real functions (f, g) such that x = f(t), y = g(t) simultaneously satisfy both equations of the system (7.1) on some real interval  $a \le t \le b$ .

The general linear system of three first-order differential equations in three unknown functions x, y, and z is of the form

$$a_{1}(t)\frac{dx}{dt} + a_{2}(t)\frac{dy}{dt} + a_{3}(t)\frac{dz}{dt} + a_{4}(t)x + a_{5}(t)y + a_{6}(t)z = F_{1}(t),$$

$$b_{1}(t)\frac{dx}{dt} + b_{2}(t)\frac{dy}{dt} + b_{3}(t)\frac{dz}{dt} + b_{4}(t)x + b_{5}(t)y + b_{6}(t)z = F_{2}(t),$$

$$c_{1}(t)\frac{dx}{dt} + c_{2}(t)\frac{dy}{dt} + c_{3}(t)\frac{dz}{dt} + c_{4}(t)x + c_{5}(t)y + c_{6}(t)z = F_{3}(t).$$
(7.2)

As in the case of systems of the form (7.1), so also in this case we shall be concerned with systems that have constant coefficients. An example of such a system is

$$\frac{dx}{dt} + \frac{dy}{dt} - 2\frac{dz}{dt} + 2x - 3y + z = t,$$

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

$$\frac{dx}{dt} + \frac{dy}{dt} - 2\frac{dz}{dt} + 2x - 3y + z = t,$$

$$2\frac{dx}{dt} - \frac{dy}{dt} + 3\frac{dz}{dt} + x + 4y - 5z = \sin t,$$

$$\frac{dx}{dt} + 2\frac{dy}{dt} + \frac{dz}{dt} - 3x + 2y - z = \cos t$$

We shall say that a solution of system (7.2) is an ordered triple of real functions (f, g, h) such that x = f(t), y = g(t), z = h(t) simultaneously satisfy all three equations of the system (7.2) on some real interval  $a \le t \le b$ .

Systems of the form (7.1) and (7.2) contained only first derivatives, and we now consider the basic linear system involving higher derivatives. This is the general linear system of two second-order differential equations in two unknown functions x and y, and is a system of the form

$$a_{1}(t)\frac{d^{2}x}{dt^{2}} + a_{2}(t)\frac{d^{2}y}{dt^{2}} + a_{3}(t)\frac{dx}{dt} + a_{4}(t)\frac{dy}{dt} + a_{5}(t)x + a_{6}(t)y = F_{1}(t),$$
  

$$b_{1}(t)\frac{d^{2}x}{dt^{2}} + b_{2}(t)\frac{d^{2}y}{dt^{2}} + b_{3}(t)\frac{dx}{dt} + b_{4}(t)\frac{dy}{dt} + b_{5}(t)x + b_{6}(t)y = F_{2}(t).$$
(7.3)

We shall be concerned with systems having constant coefficients in this case also, and an example is provided by

$$2\frac{d^2x}{dt^2} + 5\frac{d^2y}{dt^2} + 7\frac{dx}{dt} + 3\frac{dy}{dt} + 2y = 3t + 1,$$
  
$$3\frac{d^2x}{dt^2} + 2\frac{d^2y}{dt^2} - 2\frac{dy}{dt} + 4x + y = 0.$$

For given fixed positive integers m and n, we could proceed, in like manner, to exhibit other general linear systems of n mth-order differential equations in n unknown functions and give examples of each such type of system. Instead we proceed to introduce the standard type of linear system referred to in the introductory paragraph at the start of the chapter, and of which we shall make a more systematic study later. We introduce this standard type as a special case of the system (7.1) of two first-order differential equations in two unknowns functions x and y.

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

We consider the special type of linear system (7.1), which is of the form

$$\frac{dx}{dt} = a_{11}(t)x + a_{12}(t)y + F_1(t),$$

$$\frac{dy}{dt} = a_{21}(t)x + a_{22}(t)y + F_2(t).$$
(7.4)

This is the so-called *normal form* in the case of two linear differential equations in two unknown functions. The characteristic feature of such a system is apparent from the manner in which the derivatives appear in it. An example of such a system with variable coefficients is

$$\frac{dx}{dt} = t^2 x + (t+1)y + t^3,$$
$$\frac{dy}{dt} = te^t x + t^3 y - e^t,$$

while one with constant coefficients is

$$\frac{dx}{dt} = 5x + 7y + t^2,$$
$$\frac{dy}{dt} = 2x - 3y + 2t.$$

The normal form in the case of a linear system of three differential equations in three unknown functions x, y, and z is

$$\frac{dx}{dt} = a_{11}(t)x + a_{12}(t)y + a_{13}(t)z + F_1(t),$$
  
$$\frac{dy}{dt} = a_{21}(t)x + a_{22}(t)y + a_{23}(t)z + F_2(t),$$
  
$$\frac{dz}{dt} = a_{31}(t)x + a_{32}(t)y + a_{33}(t)z + F_3(t).$$

An example of such a system is the constant coefficient system

$$\frac{dx}{dt} = 3x + 2y + z + t,$$
$$\frac{dy}{dt} = 2x - 4y + 5z - t^2,$$

Prepared by K.Aarthiya, Asst Prof, Department of Mathematics, KAHE

Page 7/12

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

$$\frac{dz}{dt} = 4x + y - 3z + 2t + 1.$$

# **B. Differential Operators**

In this section we shall present a symbolic operator method for solving linear systems with constant coefficients. This method depends upon the use of so-called *differential operators*, which we now introduce.

Let x be an n-times differentiable function of the independent variable t. We denote the operation of differentiation with respect to t by the symbol D and call D a differential operator. In terms of this differential operator the derivative dx/dt is denoted by Dx. That is,

$$Dx \equiv dx/dt.$$

In like manner, we denote the second derivative of x with respect to t by  $D^2x$ . Extending this, we denote the *n*th derivative of x with respect to t by  $D^nx$ . That is,

$$D^n x = \frac{d^n x}{dt^n} \qquad (n = 1, 2, \ldots).$$

Further extending this operator notation, we write

$$(D+c)x$$
 to denote  $\frac{dx}{dt} + cx$ 

and

$$(aD^n + bD^m)x$$
 to denote  $a\frac{d^nx}{dt^n} + b\frac{d^mx}{dt^m}$ 

where a, b, and c are constants.

In this notation the general linear differential expression with constant coefficients  $a_0, a_1, \ldots, a_{n-1}, a_n$ ,

$$a_0\frac{d^nx}{dt^n}+a_1\frac{d^{n-1}x}{dt^{n-1}}+\cdots+a_{n-1}\frac{dx}{dt}+a_nx,$$

is written

$$(a_0D^n + a_1D^{n-1} + \cdots + a_{n-1}D + a_n)x.$$

Observe carefully that the operators  $D^n$ ,  $D^{n-1}$ ,..., D in this expression do not represent quantities that are to be multiplied by the function x, but rather they indicate operations (differentiations) that are to be carried out upon this function. The expression

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

$$a_0D^n + a_1D^{n-1} + \cdots + a_{n-1}D + a_n$$

by itself, where  $a_0, a_1, \ldots, a_{n-1}, a_n$  are constants, is called a linear differential operator with constant coefficients.

# Example 7.1

Consider the linear differential operator

$$3D^2 + 5D - 2$$
.

If x is a twice differentiable function of t, then

$$(3D^2 + 5D - 2)x$$
 denotes  $3\frac{d^2x}{dt^2} + 5\frac{dx}{dt} - 2x$ .

For example, if  $x = t^3$ , we have

$$(3D^{2} + 5D - 2)t^{3} = 3\frac{d^{2}}{dt^{2}}(t^{3}) + 5\frac{d}{dt}(t^{3}) - 2(t^{3}) = 18t + 15t^{2} - 2t^{3}.$$

We shall now discuss certain useful properties of the linear differential operator with constant coefficients. In order to facilitate our discussion, we shall let L denote this operator. That is,

$$L \equiv a_0 D^n + a_1 D^{n-1} + \dots + a_{n-1} D + a_n,$$

where  $a_0, a_1, \ldots, a_{n-1}, a_n$  are constants. Now suppose that  $f_1$  and  $f_2$  are both *n*-times differentiable functions of t and  $c_1$  and  $c_2$  are constants. Then it can be shown that

$$L[c_1f_1 + c_2f_2] = c_1L[f_1] + c_2L[f_2].$$

For example, if the operator  $L \equiv 3D^2 + 5D - 2$  is applied to  $3t^2 + 2 \sin t$ , then

$$L[3t^{2} + 2\sin t] = 3L[t^{2}] + 2L[\sin t]$$

or

$$(3D^2 + 5D - 2)(3t^2 + 2\sin t) = 3(3D^2 + 5D - 2)t^2 + 2(3D^2 + 5D - 2)\sin t.$$

Now let

$$L_1 \equiv a_0 D^m + a_1 D^{m-1} + \dots + a_{m-1} D + a_m$$

and

$$L_2 \equiv b_0 D^n + b_1 D^{n-1} + \dots + b_{n-1} D + b_n$$

be two linear differential operators with constant coefficients  $a_0, a_1, \ldots, a_{m-1}, a_m$ , and  $b_0, b_1, \ldots, b_{n-1}, b_n$ , respectively. Let

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

$$L_1(r) \equiv a_0 r^m + a_1 r^{m-1} + \dots + a_{m-1} r + a_m$$

and

$$L_{2}(r) \equiv b_{0}r^{n} + b_{1}r^{n-1} + \dots + b_{n-1}r + b_{n}$$

be the two polynomials in the quantity r obtained from the operators  $L_1$  and  $L_2$ , respectively, by formally replacing D by r,  $D^2$  by  $r^2, \ldots, D^k$  by  $r^k$ . Let us denote the product of the polynomials  $L_1(r)$  and  $L_2(r)$  by L(r), that is,

$$L(r) = L_1(r)L_2(r).$$

Then, if f is a function possessing n + m derivatives, it can be shown that

$$L_1 L_2 f = L_2 L_1 f = L f, (7.10)$$

where L is the operator obtained from the "product polynomial" L(r) by formally replacing r by  $D, r^2$  by  $D^2, \ldots, r^{m+n}$  by  $D^{m+n}$ . Equation (7.10) indicates two important properties of linear differential operators with constant coefficients. First, it states the effect of first operating on f by  $L_2$  and then operating on the resulting function by  $L_1$  is the same as that which results from first operating on f by  $L_1$  and then operating on

this resulting function by  $L_2$ . Second, Equation (7.10) states that the effect of first operating on f by either  $L_1$  or  $L_2$  and then operating on the resulting function by the other is the same as that which results from operating on f by the "product operator" L.

# C. An Operator Method for Linear Systems with Constant Coefficients

We now proceed to explain a symbolic operator method for solving linear systems with constant coefficients. We shall outline the procedure of this method on a strictly formal basis and shall make no attempt to justify it.

We consider a linear system of the form

$$L_{1}x + L_{2}y = f_{1}(t),$$
  

$$L_{3}x + L_{4}y = f_{2}(t),$$
(7.11)

where  $L_1$ ,  $L_2$ ,  $L_3$ , and  $L_4$  are linear differential operators with constant coefficients.

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

That is,  $L_1$ ,  $L_2$ ,  $L_3$ , and  $L_4$  are operators of the forms

$$L_{1} \equiv a_{0}D^{m} + a_{1}D^{m-1} + \dots + a_{m-1}D + a_{m},$$

$$L_{2} \equiv b_{0}D^{n} + b_{1}D^{n-1} + \dots + b_{n-1}D + b_{n},$$

$$L_{3} \equiv \alpha_{0}D^{p} + \alpha_{1}D^{p-1} + \dots + \alpha_{p-1}D + \alpha_{p},$$

$$L_{4} \equiv \beta_{0}D^{q} + \beta_{1}D^{q-1} + \dots + \beta_{q-1}D + \beta_{q},$$

where the *a*'s, *b*'s,  $\alpha$ 's, and  $\beta$ 's are constants.

A simple example of a system which may be expressed in the form (7.11) is provided by

$$2\frac{dx}{dt} - 2\frac{dy}{dt} - 3x = t,$$
$$2\frac{dx}{dt} + 2\frac{dy}{dt} + 3x + 8y = 2.$$

Introducing operator notation this system takes the form

$$(2D-3)x - 2Dy = t,$$
  
 $(2D+3)x + (2D+8)y = 2.$ 

This is clearly of the form (7.11), where  $L_1 \equiv 2D - 3$ ,  $L_2 \equiv -2D$ ,  $L_3 \equiv 2D + 3$ , and  $L_4 \equiv 2D + 8$ .

Returning now to the general system (7.11), we apply the operator  $L_4$  to the first equation of (7.11) and the operator  $L_2$  to the second equation of (7.11), obtaining

$$L_4L_1x + L_4L_2y = L_4f_1,$$
  
$$L_2L_3x + L_2L_4y = L_2f_2.$$

We now subtract the second of these equations from the first. Since  $L_4L_2y = L_2L_4y$ , we obtain

$$L_4 L_1 x - L_2 L_3 x = L_4 f_1 - L_2 f_2,$$

or

$$(L_1L_4 - L_2L_3)x = L_4f_1 - L_2f_2.$$
(7.12)

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: IV BATCH-2017-2020

The expression  $L_1L_4 - L_2L_3$  in the left member of this equation is itself a linear differential operator with constant coefficients. We assume that it is neither zero nor a nonzero constant and denote it by  $L_5$ . If we further assume that the functions  $f_1$  and  $f_2$  are such that the right member  $L_4f_1 - L_2f_2$  of (7.12) exists, then this member is some function, say  $g_1$ , of t. Then Equation (7.12) may be written

$$L_5 x = g_1.$$
 (7.13)

Equation (7.13) is a linear differential equation with constant coefficients in the single dependent variable x. We thus observe that our procedure has eliminated the other dependent variable y. We now solve the differential equation (7.13) for x using the methods developed in Chapter 4. Suppose Equation (7.13) is of order N. Then the general solution of (7.13) is of the form

$$x = c_1 u_1 + c_2 u_2 + \dots + c_N u_N + U_1, \tag{7.14}$$

where  $u_1, u_2, ..., u_N$  are N linearly independent solutions of the homogeneous linear equation  $L_5 x = 0, c_1, c_2, ..., c_N$  are arbitrary constants, and  $U_1$  is a particular solution of  $L_5 x = g_1$ .

We again return to the system (7.11) and this time apply the operators  $L_3$  and  $L_1$  to the first and second equations, respectively, of the system. We obtain

$$L_3L_1x + L_3L_2y = L_3f_1,$$
  
$$L_1L_3x + L_1L_4y = L_1f_2.$$

Subtracting the first of these from the second, we obtain

$$(L_1L_4 - L_2L_3)y = L_1f_2 - L_3f_1.$$

Assuming that  $f_1$  and  $f_2$  are such that the right member  $L_1f_2 - L_3f_1$  of this equation exists, we may express it as some function, say  $g_2$ , of t. Then this equation may be written

$$L_5 y = g_2, (7.15)$$

where  $L_5$  denotes the operator  $L_1L_4 - L_2L_3$ . Equation (7.15) is a linear differential equation with constant coefficients in the single dependent variable y. This time we have eliminated the dependent variable x. Solving the differential equation (7.15) for y, we obtain its general solution in the form

$$y = k_1 u_1 + k_2 u_2 + \dots + k_N u_N + U_2, \qquad (7.16)$$

where  $u_1, u_2, ..., u_N$  are the N linearly independent solutions of  $L_5 y = 0$  (or  $L_5 x = 0$ ) that already appear in (7.14),  $k_1, k_2, ..., k_N$  are arbitrary constants, and  $U_2$  is a particular solution of  $L_5 y = g_2$ .

QUESTION	OPT 1	OPT 2	OPT 3	OPT 4	ANSWER
The second order partial differential equation is said to be if B <sup>2</sup> -4AC>0	parabolic	hyperbolic	elliptic	none	hyperbolic
The second order partial differential equation is said to be if B <sup>2</sup> -4AC=0	parabolic	hyperbolic	elliptic	none	parabplic
The second order partial differential equation is said to be if B <sup>2</sup> -4AC<0	parabolic	hyperbolic	elliptic	none	elliptic
The second order partial differential equation is said to be elliptic if $B^2$ -4AC=	>0	<0	0	>=0	<0
The second order partial differential equation is said to be paraolic if $B^2$ -4AC=	>0	<0	0	>=0	0
The second order partial differential equation is said to be hyperolic if $B^2$ -4AC=	>0	<0	0	>=0	>0
The second order linear partial differential equation, the coefficients areconstants	real	imaginary	known	unknown	real
The second order linear partial differential equation, the are real constants	coefficients	constants	numbers	operators	coefficients
The order linear partial differential equation, the coefficients are real constants	first	second	third	fourth	second
The second order linear partial differential equation, the coefficients are real	coefficients	constants	numbers	operators	constants
The simplex form by making a change in the independent variable in second order linear partial differential equation is called	canonical	parabola	hyperbola	elliptic	canonical
The simplex form by making a in the independent variable in second order linear partial differential equation is called canonical	change	parabola	hyperbola	elliptic	change
The form by making a change in the independent variable in second order linear partial differential equation is called canonical	simplex	parabola	hyperbola	elliptic	simplex
The simplex form by making a change in the variable in second order linear partial differential equation is called canonical	independent	parabola	hyperbola	elliptic	independent
The simplex form by making a change in the independent variable in second order linear differential equation is called canonical	partial	parabola	hyperbola	elliptic	partial
The simplex form by making a change in the independent variable in order linear partial differential equation is called canonical	second	first	second	third	second
The set of five functions is called The set of functions is called strip The set of five is called strip The set of five functions is called strip	strip five functions set	five strip five five	functions functions strip functions	set set strip	strip five functions set
The infinite sector is called the range of influence of the point.	Р	Q	R	S	R
The theorem can be applied with continuous data by using polynomial approximations only if a small variation in the initial data leads to a small change in the solution.	Cauchy Kowalewskay a	Cauchy	existence	uniqueness	Cauchy Kowalewskay a
The necessary condition for obtaining the second derivatives is that the curve must not be a characteristic curve.	Ll	R0	P0	LO	L0
The equation A $(dy/dx)^2$ -B $(dy/dx)$ +C=0 is called the	characteristic curve	characteristic equation	finite equation	finite curve	characteristic equation
The solution of the is a surface called an integral surface in the (x, y, u) space passing through L.	Cauchy Kowalewskay a	Cauchy problem	existence	uniqueness	Cauchy problem
The function $f(\lambda)$ and $g(\lambda)$ in $u = f(\lambda)$ and $\partial u/\partial n = g(\lambda)$ are called data	initial	final	cauchy	complete	cauchy

The Cauchy problem involves second order partial differential equations in independent variables.	0	1	2	3	2
The Cauchy Kowalewskaya theorem can be applied with data by using polynomial approximations only if a small variation in the initial data leads to a small change in the solution.	discontinuous	continuous	infinite	finite	continuous
The necessary condition for obtaining the derivatives is that the curve L0 must not be a characteristic curve.	first	second	third	fourth	second
The equation is called the characteristic equation.	A $(dy/dx)^2+B(dy/dx)+C=0$	A $(dy/dx)^2$ - B(dy/dx)-C=0	$A (dy/dx)^{2}-B(dy/dx)=0$	$A (dy/dx)^{2}-B(dy/dx)+C=0$	$A (dy/dx)^{2}-B(dy/dx)+C=0$
The solution of the Cauchy problem is a surface called an surface in the (x, y, u) space passing through L.	Complete	integral	bounded	unbounded	integral
The function in $u = f(\lambda)$ and $\partial u/\partial n = g(\lambda)$ are called cauchy data.	f(λ)	g(λ)	both $f(\lambda)$ and $g(\lambda)$	neither $f(\lambda)$ nor $g(\lambda)$	both $f(\lambda)$ and $g(\lambda)$
Linear ordinary differential equations are further classified according to the nature of the coefficients of thevariables and its derivatives.	- single	dependent	independent	constant	dependent
The standard form of first order differential equations derivative form is	(dy/dx)=f(x)	(dx/dy)=f(x,y	(dy/dy)=f(x,y	(dx/dy)=f(y)	(dy/dy)=f(x,y
The general solution of equation is called the complementary function of equation.	single	homogeneous	nonsingular	nonhomogene ous	homogeneous
Polynomial ar2+br+c=0 is called	characteristic polynomial	trivial polynomial	determinant polynomial	singular polynomial	characteristi
A non linear ordinary differential equation is an ordinary differential equation that is not	differential	integral	linear	non linear	linear
General solution of higher order linear differential equation depends on	arbitrary constant	coefficient	type of roots	method to which solved	type of roots
An equation of the form is called an equation with variables separable or simply a separable equations.	F(x)G(y) $dx+f(x)g(y)$ $dy=0$	G(y) dx+ g(y) dy=0	F(x)G(y) dx-g(y) dy=0	F(x)G(y) dx=0	$\begin{array}{l} F(x)G(y)\\ dx+f(x)g(y)\\ dy=0 \end{array}$
Any linear combination of solutions of the homogeneous linear differential equation is also a of homogeneous equation.	value	separable	solution	exact	solution

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: V BATCH-2017-2020

# UNIT V SYLLABUS

Basic Theory of linear systems in normal form : Homogeneous linear systems with constant coefficients -Two Equations in two unknown functions -The method of successive approximations -The Euler method-The modified – Euler method - The Runge-Kutta method.

# BASIC THEORY OF LINEAR SYSTEMS IN NORMAL FORM: TWO EQUATIONS IN TWO UNKNOWN FUNCTIONS

# A. Introduction

We shall begin by considering a basic type of system of two linear differential equations in two unknown functions. This system is of the form

$$\frac{dx}{dt} = a_{11}(t)x + a_{12}(t)y + F_1(t),$$

$$\frac{dy}{dt} = a_{21}(t)x + a_{22}(t)y + F_2(t).$$
(7.61)

We shall assume that the functions  $a_{11}$ ,  $a_{12}$ ,  $F_1$ ,  $a_{21}$ ,  $a_{22}$ , and  $F_2$  are all continuous on a real interval  $a \le t \le b$ . If  $F_1(t)$  and  $F_2(t)$  are zero for all t, then the system (7.61) is called *homogeneous*; otherwise, the system is said to be *nonhomogeneous*.

# Example 7.8

The system

$$\frac{dx}{dt} = 2x - y,$$

$$\frac{dy}{dt} = 3x + 6y,$$
(7.62)

is homogeneous; the system

# CLASS: II B.Sc.COURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: VBATCH-2017-2020

$$\frac{dx}{dt} = 2x - y - 5t,$$

$$\frac{dy}{dt} = 3x + 6y - 4,$$
(7.63)

is nonhomogeneous.

#### DEFINITION

By a solution of the system (7.61) we shall mean an ordered pair of real functions

(*f*, *g*),

each having a continuous derivative on the real interval  $a \le t \le b$ , such that

$$\frac{df(t)}{dt} = a_{11}(t)f(t) + a_{12}(t)g(t) + F_1(t),$$
$$\frac{dg(t)}{dt} = a_{21}(t)f(t) + a_{22}(t)g(t) + F_2(t),$$

for all t such that  $a \leq t \leq b$ . In other words,

$$\begin{aligned} \mathbf{x} &= f(t),\\ \mathbf{y} &= g(t), \end{aligned} \tag{7.65}$$

simultaneously satisfy both equations of the system (7.61) identically for  $a \le t \le b$ .

Notation. We shall use the notation

$$x = f(t),$$
  

$$y = g(t),$$
(7.65)

to denote a solution of the system (7.61) and shall speak of "the solution

 $\begin{aligned} x &= f(t), \\ y &= g(t). \end{aligned}$ 

Whenever we do this, we must remember that the solution thus referred to is really the ordered pair of functions (f, g) such that (7.65) simultaneously satisfy both equations of the system (7.61) identically on  $a \le t \le b$ .

# CLASS: II B.Sc.COURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: VBATCH-2017-2020

# Example 7.9

The ordered pair of functions defined for all t by  $(e^{5t}, -3e^{5t})$ , which we denote by

$$x = e^{5t}, 
 y = -3e^{5t}.$$
(7.66)

is a solution of the system (7.62). That is,

$$x = e^{5t},$$
  
 $y = -3e^{5t},$ 
(7.66)

simultaneously satisfy both equations of the system (7.62). Let us verify this by directly substituting (7.66) into (7.62). We have

$$\frac{d}{dt}(e^{5t}) = 2(e^{5t}) - (-3e^{5t}),$$
  
$$\frac{d}{dt}(-3e^{5t}) = 3(e^{5t}) + 6(-3e^{5t}),$$
  
$$5e^{5t} = 2e^{5t} + 3e^{5t},$$
  
$$-15e^{5t} = 3e^{5t} - 18e^{5t}.$$

or

Hence (7.66) is indeed a solution of the system (7.62). The reader should verify that the ordered pair of functions defined for all t by  $(e^{3t}, -e^{3t})$ , which we denote by

$$x = e^{3t},$$
  
$$y = -e^{3t},$$

is also a solution of the system (7.62).

We shall now proceed to survey the basic theory of linear systems. We shall observe a close analogy between this theory and that introduced in Section 4.1 for the single linear equation of higher order. Theorem 7.1 is the basic existence theorem dealing with the system (7.61).

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: V BATCH-2017-2020

# THEOREM 7.1

**Hypothesis.** Let the functions  $a_{11}$ ,  $a_{12}$ ,  $F_1$ ,  $a_{21}$ ,  $a_{22}$ , and  $F_2$  in the system (7.61) all be continuous on the interval  $a \le t \le b$ . Let  $t_0$  be any point of the interval  $a \le t \le b$ ; and let  $c_1$  and  $c_2$  be two arbitrary constants.

**Conclusion.** There exists a unique solution

 $\begin{aligned} x &= f(t), \\ y &= g(t), \end{aligned}$ 

of the system (7.61) such that

$$f(t_0) = c_1$$
 and  $g(t_0) = c_2$ ,

and this solution is defined on the entire interval  $a \leq t \leq b$ .

## **B. Homogeneous Linear Systems**

We shall now assume that  $F_1(t)$  and  $F_2(t)$  in the system (7.61) are both zero for all t and consider the basic theory of the resulting homogeneous linear system

$$\frac{dx}{dt} = a_{11}(t)x + a_{12}(t)y,$$

$$\frac{dy}{dt} = a_{21}(t)x + a_{22}(t)y$$

THEOREM 7.2

Hypothesis. Let

$$x = f_1(t),$$
  $x = f_2(t),$   
and  $y = g_1(t),$   $y = g_2(t),$  (7.68)

be two solutions of the homogeneous linear system (7.67). Let  $c_1$  and  $c_2$  be two arbitrary constants.

Conclusion. Then

$$\begin{aligned} x &= c_1 f_1(t) + c_2 f_2(t), \\ y &= c_1 g_1(t) + c_2 g_2(t), \end{aligned}$$
 (7.69)

is also a solution of the system (7.67).

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: V BATCH-2017-2020

# DEFINITION

The solution (7.69) is called a linear combination of the solutions (7.68). This definition enables us to express Theorem 7.2 in the following alternative form.

# THEOREM 7.2 RESTATED

Any linear combination of two solutions of the homogeneous linear system (7.67) is itself a solution of the system (7.67).

# Example 7.11

We have already observed that

$$x = e^{5t}$$
,  $x = e^{3t}$ ,  
and  
 $y = -3e^{5t}$ ,  $y = -e^{3t}$ ,

are solutions of the homogeneous linear system (7.62). Theorem 7.2 tells us that

$$x = c_1 e^{5t} + c_2 e^{3t},$$
  
$$y = -3c_1 e^{5t} - c_2 e^{3t},$$

where  $c_1$  and  $c_2$  are arbitrary constants, is also a solution of the system (7.62). For

example, if  $c_1 = 4$  and  $c_2 = -2$ , we have the solution

$$x = 4e^{5t} - 2e^{3t},$$
  
$$y = -12e^{5t} + 2e^{3t}.$$

# DEFINITION

Let

$$x = f_1(t), \qquad x = f_2(t),$$
  
and  
$$y = g_1(t), \qquad y = g_2(t),$$

# CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: V BATCH-2017-2020

be two solutions of the homogeneous linear system (7.67). These two solutions are linearly dependent on the interval  $a \le t \le b$  if there exist constants  $c_1$  and  $c_2$ , not both zero, such that

$$c_1 f_1(t) + c_2 f_2(t) = 0,$$
  

$$c_1 g_1(t) + c_2 g_2(t) = 0,$$
(7.70)

for all t such that  $a \leq t \leq b$ .

# DEFINITION

Let

$$\begin{aligned} x &= f_1(t), & x &= f_2(t), \\ and & \\ y &= g_1(t), & y &= g_2(t), \end{aligned}$$

be two solutions of the homogeneous linear system (7.67). These two solutions are linearly independent on  $a \le t \le b$  if they are not linearly dependent on  $a \le t \le b$ . That is, the solutions  $x = f_1(t)$ ,  $y = g_1(t)$  and  $x = f_2(t)$ ,  $y = g_2(t)$  are linearly independent on  $a \le t \le b$  if

$$c_1 f_1(t) + c_2 f_2(t) = 0,$$
  

$$c_1 g_1(t) + c_2 g_2(t) = 0,$$
(7.71)

for all t such that  $a \leq t \leq b$  implies that

$$c_1 = c_2 = 0.$$

# Example 7.12

The solutions

$$x = e^{5t}$$
,  $x = 2e^{5t}$ ,  
and  
 $y = -3e^{5t}$ ,  $y = -6e^{5t}$ ,

of the system (7.62) are linearly dependent on every interval  $a \le t \le b$ . For in this case

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: V BATCH-2017-2020

the conditions (7.70) become

$$c_1 e^{5t} + 2c_2 e^{5t} = 0,$$
  
-3c\_1 e^{5t} - 6c\_2 e^{5t} = 0, (7.72)

and clearly there exist constants  $c_1$  and  $c_2$ , not both zero, such that the conditions (7.72) hold on  $a \le t \le b$ . For example, let  $c_1 = 2$  and  $c_2 = -1$ .

On the other hand, the solutions

$$x = e^{5t}, x = e^{3t},$$
  
and  
$$y = -3e^{5t}, y = -e^{3t},$$

of system (7.62) are linearly independent on  $a \le t \le b$ . For in this case the conditions (7.71) are

$$c_1 e^{5t} + c_2 e^{3t} = 0,$$
  
-3c\_1 e^{5t} - c\_2 e^{3t} = 0.

If these conditions hold for all t such that  $a \le t \le b$ , then we must have  $c_1 = c_2 = 0$ .

We now state the following basic theorem concerning sets of linearly independent solutions of the homogeneous linear system (7.67).

# **THEOREM 7.3**

There exist sets of two linearly independent solutions of the homogeneous linear system (7.67). Every solution of the system (7.67) can be written as a linear combination of any two linearly independent solutions of (7.67).

# Example 7.13

We have seen that

$$x = e^{5t}$$
,  $x = e^{3t}$ ,  
and  
 $y = -3e^{5t}$ ,  $y = -e^{3t}$ ,

constitute a pair of linearly independent solutions of the system (7.62). This illustrates the first part of Theorem 7.3. The second part of the theorem tells us that every solution of the system (7.62) can be written in the form

CLASS: II B.Sc. COURSE CODE: 17MMU402

# COURSE NAME: PDE AND SYSTEM OF ODE UNIT: V BATCH-2017-2020

$$x = c_1 e^{5t} + c_2 e^{3t},$$
  
$$y = -3c_1 e^{5t} - c_2 e^{3t}$$

where  $c_1$  and  $c_2$  are suitably chosen constants.

Recall that in Section 4.1 in connection with the single *n*th-order homogeneous linear differential equation, we defined the general solution of such an equation to be a linear combination of *n* linearly independent solutions. As a result of Theorems 7.2 and 7.3 we now give an analogous definition of general solution for the homogeneous linear system (7.67).

# DEFINITION

Let

$$x = f_1(t), \qquad x = f_2(t),$$
  
and  
$$y = g_1(t), \qquad y = g_2(t),$$

be two linearly independent solutions of the homogeneous linear system (7.67). Let  $c_1$  and  $c_2$  be two arbitrary constants. Then the solution

$$x = c_1 f_1(t) + c_2 f_2(t),$$
  

$$y = c_1 g_1(t) + c_2 g_2(t),$$

is called a general solution of the system (7.67).

# Example 7.14

Since

$$x = e^{5t}, \qquad x = e^{3t},$$
  
and  
$$y = -3e^{5t}, \qquad y = -e^{3t},$$

are linearly independent solutions of the system (7.62), we may write the general solution of (7.62) in the form

$$x = c_1 e^{5t} + c_2 e^{3t},$$
  
$$y = -3c_1 e^{5t} - c_2 e^{3t},$$

where  $c_1$  and  $c_2$  are arbitrary constants.

# CLASS: II B.Sc.COURSE NAME: PDE AND SYSTEM OF ODECOURSE CODE: 17MMU402UNIT: VBATCH-2017-2020

# DEFINITION

Let

$$x = f_1(t),$$
  $x = f_2(t),$   
and  
 $y = g_1(t),$   $y = g_2(t),$ 

be two solutions of the homogeneous linear system (7.67). The determinant

 $\begin{vmatrix} f_1(t) & f_2(t) \\ g_1(t) & g_2(t) \end{vmatrix}$  (7.73)

is called the Wronskian of these two solutions. We denote it by W(t). THEOREM 7.4

Two solutions

$$x = f_1(t),$$
  $x = f_2(t),$   
and  
 $y = g_1(t),$   $y = g_2(t),$ 

of the homogeneous linear system (7.67) are linearly independent on an interval  $a \le t \le b$  if and only if their Wronskian determinant

$$W(t) = \begin{vmatrix} f_1(t) & f_2(t) \\ g_1(t) & g_2(t) \end{vmatrix}$$
(7.73)

. . .

is different from zero for all t such that  $a \le t \le b$ .

THEOREM 7.5

Let W(t) be the Wronskian of two solutions of homogeneous linear system (7.67) on an interval  $a \le t \le b$ . Then either W(t) = 0 for all  $t \in [a, b]$  or W(t) = 0 for no  $t \in [a, b]$ .

QUESTION	OPT 1	OPT 2	OPT 3	OPT 4	ANSWER
A partial differential equation has independent variables	one or more	two or more	more than one	none of these	two or more
The general linear system on the interval of	a<=t<=b	a <t<b< td=""><td>a&gt;t&gt;b</td><td>none of these</td><td>a&lt;=t&lt;=b</td></t<b<>	a>t>b	none of these	a<=t<=b
The normal form of linear system of differential equation is the function of	x,y,z	x	у	Z	x,y,z
Which method is used for solving linear system with constant coefficient	symbolic operator	constant method	coefficient method	none of these	symbolic operator
A linear operator with	constant	constant	linear	non linear	constant
The method which depends upon the symbolic operator is called	linear operator	non linear operator	differential operator	none of these	differential operator
A equation involving of variables with respect to a one or more independent variable is called differential equation	one or more	one	two	three	one or more
The canonical form is hyperolic if	a=-c	a=0	a=c	a>0	a=-c
The canonical form is paraolic if	a=-c	a=0	a=c	a>0	a=0
The canonical form is elliptic if	a=-c	a=0	a=c	a>0	a=c
The required solution for the product will be in the form	u(x,y)=X(x)Y(x)	u(x,y)=X(x)/Y	u(x,y)=X(x)	u(x,y)=X(x)+	u(x,y)=X(x)Y(
The function $u(x, y, z, t)$ is used to represent the	y)	(y)	I (y)	1 (y)	y)
displacement at a time of a particle whose position at is (x,y,z)	rest	motion	object	moving	rest
The equation for conduction tells us that the rate ofin joules per second	melting	heat transfer	heat conduction	cooling	heat transfer
The heat equation is a consequence of law	fourier law	cauchy law	eulers law	conservation	fourier law
The heat equation is a consequence of fourier law of	constant	melting	heating	conduction	conduction
An is a function where domain is a set of function	operator	method	linear operator	non linear operator	operator
The equation for conduction tells us that the rate of heat transfer in	joules per second	kilowatts	meter	kilometer	joules per second
The equation of differentiation with respect to t is denoted by the symbol	А	В	С	D	D
The dependent variable is expressed in the separable form $u(x, y) = X(x) Y(y)$ .	u(y, z)	u(x', y')	u(x, y)	u(y', z')	u(x, y)
The derivative dx/dt is denoted by D is called as operator.	Dy differential	Dx logical	Dy' relational	Dx' boolean	Dx differential
The second derivative of x with respect to t is denoted by	$D^2y$	$D^2x$	$D^2y'$	$D^2x'$	$D^2x$
The dependent variable $u(x, y)$ is expressed in the separable form $u(x, y) =$ .	X(x) + Y(y)	X(x) - Y(y)	X(x) Y(y)	$X(x) \ / \ Y(y)$	X(x) Y(y)
in finding solutions of a large class of initial boundary value problems.	integration	separation	differentiation	induction	separation
In the functions $\omega_n$ represents the discrete spectrum of circular frequencies.	Euclid	Kernal	eigen	node	eigen
The n <sup>th</sup> derivative of x with respect to t is denoted by	$D^ny$	D <sup>n</sup> x	$D^ny'$	D <sup>n</sup> x'	D <sup>n</sup> x
The method of of variables is also known as the Fourier method or the method of eigen function expansion.	integration	separation	differentiatio	induction	separation
The string displays loops separated by the nodes.	1	2	n-1	n	n
In the eigen functions $\gamma_n = \omega_n/2\pi = nc/21$ represents the	angular	circular	rectangular	spherical	angular
The harmonic is called fundamental harmonic	n=0	n=1	n=2	n=3	n=1
The harmonics are called overtones	n < 1	n=1	n>1	n=0	n>1
The $\lambda_n$ are called the values of the problem.	euclid	kernal	eigen	node	eigen

The T* represents of a vibrating string.	constant tension	eigen	node	kernel	constant tension
In the method of integral transforms K is the of the function	kennel	k(x,y)	kernel	k(z)	kernel
If f and g are said to be compatible then it have solution	unique	different	linear	non linear	unique
The derivative of x with respect to t is denoted by $D^2x$	second	first	third	fourth	second
The second of x with respect to t is denoted by $D^2x$	derivative	non linear	linear	integral	derivative
The second derivative of with respect to t is denoted by $D^2x$	x,y,z	X	у	Z	X
The second derivative of x with respect to is denoted by $D^2x$	t	X	у	Z	t
The harmonic n=1 is called	fundamental harmonic	fundamental	harmonic	none of these	fundamental harmonic

	Reg. No
	(16MMU602B)
KARPAGAM ACADEMY OF	HIGHER EDUCATION
COIMBATO	RE-21
DEPARTMENT OF MA	ATHEMATICS
Sixth Seme	ster
PDE and Systems	s of ODE
I Internal Test - Dec	cember'2018
Date : 17.12.2018 (AN)	Time : 2 Hours
Class : III B. Sc Mathematics	Maximum: 50 Marks

# **PART - A (20 x 1 = 20 Marks)**

# Answer all the questions:

\_\_\_\_

1. A partial differential equation is the equation involving partial derivatives of one or more dependent variables with respect to \_\_\_\_\_\_ independent variable.

a) one most one		b) atleast one	
		1	

c) more than one d) more than two

2. A differential equation involving ordinary derivatives of one or more dependent variables with respect to a -----independent variable is called an ordinary differential equation.

a) 1 b) 2 c) 3 d) 4

3. The equation  $u_x u_{xx} + x u u_y = \sin y$  is \_\_\_\_\_

a) quasi linear b) non linear c) Cauchy d) boundary

4. The order of the equation  $u_{xx} - u_{yy} = 0$  is \_\_\_\_\_

a) 1 b) 2 c) 3 d) 4

5. The partial differential equation  $u_t - u_{xx} = 0$  with  $u(x, 0) = \sin x$ . The given condition is called \_\_\_\_\_

a) initial	b) boundary	c) Cauchy	d) linear
6. F(x, y, u, u,	$(x, u_{yy}) = 0$ is	orde	er PDE.
a) first	b) second	c) third	d) fourth
7. The equation	on $F(x, y, u, u_x, u_y) = 0$	is said to be	
a) linear quasi linear	b) non linear	c) quasi linea	d) non
8. Which one	of the following is the	homogeneous	equation?
a) xy + x = e d) ax +	b) nx + my = + by = 0	sinx c) yx =	= y +3x
9. The existen solution.	ce of the mathematica	l problem have	
a) one most or d) mor	ne b) atleast one re than two	c) mor	e than one
10. $L + M = N$ property.	A + L & LM = ML are	called	
a) associative d) clos	b) distributive	e c) con	nmutative
11. An operate	or is said to be linear if	f it satisfies	
a) $L(cu) = c L$ + $c_2u_2) = c_1L($	L(u) b) $L(u)u_1) + c_2 L(u_2)$	$(u_1 + u_2) = L(u_1)$	$+ L(u_2) c) L(c_1u_1)$
d) L( $c_1u_1 + c_2$ )	$u_2 \neq c_1 L(u_1) + c_2 L(u_2)$	)	

12. (L + M) + N = L + (M + N) is called \_\_\_\_\_ property.

a) associative b) distributive c) commutative d) closed

13. A non linear ordinary differential equation is an ordinary differential equation that is not .....

a) **linear** b) non linear c) differential d) integral

14. Linear ordinary differential equations are further classified according to the nature of the coefficients of the .....variables and its derivatives

a) single b) **dependent** c) independent d) constant

15. The order of .....derivatives involved in the differential equations is called order of the differential equation

a) Zero b) lowest c) **highest** d) infinite

16. The equation f(x, y, z, a, b) = 0 containing two arbitrary parameters is called \_\_\_\_\_\_ of an equation.

a) linear b) non linear c) partial solution d) complete solution

17. A solution which is not everywhere differentiable is called a \_\_\_\_\_\_ solution.

a) strong b) **weak** c) low d) high

18. A curve in (x, y, u)-space, whose tangent at every point coincides with the characteristic direction field (a, b, c), is called a

a) tangent curve characteristic curve	<ul><li>b) normal curve</li><li>d) uniform curve</li></ul>	c)				
19. The characteristic	direction is also c	alled as				
a) <b>Monge axis</b> b) Mon d) Monge line	a) <b>Monge axis</b> b) Monge cone c) Monge curve d) Monge line					
20. In the equation u(	$x_0(t), y_0(t)) = u_0(t),$	$u_0(t)$ is called the				
a) final data b) <b>init</b>	ial data c)	curve value d) null value				
I	PART - B (3 x 2 = 6 Marks)					
Answer all the questio 21. Define differen 22. Write the Lapla 23.	<ul> <li>Answer all the questions:</li> <li>21. Define differential equation.</li> <li>22. Write the Laplace equation.</li> <li>23.</li> <li>PART - C (3 x 8 = 24 Marks)</li> </ul>					
<b>Answer all the que</b> 24. a) Explain the c	estions: classification of first (OR)	order equations.				
b) Find the general solution of the first-order linear partial differential equation $xu_x + y u_y = u$ .						

25. a) Find a solution f of the differential equation  $\frac{dy}{dx} = 2x$ . Show that x = 1 is the solution f of the value.

b) Reduce the equation  $u_x - u_y = u$  to canonical form and obtain the general solution.

26. a) Show that the initial value problem  $\frac{dy}{dx} = -\frac{x}{y}$ , y(3) = 4. (OR)

b)

QUESTION	OPT 1	OPT 2	OPT 3	OPT 4	ANSWER
The system of the form $dx/dt = a_{11}(t)x + a_{12}(t)y + F_1(t)$ if F (t) is non-zero for all t then the system is called	homogeneous	non	linear	non linear	non
	nomogeneous	homogeneous	linear	non mea	homogeneous
Any of two solutions of the homogeneous linear system $dx/dt=a_{11}x + a_{12}y$ and $dy/dt=a_{21}x + a_{22}y$ is itself a solution.	different equations	non linear combination	linear combination	same equations	linear combination
The general solution of the of the form $x = c_1 e^{5t} + c_2 e^{3t}$ where $c_1$ and $c_2$ are	constant	variable	dependent	independent	constant
The homogeneous linear system $X' = a_1x + b_1y$ and $Y' = a_2x + b_2y$ where the coefficient $a_1$ , $b_1$ , $a_2$ , & $b_2$ are	variable	real constant	dependent	independent	real constant
The general solution of the system $dx/dt=a_1x_1 + b_1x_2$ is written as $x = A_1e^{(\lambda 1 t)} + A_2e^{(\lambda 2 t)}$ . Then roots $\lambda 1$ and $\lambda 2$ are	imaginary and unequal	conjugate complex	real and equal	real and distinct	real and distinct
The general solution of the system $dx/dt=a_1x_1 + b_1x_2$ is written as $x = (A_1 \cos bt + A_2 \sin bt)$ . Then roots $\lambda 1$ and $\lambda 2$ are	imaginary and unequal	conjugate complex	real and equal	real and distinct	conjugate complex
The general solution of the system $dx/dt=a_1x_1 + b_1x_2$ is written as $x = (A_1t + A_2) e^{\lambda t}$ then roots $\lambda 1$ and $\lambda 2$ are	imaginary and unequal	conjugate complex	real and equal	real and distinct	real and equal
Equations having a common solution are called	linear equations	homogeneous equations	simultaneous equations	non homogeneous	simultaneous equations
Complementary function of (D2 + 4)y = tan 200x is	(Acos2x + Bsin2x)	(Acos2x – Bsin2x)	(Acosh2x + Bsinh2x)	(Acosh2x – Bsinh2x)	(Acos2x + Bsin2x)
If roots of linear second order differential equation is real double root than general solution will contain	two constants & two exponentials	one constant & two exponentials	two constants & one exponential	d) one constant & one exponential	two constants & one exponential
A particular case of Runge Kutta method of second order is	Milne's method	Picard's method	Modified Euler method	Runge's method	Modified Euler method
Runge Kutta of first order is nothing but the	Euler method	modified Euler method	Taylor series	none of these	Euler method
In Runge Kutta second and fourth order methods, the values of $k_1$ and $k_2$ are	always positive	always negative	differ	same	same
Thevalues are calculated in Runge Kutta fourth order method.	$k_1, k_2, k_3, k_4$ and Dy	$k_1$ , $k_2$ and Dy	$k_1$ , $k_2$ , $k_3$ and Dy	k <sub>1</sub> and Dy	$k_1, k_2, k_3, k_4$ and Dy
The use of Runge kutta method gives to the solutions of the differential equation than Taylor's series method.	Slow convergence	quick convergence	oscillation	divergence	quick convergence
In Runge – kutta method the value x is taken as	h = x0 - x	x0=x+h	$\mathbf{x} = \mathbf{x}0 + \mathbf{h}$	$\mathbf{h} = \mathbf{x}0 + \mathbf{x}$	x=x0+h
The is nothing but the modified Euler method.	Taylor series method	Runge kutta method of fourth order	Runge kutta method of third order	Runge kutta method of second order	Runge kutta method of second order
If dy/dx is a function x alone, then fourth order Runge – Kutta method reduces to	Euler method	Taylor series	Simpson method	Trapezoidal rule	Simpson method
In Runge Kutta methods, the derivatives of order are not require and we require only the given function values at different points.	lower	higher	middle	zero	higher
The formula of Dy in second order Runge Kutta method is given by	k1	k2	k3	k4	k2
Karpagan Academy Of Higher Education. Department Of Mathematics. Answer ky. CTA-I. Subject: PDE & System of ODE (17MMU402 & 16MMU402B) Class : II. B.Sc. Mathematics & II. B.Sc. Mathematics.

· 10 --

PART-A.

- 1. C) More than one
- 2. 0) 1
- 3. a) quas? linear
- 4. 6) 2
- 5. a) Initial
- 6. -Second
- 7. quasi lenear. (c)
- 8. d) ax+by=0
- 9. b) at least one
- 10. c) commutative
- 11. c)  $L(CC, u_1 + C_2 U_2) = C, L(u_1) + C_2 L(U_2)$
- 12. a) associative
- 13. a) finear
- 14. dependent
- 15. c) highest
- b. . complete solution
- 17. bob) weak
- 18. Charactertistic curve

- 19. a) Monge axis
- 20. D initial data.

### PART-B.

2D. Differential equation:-

A equation involving one or more dependent Variable with suspect to the one or more independent Variable 1  $E_{x}: \frac{d^{2}y}{dx^{2}} + \frac{dy}{dx} + y = 0.$ 

22. Laplace equation: -

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0.$$

23. Differential Operators:

$$L(u) = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}$$
  

$$M(u) = \frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial x} + x \frac{\partial u}{\partial y}.$$

PART-C.

24) (a) Classification of first order equation: (i) Linear differential equation (ii) Non Linear differential equation

(111) Quali linear differential equation  
(W) Homogeneous differential equation  
(V) Non Homogeneous differential equations.  
(D) Griven 
$$\chi U_{1}+YU_{2}=U$$
.  
We know that  $a(z_{1},y_{1},y_{2})U_{2}+b(x,y_{1},y_{2})U_{2}=U \rightarrow O$ .  
 $\therefore \phi = const = c, & \psi = const = c_{2}$ .  
 $\frac{d\chi}{a} = \frac{dy}{b} - \frac{du}{c} \rightarrow O$ .  
 $\frac{d\chi}{a} = \frac{dy}{b} - \frac{du}{c} \rightarrow O$ .  
 $\frac{d\chi}{a} = \frac{dy}{b} - \frac{du}{c} \rightarrow O$ .  
 $\frac{d\chi}{a} = \frac{dy}{b} + \frac{du}{c} + c\phi = 0$ .  
 $\frac{d\chi}{a} = \frac{dy}{b} + c\phi = 0$ .  
 $\frac{d\chi}{a} + b\phi + c\phi = 0 \rightarrow O$ .  
 $\therefore a\phi_{2} + b\phi_{2} + c\phi = 0 \rightarrow O$ .  
 $\therefore a\phi_{2} + b\phi_{2} + c\phi = 0 \rightarrow O$ .  
 $\therefore a\phi_{2} + b\phi_{2} + c\phi = 0 \rightarrow O$ .  
 $\frac{d\phi_{1}}{d(u, y)} = \frac{d}{\partial(c\phi_{1}, y)} = \frac{d(\phi_{1}, \psi)}{\partial(c_{1}, y)} = \frac{d(\phi_{1}, \psi)}{\partial(c_{1}$ 

25) (a). 
$$\frac{dy}{dz} = 22$$
 and  $z = 1$  at  $h$ .  
Given  $\frac{dy}{dz} = 22$   $\longrightarrow$  (b).  
 $\frac{dy}{dz} = 22$   
 $\Rightarrow y = x^2 + c - \sqrt{3}$ .  
From (c) in Subl-  $z = 1$  and  $y = h$  in (c).  
 $A = 1^2 + c$   
 $c = 3$ .  
 $\overline{f \cdot y = 2^2 + 3}$ .  
2b) w. Given  $\frac{dy}{dx} = -\frac{x}{y} \rightarrow 0$ .  
 $T : y(0) = h \rightarrow 0$ .  
 $T : y(0) = h \rightarrow 0$ .  
 $\frac{dy}{dx} = -\frac{2}{y}$ .  
 $y dy = -x dx$ .  
 $\int y dy = -x dx$ .  
 $\int (c = \pm 5] \rightarrow (b)$ .  
 $x^2 + y^2 = (\pm 5)^2$   
 $\Rightarrow y = \pm \sqrt{25 - x^2} \Rightarrow y = \sqrt{25 - x^2}$ .

Reg. No (16MMU602B/17MMU402) KARPAGAM ACADEMY OF HIGHER EDUCATION COIMBATORE-21 DEPARTMENT OF MATHEMATICS PDE and Systems of ODE II Internal Test – Feb' 2018 Date : 04.02.2019 (AN) Time : 2 Hours	<ul> <li>6. Z=X(x)Y(y) is called of variables a)integration b)separation c) differentiation d)induction</li> <li>7. The separation principle can readily be extended to number of variables. a)smaller b)unique c)larger d)contrary</li> <li>8. The partial differential equation xy dz/dx = 5 d<sup>2</sup>z/dy<sup>2</sup> is classified as</li> </ul>		
Class : III B. Sc. / II B.Sc.Maximum: 50 MarksPART - A (20 x 1 = 20 Marks)Answer all the questions:1. A solution to the partial differential equation $\frac{\partial^2 u}{\partial x^2} = 9 \frac{\partial^2 u}{\partial y^2}$ is	a) elliptic b) parabolic c) hyperbolic d) sphere 9. If $f(x,p) = g(y,q)$ is calledequation a) Clairaut b) Charpit c) Crout d) Separable 10. $L(z)+f(x,y,z,p,q)=0$ where L is the operator a) Laplace b) Differential		
a) $\cos(3x - y)$ c) $\sin(3x - y)$ 2. The partial differential equation $5\frac{\partial^2 z}{\partial x^2} + 6\frac{\partial^2 z}{\partial y^2} = xy$ is classified	a) Laplace b) Differential c) Lagrange d) Longdivision 11. The infinite sector is called the range of influence of the point. a)P b) Q c) R d) S		
<ul> <li>as</li> <li>a) elliptic</li> <li>b) parabolic</li> <li>c) hyperbolic</li> <li>d) sphere</li> </ul> 3. A partial differential equation is one which involves derivatives. <ul> <li>a) Single</li> <li>b) ordinary</li> <li>c) partial</li> <li>d) linear</li> </ul> 4. The order of PDE to be the order of the derivative of order occurring in it.	<ul> <li>12. Thetheorem can be applied with continuous data by using polynomial approximations only if a small variation in the initial data leads to a small change in the solution.</li> <li>a)Cauchy Kowalewskaya b) Cauchy</li> <li>c) Existence d) Uniqueness</li> <li>13. The necessary condition for obtaining the second derivatives is that the curve must not be a characteristic curve.</li> <li>a)L<sub>1</sub> b) R<sub>0</sub> c) P<sub>0</sub> d) L<sub>0</sub></li> </ul>		
<ul> <li>5. The variational approach to value problem is useful in the derivation of approximating solution</li> <li>a) Euclid b) Kernal c) eigen d) node</li> </ul>	14. The equation $A\left(\frac{dy}{dx}\right)^2 - B\left(\frac{dy}{dx}\right) + C = 0$ is called the a)characteristic curve b) characteristic equation c) finite equation d) finite curve		

- 15. The solution of the \_\_\_\_\_\_ is a surface called an integral surface in the (x, y, u) space passing through L.
  a)Cauchy Kowalewskaya
  b) Cauchy problem
  c) Existence
  d) Uniqueness
- 16. The function  $f(\lambda)$  and  $g(\lambda)$  in  $u = f(\lambda)$  and  $\frac{\partial u}{\partial n} = g(\lambda)$  are called data.
  - a)initial b) final c) cauchy d) complete
- 17. The Cauchy problem involves second order partial differential equations in----- independent variables.

a)Two b) one c) three d) zero

- 19. The necessary condition for obtaining the \_\_\_\_\_derivatives is that the curve  $L_0$  must not be a characteristic curve.

a)fourthb) thirdc) secondd) first20. The equation\_\_\_\_\_\_is called the characteristic equation.

a) 
$$A\left(\frac{dy}{dx}\right)^2 - B\left(\frac{dy}{dx}\right) + C = 0$$
  
b)  $A\left(\frac{dy}{dx}\right)^2 + B\left(\frac{dy}{dx}\right) + C = 0$   
c)  $A\left(\frac{dy}{dx}\right)^2 - B\left(\frac{dy}{dx}\right) - C = 0$   
d)  $A\left(\frac{dy}{dx}\right)^2 - B\left(\frac{dy}{dx}\right) = 0$ 

PART - B (3 x 2 = 6 Marks)

### Answer all the questions:

- 21. Define initial value problem.
- 22. Write the heat equation.

23. Write the general solution of initial boundary value problem using a semi-infinite string with a fixed end.

### **PART – C (3 x 8 = 24 Marks)**

### Answer all the questions:

24. a) Solve  $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u$  by method of separation of variables where  $u(x, 0) = 6e^{-3x}$ . (OR)

- b) Derive wave equation.
- 25. a)Use the separation of variables u(x, y) = f(x) + g(y) to solve the equation  $u_x^2 + u_y^2 = 1$ .

(**OR**)

b) Transform the equation  $y^2u_{xx} - x^2u_{yy} = 0$  into canonical form.

26. a) Derive the canonical form of the elliptic equation.

b) Transform the equation  $4u_{xx} + 5u_{xy} + u_{yy} + u_x + u_y = 2$  into canonical form.

Reg. No (16MMU602B/17MMU402) KARPAGAM ACADEMY OF HIGHER EDUCATION							
COIMBATORE-21 DEPARTMENT OF MATHEMATICS PDE and Systems of ODE III Internal Test March 2019							
Da Cl	ate : 11.03.2 ass : III B.So	019 (AN) c./II B.Sc.Mat	hematics	Time : 2 Hours Maximum: 50 Marks	9		
	PART - A (20 x 1 = 20 Marks)						
A	nswer all the	questions:					
1.	The equation symbol	of differentiatio	n with respect t	o t is denoted by the	1		
	a)A	b) B	c) C	d) D			
2.	The depender	nt variable	is express	sed in the separable form			
	$\mathbf{u}(\mathbf{x}, \mathbf{y}) = \mathbf{X}(\mathbf{x}) \mathbf{Y}(\mathbf{y}).$						
	a)u(y, z)	b) u(x', y')	c) u(x, y)	d) u(y', z')			
3.	The derivativ	e dx/dt is denote	ed by				
	a)Dy	b) Dx	c) Dy'	d) Dx'	1		
4.	4. D is called as operator.						
	a)differential		b) logical	b) logical			
	c) relational		d) Boolean				
5.	. The system of the form $\frac{dx}{dt} = a_{11}(t)x + a_{12}(t)y + F_1(t)$ if $F_1(t)$ is non zero						
	for all t, then the system is called						
	a)homogeneo	us	b) non homogeneous		1		
	c) linear		d) nonlinear				
6.	Any	of two solutions of the homogeneous linear system 1					
	$\frac{dx}{dt} = a_{11}x + a_{12}y$ and $\frac{dy}{dt} = a_{21}x + a_{22}y$ is itself a solution.						
	a)different eq	uations	b) non linear	b) non linear combination			
	c) linear com	bination	d) same equations				

7.	The general solution of the of the form $x = c_1e^{5t} + c_2e^{3t}$ where $c_1$ and $c_2$						
	are						
	a)constant		b) variable				
	c) dependent		d) independent				
8.	The homogeneo	The homogeneous linear system $X' = a_1x + b_1y$ and $Y' = a_2x + b_2y$ where					
	the coefficient $a_1$ , $b_1$ , $a_2$ , & $b_2$ are						
	a)variable		b) real constant				
	c) imaginary constant		d) zero				
Э.	The second derivative of x with respect to t is denoted by						
	a)D <sup>2</sup> y	b) $D^2x$	c) D <sup>2</sup> y'	d) D <sup>2</sup> x'			
10.	The dependent	variable u(x, y)	is expressed in t	he separable form			
	u(x,y) =						
	a)X(x) + Y(y)	b) X(x) - Y(y)	c) $X(x) Y(y)$	d) X(x) / Y(y)			
11.	1. The method of of variables is widely used in finding						
	solutions of a large class of initial boundary value problems.						
	a)integration b) separation						
	c) differentiation		d) induction				
12.	. In the functions $\omega_n$ represents the discrete spectrum of						
	circular frequen	icies.					
	a)Euclid	b) Kernal	c) eigen	d) node			
13.	The n <sup>th</sup> derivativ	ve of x with resp	ect to t is denote	d by			
	a)D <sup>n</sup> y	b) D <sup>n</sup> x	c) D <sup>n</sup> y'	d) D <sup>n</sup> x'			
14.	The method of of variables is also known as the Fourier						
	method or the method of eigen function expansion.						
	a)integration	ntegration b) separation					
	c) differentiati	on	d) induction				
15.	The string displ	The string displays loops separated by the nodes.					
	a)1	b) 2	c) n-1	d) n			
16.	In the eigen functions $\gamma_n = \frac{\omega_n}{2\pi} = \frac{nc}{21}$ represents thefrequencies.						
	a)angular	b) circular	c) rectangular	d) spherical			

17. The harmonics ----- are called overtones.

a)n < 1 b) n = 1 c) n > 1 d) n = 0

18. The  $\lambda_n$  are called the ------ values of the problem.

a)euclid b) kernal c) eigen d) node

19. The T<sup>\*</sup> represents ------ of a vibrating string.
a)constant tension
b) eigen
c) node
d) kernel
20. In the method of integral transforms K is the-----of the function.

a)kennal b) k(x,y) c) kernel d) k(z)

### PART - B (3 x 2 = 6 Marks)

### Answer all the questions:

- 21. Write the normal form of three linear differential equations in two unknown functions.
- 22. Write the linear differential operators with constant coefficients.
- 23. Define linearly independent solutions.

### **PART – C (3 x 8 = 24 Marks)**

### Answer all the questions:

### 24. a) Reduce the second order homogeneous partial differential equation to two ordinary differential using method of separation of variables.

### (**OR**)

b) Write a note on types of linear systems.

25. a) Explain the eigen value problem for wave equation.

(OR)  
b) Solve the system 
$$\frac{dx}{dt} + \frac{dy}{dt} - x = -2t$$
 and  $\frac{dx}{dt} + \frac{dy}{dt} - 3x - y = t^2$ 

26. a) Apply the Euler method to the initial value problem  $\frac{dy}{dx} = x - 2y$ , y(0)=1. Employ the method to approximate the value of the solution y at x = 0.1, 0.2, 0.3 and 0.4 using h = 0.1. Obtain results to three figures after the decimal point.

### (OR)

b) Find the general solution of the system of equations  $\frac{dx}{dt} = 4x - y, \frac{dy}{dt} = x + 2y.$ 

Karpagan Acadamy & Higher Education. Department & Mathematics. Answer key. CIA-TIL Sub: PDE & System of ODE (147MMU402 & 16MMU602B] Class: I.B.Sc. Mathematics & IV.B.Sc. Mathematics. PART-ITA.

- 1. d) D
- 2. c) u(x,y)
- 3) b) Dr
- A) a) differtantial
- 5) b) non homogeneous
- 6). c) finear combination.
- 7) a) constant
- 8) b) greal Constant
- 9 b) D2
- 10) C) X(x) Y(y)
- 11). b) Separation
- 12 c) eigen
- B) b) D'2
- 14) Separation
- 15). d) 1
- (b) a) anguhr.

- H). c) n>1.
- 18. c)eigen 19. a) constant tension

20. c) kernel.

PART-B.

 $\begin{aligned} \partial t &= Q_{11}(t) + Q_{12}(t) + Q_{13}(t) + F_{1}(t) \\ &= \frac{dy}{dt} = Q_{21}(t) + Q_{22}(t) + Q_{23}(t) + F_{2}(t) \\ &= \frac{dz}{dt} = Q_{21}(t) + Q_{22}(t) + Q_{23}(t) + F_{2}(t) \\ &= \frac{dz}{dt} = Q_{31}(t) + Q_{32}(t) + Q_{33}(t) + F_{3}(t) \\ &= \frac{dz}{dt} = Q_{31}(t) + Q_{32}(t) + Q_{33}(t) + Q_{33}(t) + F_{3}(t) \end{aligned}$ 

$$22. \quad a_0 D^n + a_1 D^{n-1} + \dots + a_{n-1} D + a_n = 0$$

28. Linear independent solution consist of the dependent Variable y and independent Variable x.  $\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} + 2x = 3.$ 

 $d_{4}$ ) (b). Types of linear System:  $a_1\left(\frac{dx}{dt}\right) + a_2\left(\frac{dy}{dt}\right) + a_3\left(x\right) + a_4(y) = F(t).$  $b_1\left(\frac{dx}{dt}\right) + b_2\left(\frac{dy}{dt}\right) + b_3(x) + b_4(y) = F_2(t).$ 

$$dx = a_1 x + a_2 y + a_3 z + F_1(t).$$

$$dy = b_1 x + b_2 y + b_3 z + F_2(t).$$

$$dz = c_1 x + c_2 y + c_3 z + F_3(t).$$

$$dz = c_1 x + c_2 y + c_3 z + F_3(t).$$

$$dz = c_1 x + c_2 y + c_3 z + F_3(t).$$

$$dx = x^{-1} \tau \quad x \quad u_{t-1} x = 0 \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad T(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad T(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad T(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad T(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad -\chi(t) \quad \longrightarrow \\ (x_1, t) = \chi(x) \quad -\chi(t) \quad -$$

Give 3: 
$$m = \pm \sqrt{-\lambda}$$
  
 $0 = B \sin \sqrt{-\lambda} I$ .  
 $u(x,t) = \left[ \cos \frac{n\pi}{2} ct + D \sin \frac{n\pi ct}{2} \right] B \sin \frac{n\pi x}{2}$   
 $= \sum_{n=1}^{\infty} \left[ a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} \left[ a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} \left[ a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct + b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct - b_n \sin \frac{n\pi}{2} ct \right]$   
 $a_n = \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{2} ct - b_n \sin \frac$ 

Reg. No.....

### [16MMU602B]

# KARPAGAM ACADEMY OF HIGHER EDUCATION

Pollachi Main Road, Eachanari Post, Coimbatore - 641 021 (For the candidates admitted from 2016 onwards) (Established Under Section 3 of UGC Act, 1956) (Deemed to be University)

## **B.Sc., DEGREE EXAMINATION, APRIL 2019**

Sixth Semester

MATHEMATICS

## PDE AND SYSTEMS OF ODE

Time: 3 hours

Maximum : 60 marks

(Question Nos. 1 to 20 Online Examinations) PART - A (20 x 1 = 20 Marks) (30 Minutes)

PART B (5 x 2 = 10 Marks) (2 ½ Hours) Answer ALL the Questions

Write the Laplace equation and Poisson equation.
 Define initial value problem.

23. Write the general solution of initial boundary value problem using a semi-infinite

string with a fixed end

25. Define linearly independent solutions. 24. Define fundamental harmonic.

### PART C (5 x 6 = 30 Marks) Answer ALL the Questions

26. a. Show that the family of the spheres  $(x-a)^2 + (y-b)^2 + z^2 = r^2$  satisfies the first order, non liner partial differential equation  $z^2(p^2 + q^2 + 1) = r^2$ .

b. If  $\phi = \phi(x, y, z)$  and  $\psi = \psi(x, y, z)$  are two given functions of x, y, and z and if  $f(\phi, \psi) = 0$ , then prove that z = z(x, y) satisfies a first-order , partial differential

equation  $p \frac{\partial(\phi, \psi)}{\partial(y, z)} + q \frac{\partial(\phi, \psi)}{\partial(z, x)} = \frac{\partial(\phi, \psi)}{\partial(x, y)}$ .

27. a. Solve the IVP  $u_x + 2u_y = 0$ , using  $u(0, y) = 4e^{-2y}$ 

b. Derive wave equation.

2

28. a. Transform the equation  $4u_{xx} + 5u_{xy} + u_{yy} + u_x + u_y = 2$  into canonical form. b. Find the solution of the initial boundary value problem using a semi-infinite string with a free end at x = 0. ç

29. a. Derive struck string problem.

b. Solve the system  $2\frac{dx}{dt} - 2\frac{dy}{dt} - 3x = t$  and  $2\frac{dx}{dt} + 2\frac{dy}{dt} + 3x + 8y = 2$ . ç

30. a. Find the general solution of the system of equations  $\frac{dx}{dt} = 4x - y \cdot \frac{dy}{dt} = x + 2y$ 

ç

b. Apply the Euler method to the initial value problem  $\frac{dy}{dx} = 2x + y$ , y(0) = 1. Use the method to approximate the value of the solution y at x = 0.2, 0.4 and 0.6

using h = 0.1. Obtain results to three figures after the decimal point.