

(Deemed to be University Established Under Section 3 of UGC Act 1956) Coimbatore – 641 021.

### LECTURE PLAN DEPARTMENT OF COMMERCE

STAFF NAME: Dr. S. Santhosh Kumar SUBJECT NAME: BUSINESS MATHEMATICS AND STAISTICS SUB.CODE:18BAU202 SEMESTER: II CLASS: I B.B.A.

S. No.	Lecture Duration Period	Topics to be Covered	Support Material/Page No.
		UNIT-I	
1	1	Introduction and definition of a matrix.	R4:Chap-15 Pg.No:280-281
2	1	Types of matrices	R4:Chap-15 Pg.No:281-284
3	1	Algebra of matrices	R4:Chap-15 Pg.No:285-286
4	1	Calculation of values of determinants up to third order;	R4:Chap-15 Pg.No:297-301
5	1	Tutorial-I	
6	1	Adjoint of a matrix	R4:Chap-15 Pg.No:306-307
7	1	Problems on finding inverse of a matrix through ad joint	R4:Chap-15 Pg.No:307-313
8	1	Continuation on problems on finding inverse of a matrix through ad joint	R6:Chap:1:Pg.No: 1.9-1.14
9	1	Applications of matrices to solution of simple business and economic problems	R6:Chap:4:Pg.No:4.1-4.2
10	1	Tutorial-II	
11	1	Simple and compound interest Rates of interest	R6:Chap:4:Pg.No:4.3-4.15
12	1	Continuation on Simple and compound interest Rates of interest	R6:Chap:4:Pg.No:4.32-4.35
13	1	Problems on Compounding and discounting of a sum using different types of rates	R6:Chap:4:Pg.No:4.13-4.22
14	1	Continuation of problems on Compounding and discounting of a sum using different types of rates	R6:Chap:4:Pg.No:4.22-4.31

15	1	Tutorial-III	
16	1	Recapitulation & discussion of possible questions	
	1	Fotal No of Hours Planned For Unit I=16	
		UNIT-II	
1	1	Introduction of Mathematical Functions	R6:Chap:2:Pg.No:2.3-2.4
2	1	Types of Mathematical Functions: linear, quadratic, polynomial formula and its problems	R6:Chap:2:Pg.No:2.4-2.5
3	1	Concept of limit of a function	R4:Chap:17:Pg.No:363-364
4	1	Concept of Continuity of a function	R4:Chap:17:Pg.No:364-365
5	1	Tutorial-I	
6	1	Concept of differentiation	R4:Chap:17:Pg.No:366-371
7	1	Problems on differentiation	R4:Chap:17:Pg.No:372-377
8	1	Rules of differentiation- simple standard forms	R6:Chap:2:Pg.No:2.6-2.8
9	1	Continuation rules of differentiation- simple standard forms	R6:Chap:2:Pg.No:2.8-2.10
10	1	Tutorial-II	
11	1	Applications of Differentiation in elasticity of demand and supply	R7:Chap:6:Pg.No:285-286
12	1	Maxima and Minima of functions	R7:Chap:6:Pg.No:287-286
13	1	Maxima and Minima of functions (involving second order derivatives) relating to cost, revenue and profit	R7:Chap:6:Pg.No:288-290
14	1	Maxima and Minima of functions (involving third order derivatives) relating to cost, revenue and profit	R7:Chap:6:Pg.No:293-294
15	1	Tutorial-III	
16	1	Recapitulation & discussion of possible questions	
	Total No	of Hours Planned For Unit II=16	
		UNIT-V	
1	1	Introduction on index numbers Meaning and its uses	R5:Chap:17:Pg.No:587-590

2	1	Construction of index numbers	R5:Chap:17:Pg.No:598-599
3	1	Problems on unweighted price index	R5:Chap:17:Pg.No:599-602
4	1	Problems on weighted price index	R5:Chap:17:Pg.No:605-607
5	1	Tutorial-I	
6	1	Tests of adequacy of index numbers	R5:Chap:17:Pg.No:617-618
7	1	Continuation of tests of adequacy of index numbers	R5:Chap:17:Pg.No:617-619
8	1	Construction of consumer price indices.	R5:Chap:17:Pg.No:632-634
9	1	Components of time series; additive and multiplicative models	R5:Chap:16:Pg.No:550-552
10	1	Tutorial-II	
11	1	Trend analysis: Finding trend by moving average method	R5:Chap:16:Pg.No:565-566
12	1	Fitting of linear trend line using principle of least squares	R5:Chap:16:Pg.No:566-567
13	1	Recapitulation & discussion of possible questions	
14	1	Discussion of previous ESE question papers	
15	1	Discussion of previous ESE question papers	
16	1	Discussion of previous ESE question papers	
	Total No	o of Hours Planned for unit V=16	
Total Plann ed Hours	48		

### **Suggested Reading**

- R1. Sreyashi Ghosh and Sujata Sinha (2018), Business Mathematics and Statistics, 1st edition,Oxford University Press; New Delhi.
- R2. Asim Kumar Manna (2018), Business Mathematics and Statistics, 1st edition, McGraw Hill Education, New Delhi.
- R3. S.P. Gupta and P.K. Gupta (2013), Business Statistics and Business Mathematics, S Chand Publishing, New Delhi.
- R4. Mariappan (2015), Business Mathematics, 1st edition, Pearson Education, NewDelhi.
- R5. J.K.Sharma, (2014) Business statistics, 4th edition, Vikas Publishing House, New Delhi

- R6. N. D. Vohra.(2013) Business Mathematics and Statistics. [Reprint]. New Delhi, McGraw Hill Education (India) Pvt Ltd.
- R7. Navainitham Pa.(2004).Business Mathematics & Statistics(3<sup>rd</sup> edition),Jai Publishers,India

**Class Representative** 

Signature of the Faculty

Tutor

**Programme Co-ordinator** 

HOD

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### <u>UNIT – I</u>

### SYLLABUS

### Matrices & Basic Mathematics of Finance

Definition of a matrix.Types of matrices; Algebra of matrices. Calculation of values of determinants up to third order; Adjoint of a matrix; Finding inverse of a matrix through ad joint; Applications of matrices to solution of simple business and economic problems- Simple and compound interest Rates of interest – nominal, effective and continuous – their interrelationships; Compounding and discounting of a sum using different types of rates

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### Matrix

An array of **mn** numbers arranged in **m** rows and **n** columns and bounded by square bracket [] brackets () or  $\| \|$  is called **m** by **n** matrix and is represented as

	a <sub>11</sub>	a <sub>12</sub>	 a <sub>1n</sub> –		( a <sub>11</sub>	a <sub>12</sub>	 $a_{1n}$	
	a 21	a 22	 $a_{2n}$		a <sub>21</sub>	a 22	 a <sub>2n</sub>	
۸ –	a <sub>31</sub>	a <sub>32</sub>	 a <sub>3n</sub>	_			 	
A –	 a <sub>i1</sub>	 a <sub>i2</sub>	   a <sub>in</sub>		a <sub>i1</sub>	$a_{i2}$	 a <sub>in</sub>	(1)
	•• <sub>11</sub>		 •• <sub>in</sub>				 	
	$a_{m1}$	a <sub>m2</sub>	 a <sub>mn</sub>		$(a_{m1})$	$a_{m2}$	 a <sub>mn</sub> )	

(1) is known as  $m \times n$  matrix in which there are m rows and n columns. Each member of  $m \times n$  matrix is known as an element of the matrix.

Note:

1. In general, we denote a Matrix by capital letter  $A = [a_{ij}]$ , where  $a_{ij}$  are elements of Matrix in which its position is in i<sup>th</sup> row and j<sup>th</sup> column i.e. first suffix denote row number and second suffix denote column number.

2. The elements  $a_{11}$ ,  $a_{22}$ ,..., $a_{nn}$  in which both suffix are same called diagonal elements, all other elements in which suffix are not same are called non diagonal elements.

3. The line along which the diagonal element lie is called the Principal Diagonal.

### **Different types of Matrices**

Zero Matrix or Null Matrix. A Matrix in which each elements is equal to zero is called a zero matrix or null matrix.

e.g.,

(0, 0)	<b>0</b>	$\left( 0 \right)$	0)		0	0	0	
$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$	olor	0	0	or	0	0	0	
	0)	0	0		0	0	0)	

are zero matrix respectively of order  $2 \times 3$ ;  $3 \times 2$  and  $3 \times 3$ .

In general we denote zero matrix of order  $m \times n$  by  $O_{m \times n}$ . Matrix other than Zero Matrix are called Non-Zero Matrix.

Square matrix. Matrix in which number of row becomes equal to number of column is called square matrix i.e. If matrix A is of type  $m \times n$ , where m = n then the matrix is called square matrix otherwise it is called rectangular matrix.

**Row Matrix.** A matrix of type  $1 \times n$ , having only one row is called row matrix. For example  $\begin{pmatrix} 1 & -2 & 3 \end{pmatrix}$  is a row matrix.

**Column Matrix.** A matrix of type m × 1, having only one column is called column matrix. For example  $\begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}$  is a

column matrix.

**Diagonal Matrix.** A square matrix in which all non diagonal elements are equal to zero is called diagonal matrix i.e.

A square matrix  $A = [a_{ij}]$ , is diagonal matrix if  $a_{ij} = 0$  for  $i \neq j$ . Thus

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

 $\begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \text{or} \begin{pmatrix} 2 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 3 \end{pmatrix} \text{ are diagonal matrices.}$ 

**Scalar Matrix.** Diagonal matrices in which all diagonal elements are equal are called scalar matrix i.e. A square matrix  $A = [a_{ij}]$ , is scalar matrix if  $a_{ij} = 0$  for  $i \neq j$  and  $a_{ij} = k$ , for i = j. Thus

 $(2 \ 0 \ 0)$ 

0 2 0 is scalar matrix.

 $\begin{pmatrix} 0 & 0 & 2 \end{pmatrix}$ 

Unit Matrix or Identity Matrix. A scalar matrix in which all diagonal elements are unity are called Unit matrix or Identity matrix generally denoted by  $I_n$ .

A square matrix  $A = [a_{ij}]$ , is Identity matrix if  $a_{ij} = 0$  for  $i \neq j$  and  $a_{ij} = 1$ , for i = j. Thus

- $(1 \ 0 \ 0)$
- 0 1 0, is identity matrix of order  $3 \times 3$ .

 $\begin{pmatrix} 0 & 0 & 1 \end{pmatrix}$ 

Triangular matrix are of two types:

(a) Upper Triangular Matrix. It is a matrix in which all elements below the principal diagonal are zero

e.g.

(b) Lower Triangular Matrix. It is a Matrix in which all elements above the principal diagonal are zero

e.g.

**Sub Matrix:** A matrix B obtained by deleting some rows or some column or both of a matrix A, is called a sub matrix of A.

For example. If  $A = \begin{pmatrix} 2 & 1 & 5 \\ 0 & -7 & 9 \\ -3 & 4 & 3 \end{pmatrix}$  then the matrices  $\begin{pmatrix} 2 & 1 \\ 0 & -7 \end{pmatrix}$ ,  $\begin{pmatrix} 2 & 1 & 5 \\ 0 & -7 & 9 \end{pmatrix}$  etc. are sub matrix of A.

### Transpose of a matrix

If matrix A is of type  $m \times n$ , then the matrix obtained by interchanging the rows and the columns of A is known as Transpose of Matrix A, denoted by A' or  $A^{T}$  i.e.

 $A = [a_{ij}]$  of m × n order then

A or  $A^{T} = [a_{ji}]$  of n × m order Matrix,

Now if A', B' be the transpose of matrix A and B respectively, then

(i) A = (A')' i.e. the transpose of transpose of a matrix A is matrix A itself.

(ii) (A + B)' = A' + B' i.e. the transpose of the sum of two matrices is equal to the sum of their transposes. (iii) (kA)'=kA', where k is a scalar.

(iv) (AB)' = B'A' i.e. the transpose of the product of two matrices is equal to the product of their transposes, taken in reversed order.

### **Conjugate of a Matrix**

Prepared by : Dr. Santhosh Kumar , Asst Prof, Department of Mathematics, KAHE

Page 3/31

# CLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IBATCH-2018-2021

If A be a matrix of order  $m \times n$ , over complex number system, then the matrix obtained from A by replacing each of its elements by their corresponding complex conjugates is called the conjugate of A and is denoted by  $\overline{A}$ , where  $\overline{A}$  is also of same order  $m \times n$ . If  $\overline{A}$ ,  $\overline{A}$  be the conjugate matrices of A, B respectively, then

(i) 
$$(\overline{A}) = A$$
.

(ii)  $\overline{(A+B)} = \overline{A} + \overline{B}$ , where A and B are conformable for addition.

- (iii)  $\overline{(kA)} = \overline{k}.\overline{A}$ , where k is any complex number.
- (iv)  $\overline{AB} = \overline{A}.\overline{B}$ , where A and B are conformable for multiplication.

### Transpose Conjugate of a Matrix

The transpose of the conjugate or conjugate of the transpose of a matrix A is called Transpose conjugate of A and is denoted by  $A^{\theta}$ . Thus

$$A^{\theta} = (\overline{A})' = \overline{(A')}.$$

If  $A^{\theta}$ ,  $B^{\theta}$  denote the transposed conjugate of A, B respectively, then

(i) 
$$(\mathbf{A}^{\theta})^{\theta} = \mathbf{A}$$

- (ii)  $(A+B)^{\theta} = A^{\theta} + B^{\theta}$ , where A and B are conformable for addition.
- (iii)  $(kA)^{\theta} = \overline{k} \cdot A^{\theta}$ , where k is any complex number.
- (iv)  $(AB)^{\theta} = B^{\theta}.A^{\theta}$ , where A and B are conformable for multiplication.

### Adjoint of a Square Matrix

If A is an n-rowed square matrix, then adjoint of A is defined as a transpose of matrix obtained by replacing each of its elements by its cofactors.

**Theorem 1.1:** If A be an n-square matrix, then A(adj. A)=  $(adj.A)A=|A|I_n$ , where  $I_n$  denotes the unit matrix of order n.

Theorem 1.2: If A and B are square matrix of the same order n, then adj. (AB)=(adj.B)(adj.A).

### **Inverse of Square Matrix**

Let A be n-square matrix, if there exist an n-square matrix B such that

 $AB = BA = I_n$ , then the matrix A is called invertible and the matrix B is called inverse of A. Inverse of a square matrix is denoted by  $A^{-1}$ .

Note. 1. From definition it is clear that A is the inverse of B.

2. A non-square matrix does not have any inverse.

### Singular and Non Singular Matrices

A square matrix A is said to be singular or non singular according as |A| = 0 or  $|A| \neq 0$ .

### **Orthogonal Matrix**

Any square matrix A is said to be orthogonal if  $AA^{=}A'A=I$ , this indicates that the row vectors (column vectors) of an orthogonal matrix A are mutually orthogonal unit vectors.

### **Unitary Matrix**

Any square A with complex elements is said to be unitary if  $A \cdot A^{\theta} = A^{\theta} A = I$ .

### CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### Symmetric And Skew Symmetric Matrices Symmetric Matrix

A square matrix  $A = [a_{ij}]$  is said to be symmetric if  $a_{ij} = a_{ji}$  for all i and j.

Examples  $\begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}, \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 4 \\ 3 & 4 & 7 \end{bmatrix}$ 

### **Skew Symmetric Matrix**

If a square matrix A has its elements such that $a_{ij} =$	$-a_{ji}$ for i and j and the leading diagonal elements are
	$\begin{bmatrix} 0 & 2 & -3 \end{bmatrix} \begin{bmatrix} 0 & h & g \end{bmatrix}$
zeros, then matrix A is known as skew matrix. For example	$\begin{vmatrix} -2 & 0 & 1 \end{vmatrix}$ , $\begin{vmatrix} -h & 0 & -f \end{vmatrix}$ are skew
	$\begin{bmatrix} 3 & -1 & 0 \end{bmatrix} = \begin{bmatrix} -g & f & 0 \end{bmatrix}$

symmetric matrices.

Example

Some examples of matrices are:

$$A = \begin{pmatrix} -4 & 3 & -6 & 9 \end{pmatrix} \quad B = \begin{pmatrix} 5 & -2 \\ 3 & -7 \end{pmatrix} \qquad C = \begin{pmatrix} 4 & -1 & 6 \\ 2 & 7 & -4 \\ 0 & 1 & -3 \end{pmatrix}$$

$$A = \begin{pmatrix} 4 & -3 & 6 \\ -8 & 5 & -9 \end{pmatrix} B = \begin{pmatrix} -5 & 6 & -2 \\ 3 & 7 & -4 \end{pmatrix}$$
  
1. (A)+(B) Given:

$$A + B = \begin{pmatrix} 4 + -5 & -3 + 6 & 6 + -2 \\ -8 + 3 & 5 + 7 & -9 + -4 \end{pmatrix} \Longrightarrow A + B = \begin{pmatrix} -1 & 3 & 4 \\ -5 & 12 & -13 \end{pmatrix}$$

Solution:


$$A = \begin{pmatrix} 6 & -7 \\ -4 & 5 \\ -3 & 2 \end{pmatrix} B = \begin{pmatrix} -8 & 3 \\ 3 & -1 \\ 2 & -8 \end{pmatrix}$$
  
2. (A)-(B) Given:


$$A - B = \begin{pmatrix} 6 - -8 & -7 - 3 \\ -4 - 3 & 5 - -1 \\ -3 - 2 & 2 - -8 \end{pmatrix} \implies A - B = \begin{pmatrix} 14 & -10 \\ -7 & 6 \\ -5 & 10 \end{pmatrix}$$

Solution:

$$5 \begin{pmatrix} -4 & 3 \\ 6 & -2 \end{pmatrix} \Rightarrow \text{Solution:} \begin{pmatrix} -20 & 15 \\ 30 & -10 \end{pmatrix}$$

$A = \begin{pmatrix} 6 & -2 & 3 \\ -4 & 2 & 5 \end{pmatrix}^{B} = \begin{pmatrix} 2 & -3 \\ 4 & -5 \\ 1 & -6 \end{pmatrix}$ 2. Given: $A = \begin{bmatrix} 1 & 0 \\ 3 & -5 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & 1 \\ 4 & -3 \end{bmatrix}$ Find: (a) $A + B$ $A + B = \begin{bmatrix} -1 & 1 \\ 7 & -8 \end{bmatrix}$ (b) $2A - 3B$ $2A - 3B = \begin{bmatrix} 8 & -3 \\ -6 & -1 \end{bmatrix}$ (c) $AB$ $AB = \begin{bmatrix} -2 & 1 \\ -26 & 18 \end{bmatrix}$ (d) $AB + BA$ $AB + BA = \begin{bmatrix} -1 & -4 \\ -31 & 33 \end{bmatrix}$ (e) $A^{T}B$ (where $A^{T}$ denotes the transpose of $A$ ) $A^{T}B = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) $B^{T}A$ $B^{T}A = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	KARPAC CLASS: I B.B.A. COURSE CODE: 18BAU202		DUCATION SS MATHEMATICS AND STATISTICS BATCH-2018-2021
Find: (a) $\mathbf{A} + \mathbf{B}$ $\mathbf{A} + \mathbf{B} = \begin{bmatrix} -1 & 1 \\ 7 & -8 \end{bmatrix}$ (b) $2\mathbf{A} - 3\mathbf{B}$ $2\mathbf{A} - 3\mathbf{B} = \begin{bmatrix} 8 & -3 \\ -6 & -1 \end{bmatrix}$ (c) $\mathbf{AB}$ $\mathbf{AB} = \begin{bmatrix} -2 & 1 \\ -26 & 18 \end{bmatrix}$ (d) $\mathbf{AB} + \mathbf{BA}$ $\mathbf{AB} + \mathbf{BA} = \begin{bmatrix} -1 & -4 \\ -31 & 33 \end{bmatrix}$ (e) $\mathbf{A}^{T}\mathbf{B}$ (where $\mathbf{A}^{T}$ denotes the transpose of $\mathbf{A}$ ) $\mathbf{A}^{T}\mathbf{B} = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) $\mathbf{B}^{T}\mathbf{A}$ $\mathbf{B}^{T}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	4. $(A)(B)$ Given:	$A = \begin{pmatrix} 6 & -2 & 3 \\ -4 & 2 & 5 \end{pmatrix} B = \begin{pmatrix} 2 & -3 \\ 4 & -5 \\ 1 & -6 \end{pmatrix}$	
(a) $\mathbf{A} + \mathbf{B}$ $\mathbf{A} + \mathbf{B} = \begin{bmatrix} -1 & 1 \\ 7 & -8 \end{bmatrix}$ (b) $2\mathbf{A} - 3\mathbf{B}$ $2\mathbf{A} - 3\mathbf{B} = \begin{bmatrix} 8 & -3 \\ -6 & -1 \end{bmatrix}$ (c) $\mathbf{A}\mathbf{B}$ $\mathbf{A}\mathbf{B} = \begin{bmatrix} -2 & 1 \\ -26 & 18 \end{bmatrix}$ (d) $\mathbf{A}\mathbf{B} + \mathbf{B}\mathbf{A}$ $\mathbf{A}\mathbf{B} + \mathbf{B}\mathbf{A} = \begin{bmatrix} -1 & -4 \\ -31 & 33 \end{bmatrix}$ (e) $\mathbf{A}^{\mathrm{T}}\mathbf{B}$ (where $\mathbf{A}^{\mathrm{T}}$ denotes the transpose of $\mathbf{A}$ ) $\mathbf{A}^{\mathrm{T}}\mathbf{B} = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) $\mathbf{B}^{\mathrm{T}}\mathbf{A}$ $\mathbf{B}^{\mathrm{T}}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$		and $\mathbf{B} = \begin{bmatrix} -2 & 1 \\ 4 & -3 \end{bmatrix}$	
(c) AB $AB = \begin{bmatrix} -2 & 1 \\ -26 & 18 \end{bmatrix}$ (d) AB + BA $AB + BA = \begin{bmatrix} -1 & -4 \\ -31 & 33 \end{bmatrix}$ (e) A <sup>T</sup> B (where A <sup>T</sup> denotes the transpose of A) $A^TB = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) B <sup>T</sup> A $B^TA = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$		$\mathbf{A} + \mathbf{B} = \begin{bmatrix} -1 & 1 \\ 7 & -8 \end{bmatrix}$	
(d) $\mathbf{AB} + \mathbf{BA}$ $\mathbf{AB} + \mathbf{BA} = \begin{bmatrix} -1 & -4 \\ -31 & 33 \end{bmatrix}$ (e) $\mathbf{A}^{\mathrm{T}}\mathbf{B}$ (where $\mathbf{A}^{\mathrm{T}}$ denotes the transpose of $\mathbf{A}$ ) $\mathbf{A}^{\mathrm{T}}\mathbf{B} = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) $\mathbf{B}^{\mathrm{T}}\mathbf{A}$ $\mathbf{B}^{\mathrm{T}}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	(b) 2 <b>A</b> -3 <b>B</b>	$2\mathbf{A} - 3\mathbf{B} = \begin{bmatrix} 8 & -3 \\ -6 & -1 \end{bmatrix}$	
(e) $\mathbf{A}^{\mathrm{T}}\mathbf{B}$ (where $\mathbf{A}^{\mathrm{T}}$ denotes the transpose of $\mathbf{A}$ ) $\mathbf{A}^{\mathrm{T}}\mathbf{B} = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) $\mathbf{B}^{\mathrm{T}}\mathbf{A}$ $\mathbf{B}^{\mathrm{T}}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	(c) <b>AB</b>	$\mathbf{AB} = \begin{bmatrix} -2 & 1 \\ -26 & 18 \end{bmatrix}$	
$\mathbf{A}^{T}\mathbf{B} = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$ (f) $\mathbf{B}^{T}\mathbf{A}$ $\mathbf{B}^{T}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	(d) <b>AB</b> + <b>BA</b>	$\mathbf{AB} + \mathbf{BA} = \begin{bmatrix} -1 & -4 \\ -31 & 33 \end{bmatrix}$	
(f) $\mathbf{B}^{\mathrm{T}}\mathbf{A}$ $\mathbf{B}^{\mathrm{T}}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	(e) $\mathbf{A}^{\mathrm{T}}\mathbf{B}$ (where $\mathbf{A}^{\mathrm{T}}$	denotes the transpose of A)	
		$\mathbf{A}^{T}\mathbf{B} = \begin{bmatrix} 10 & -8 \\ -20 & 15 \end{bmatrix}$	
	(f) <b>B</b> <sup>T</sup> <b>A</b>	$\mathbf{B}^{T}\mathbf{A} = \begin{bmatrix} 10 & -20 \\ -8 & 15 \end{bmatrix}$	
(g) $\mathbf{A}^2 \mathbf{B}$ $\mathbf{A}^2 \mathbf{B} = \begin{bmatrix} 1 & 1 \\ 124 & -87 \end{bmatrix}$	(g) <b>A</b> <sup>2</sup> <b>B</b>	$\mathbf{A^2B} = \begin{bmatrix} -2 & 1\\ 124 & -87 \end{bmatrix}$	





CLASS: I B.B.A. COURSE CODE: 18BAU2			S MATHEMATICS AND STATISTICS BATCH-2018-2021
(d) $\begin{bmatrix} 10 & 5 & 1 \\ 2 & -1 & 4 \\ 4 & 0 & 4 \end{bmatrix}$			
De	eterminant: 4		
Mi	inors:	$\begin{bmatrix} -4 & -8 & 4 \\ 20 & 36 & -20 \\ 21 & 38 & -20 \end{bmatrix}$	
Co	ofactors:	-4     8     4       -20     36     20       21     -38     -20	
inv	verse matrix:	-1       -5       5.25         2       9       -9.5         1       5       -5	

### The Cofactor Expansion for Determinants

Every square matrix has a determinant. All matrices with zero determinant are singular. All matrices with non-zero determinant are invertible.

The determinant of a (1×1) matrix A = [a] is just detA = a.

From section 2.3, the determinant of a (2×2) matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  is detA = ad - bc.

The determinants of all higher-order matrices can be expressed in terms of lower-order determinants.

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IBATCH-2018-2021

### **Determinants and Inverse Matrices**

For any set of square matrices of the same dimensions, the determinant of a product is the product of the determinants:

 $det (AB) = (detA) (detB), \quad det (ABC) = (detA) (detB) (detC), \quad etc.$ 

It then follows that  $\det(A^k) = (\det A)^k$   $AA^{-1} = I \implies \det(AA^{-1}) = 1 \implies \det A \det(A^{-1}) = 1 \implies \det(A^{-1}) = 1 \implies \det(A^{-1}) = \frac{1}{\det A}$ 

The **adjugate** (or **adjoint**) of any square matrix A is the transpose of the matrix of cofactors of A:

$ \frac{1}{1} \begin{pmatrix} 1 \end{pmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$	adi a	b	= $d$	-b
$\operatorname{adj}(A) = \left[c_{ij}(A)\right]^{\mathrm{T}}$	[The 2×2 case is $adj c$	d	$\lfloor -c$	$a \rfloor_{.]}$

Example 1


$$A = \begin{bmatrix} 1 & 9 & 1 \\ 0 & 2 & 0 \\ 1 & -4 & 2 \end{bmatrix}$$

Compute adj (A), Aadj (A) and det (A) for

The matrix of cofactors is

$$C = \begin{bmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{bmatrix} = \begin{bmatrix} + \begin{vmatrix} 2 & 0 \\ -4 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 0 \\ -4 & 2 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ 1 & 2 \end{vmatrix} - \begin{vmatrix} 1 & 9 \\ 1 & -4 \end{vmatrix} = \begin{bmatrix} 4 & 0 & -2 \\ -22 & 1 & 13 \\ -2 & 0 & 2 \end{bmatrix} + \begin{vmatrix} 9 & 1 \\ -2 & 0 & 2 \end{vmatrix}$$

KARPAGAM	ACADEMY OF HIGHER EL	DUCATION
CLASS: I B.B.A. COURSE CODE: 18BAU202	COURSE NAME:BUSINES UNIT: I	SS MATHEMATICS AND STATISTICS BATCH-2018-2021
$\operatorname{adj} A = C^{\mathrm{T}} = \begin{bmatrix} 4 & -22 & -2 \\ 0 & 1 & 0 \\ -2 & 13 & 2 \end{bmatrix}$		
$A \operatorname{adj} A = \begin{bmatrix} 1 & 9 & 1 \\ 0 & 2 & 0 \\ 1 & -4 & 2 \end{bmatrix} \begin{bmatrix} 4 & -4 & -4 \\ 0 & -2 & -2 \end{bmatrix}$	$\begin{bmatrix} -22 & -2 \\ 1 & 0 \\ 13 & 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = 2$	I
Expand along the middle row to det $A = 0 + 2 \times c_{22} + 0 = 2$	find detA :	
number that results from the diff	ts any $2 \times 2$ matrix, then the def	ninant. This value for a $2 \times 2$ matrix is the S the numbers in each diagonal of the terminant is the result of <b>ad</b> – <b>bc</b> . The
det $A = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad$ Notation:	d-bc	
Examples: Find the determinant	t of each of the following matric	ces:
$ \begin{array}{cccc} 5 & 3 \\ 7 & 2 \end{array} \qquad \qquad 2. \begin{array}{cccc} 6 & - \\ 2 & - \\ 2 & - \end{array} $	$ \begin{array}{c} 4\\3 \end{array} \qquad \qquad 3. \begin{array}{c} \begin{pmatrix} 4 & -2\\3 & 5 \end{array} \end{pmatrix} $	$ \begin{pmatrix} -3 & 8 \\ -2 & -4 \end{pmatrix} $
(5)(2) - (7)(3)(6)(-3) - (2)(-4)(10 - 21 - 188) -11 -10 2		(8)



Do not leave a negative sign in the denominator of any fraction. Apply the rule for division of integers and place the resulting sign in the numerator of the fraction.

When using the inverse matrix to solve a system of linear equations, it is best to leave all the elements in fraction form. To confirm that the answer is correct, multiply the matrix by its inverse.

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

$$\begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ 5 & -3 \\ 2 & 2 \end{pmatrix} \Longrightarrow \begin{pmatrix} 6 + \frac{-10}{2} & -3 + \frac{6}{2} \\ 10 + \frac{-20}{2} & -5 + \frac{12}{2} \end{pmatrix} \Longrightarrow \begin{pmatrix} 6 - 5 & -3 + 3 \\ 10 - 10 & -5 + 6 \end{pmatrix} \Longrightarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

The product of

Check:

 $(A) \times (A)^{-1}$  is the identity matrix. Therefore the inverse matrix is correct.

 $A = \begin{pmatrix} 2 & -6 \\ 1 & 3 \end{pmatrix}$ 

Solution: The first step is to find the determinant of (A).

$$|A| = 6 - -6$$
$$|A| = 12$$

$$A^{-1} = \begin{pmatrix} \frac{3}{12} & \frac{6}{12} \\ \frac{-1}{12} & \frac{2}{12} \end{pmatrix} \quad orA^{-1} = \begin{pmatrix} \frac{1}{4} & \frac{1}{2} \\ \frac{-1}{12} & \frac{1}{6} \end{pmatrix}$$

When using the inverse matrix to solve a system of linear equations, it is best to leave all of the elements in fraction form with the same common denominator as shown in the first inverse rather than as reduced fractions as shown in the second inverse.

$$\begin{pmatrix} 2 & -6 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} \frac{3}{12} & \frac{6}{12} \\ \frac{-1}{12} & \frac{2}{12} \end{pmatrix} \Rightarrow \begin{pmatrix} \frac{6}{12} + \frac{6}{12} & \frac{12}{12} - \frac{12}{12} \\ \frac{3}{12} - \frac{3}{12} & \frac{6}{12} + \frac{6}{12} \end{pmatrix} \Rightarrow \begin{pmatrix} \frac{12}{12} & \frac{0}{12} \\ \frac{0}{12} & \frac{12}{12} \end{pmatrix} \Rightarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

### Check:

The product of  $(A) \times (A)^{-1}$  is the identity matrix. Therefore the inverse matrix is correct.

The multiplication above was done using  $A^{-1}$  in which all the elements had the common denominator 12. Therefore, the resulting products could be manipulated in order to produce the identity matrix.

 $A = \begin{pmatrix} -10 & 4 \\ 5 & 2 \end{pmatrix}$ 

Solution: The first step is to find the determinant of (A).

COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS CLASS: I B.B.A. COURSE CODE: 18BAU202 BATCH-2018-2021 UNIT: I

$$|A| = -20 - 20$$
  

$$|A| = -40$$
  

$$A^{-1} = \begin{pmatrix} \frac{2}{-40} & \frac{-4}{-40} \\ \frac{-5}{-40} & \frac{-10}{-40} \end{pmatrix}$$
  

$$A^{-1} = \begin{pmatrix} \frac{-2}{40} & \frac{4}{40} \\ \frac{5}{40} & \frac{10}{40} \end{pmatrix} \quad orA^{-1} = \begin{pmatrix} \frac{-1}{20} & \frac{1}{10} \\ \frac{1}{8} & \frac{1}{4} \end{pmatrix}$$

Chaele	$\begin{pmatrix} -10\\ 5 \end{pmatrix}$	$\begin{pmatrix} 4\\ 2 \end{pmatrix}$		$\frac{4}{40} \\ \frac{10}{40}$	⇒	$\left(\frac{\frac{20}{40}}{\frac{-10}{40}}\right)$		$\frac{-40}{40}$ $\frac{20}{40}$ +	$\left( \frac{+\frac{40}{40}}{\frac{20}{40}} \right)^{\frac{1}{2}}$	$\Rightarrow \begin{pmatrix} \frac{40}{40} \\ \frac{0}{40} \\ \frac{1}{40} \end{pmatrix}$	$\overline{\begin{array}{c}0\\40\\40\end{array}}$	$\Rightarrow \begin{bmatrix} 1 \\ 0 \end{bmatrix}$	$\begin{pmatrix} 0\\1 \end{pmatrix}$
Check:	$( \mathbf{J} $	2) (	<b>4</b> 0	407		<b>\ 40</b>	40	40	40 )	(40	<i>40,</i>	/	

The product of  $(A) \times (A)^{-1}$  is the identity matrix. Therefore the inverse matrix is correct.

 $A = \begin{pmatrix} 6 & -3 \\ 5 & -2 \end{pmatrix}$ 

Solution: The first step is to find the determinant of (A). |A| = -12 - -15|A| = 3

$$A^{-1} = \begin{pmatrix} -2 & 3\\ 3 & 3\\ -5 & 6\\ \hline 3 & 3 \end{pmatrix} \quad or A^{-1} = \begin{pmatrix} -2 & 1\\ -5 & 2\\ \hline 3 & 2 \end{pmatrix}$$

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

$$\begin{pmatrix} -2 & \frac{3}{3} \\ \frac{-5}{5} & \frac{6}{3} \\ \frac{-5}{5} & -2 \end{pmatrix} \left( \begin{array}{ccc} \frac{-2}{3} & \frac{3}{3} \\ \frac{-5}{3} & \frac{6}{3} \\ \frac{-10}{3} + \frac{10}{3} & \frac{15}{3} - \frac{12}{3} \\ \frac{15}{3} - \frac{12}{3} \\$$

Check:

The product of  $(A) \times (A)^{-1}$  is the identity matrix. Therefore the inverse matrix is correct.

1. Find the determinant of each of the following matrices:

a) 
$$\begin{pmatrix} 2 & -3 \\ -4 & -10 \\ & & \\ \end{pmatrix}$$
 b)  $\begin{pmatrix} 7 & 3 \\ -4 & -2 \end{pmatrix}$  c)  $\begin{pmatrix} 5 & -2 \\ -3 & 2 \end{pmatrix}$  d)  $\begin{pmatrix} -8 & 5 \\ -4 & 3 \end{pmatrix}$ 

Solution :

$$\begin{pmatrix} 2 & -3 \\ -4 & -10 \\ a \end{pmatrix} \begin{pmatrix} 7 & 3 \\ -4 & -2 \end{pmatrix} \begin{pmatrix} 5 & -2 \\ -3 & 2 \end{pmatrix} \begin{pmatrix} -8 & 5 \\ -4 & 3 \end{pmatrix}$$
  
det = -2 det = -14 - -12 det = 10 - + 6 det -24 - 20 det = -4

2. Find the inverse of each of the following  $2 \times 2$  matrices. Check your solution by performing the operation of  $(A) \times (A)^{-1}$ .

a) 
$$A = \begin{pmatrix} 4 & 3 \\ -1 & -2 \end{pmatrix}$$
 b)  $A = \begin{pmatrix} 6 & -3 \\ 7 & -4 \end{pmatrix}$  c)  $A = \begin{pmatrix} 5 & 0 \\ -4 & 2 \end{pmatrix}$  d)  $A = \begin{pmatrix} -9 & 3 \\ 5 & -2 \end{pmatrix}$   
Solution :  
a)  $A = \begin{pmatrix} 4 & 3 \\ -1 & -2 \end{pmatrix}$  b)  $A = \begin{pmatrix} 6 & -3 \\ 7 & -4 \end{pmatrix}$  c)  $A = \begin{pmatrix} 5 & 0 \\ -4 & 2 \end{pmatrix}$  d)  $A = \begin{pmatrix} -9 & 3 \\ 5 & -2 \end{pmatrix}$   
 $|A| = -8 - 3 |A| = -24 - -21 |A| = 10 - 0 |A| = 18 - 15$   
 $|A| = -5 |A| = -3 |A| = 10 |A| = 3$   
 $A^{-1} = \begin{pmatrix} -2 & -3 \\ -5 & -5 \\ \frac{1}{-5} & \frac{4}{-5} \end{pmatrix} A^{-1} = \begin{pmatrix} -4 & 3 \\ -3 & -3 \\ -7 & 6 \\ -3 & -3 \end{pmatrix} A^{-1} = \begin{pmatrix} 2 & 0 \\ 10 & 10 \\ \frac{4}{-5} & 10 \end{pmatrix} A^{-1} = \begin{pmatrix} -2 & -3 \\ 3 & 3 \\ -5 & -9 \\ 3 & 3 \end{pmatrix}$ 

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

$$A^{-1} = \begin{pmatrix} \frac{2}{5} & \frac{3}{5} \\ \frac{-1}{5} & \frac{-4}{5} \end{pmatrix} A^{-1} = \begin{pmatrix} \frac{4}{3} & \frac{-3}{3} \\ \frac{7}{3} & \frac{-6}{3} \end{pmatrix} A^{-1} = \begin{pmatrix} \frac{1}{5} & 0 \\ \frac{2}{5} & \frac{1}{2} \end{pmatrix} A^{-1} = \begin{pmatrix} \frac{-2}{3} & -1 \\ \frac{-5}{3} & -3 \end{pmatrix} A^{-1} = \begin{pmatrix} \frac{1}{7} & -1 \\ \frac{7}{3} & -2 \end{pmatrix}$$

### **Minors and cofactors of a Matrix**

Let A = 
$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

**Minor of** 
$$a_{ij} \equiv M_{ij}$$
,

### is determinant obtained by deleting ith row and jth column.

 $M_{11} = \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$  is determinant obtained by deleting 1st row and 1st column

$$M_{11} = \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}, M_{12} = \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}, M_{13} = \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$
$$M_{21} = \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix}, M_{22} = \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix}, M_{23} = \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}$$
$$M_{13} = \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}, M_{32} = \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix}, M_{33} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

Cofactor of 
$$a_{ij} = C_{ij} = (-1)^{i+j} M_{ij}$$

**Signs of Cofactors** 

For 
$$2x^2$$
 – matrix  $\begin{bmatrix} + & - \\ - & + \end{bmatrix}$ 

	ACADEMY OF HIGHE	
CLASS: I B.B.A. COURSE CODE: 18BAU202	COURSE NAME:BUS UNIT: I	INESS MATHEMATICS AND STATISTICS BATCH-2018-2021
For 3x3 – matrix		
For 4x4 – matrix	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
Example:1. Find all minors an	d cofactors of the matr	ix
$A = \begin{bmatrix} 3 & 4\\ 1 & 0\\ 2 & 5 \end{bmatrix}$	$\begin{bmatrix} -1\\3\\-4 \end{bmatrix}$	
Solution:		
$M_{11} = \begin{vmatrix} 0 & 3 \\ 5 & -4 \end{vmatrix} = -15, M_{12} = M_{21} = \begin{vmatrix} 4 & -1 \\ 5 & -4 \end{vmatrix} = -11, M_{22}$ $M_{31} = \begin{vmatrix} 4 & -1 \\ 0 & 3 \end{vmatrix} = 12, M_{32}$	$= \begin{vmatrix} 3 & -1 \\ 2 & -4 \end{vmatrix} = -10, M_{23} = \begin{vmatrix} 3 \\ -1 \\ 2 \end{vmatrix}$	$\begin{vmatrix} 3 & 4 \\ 2 & 5 \end{vmatrix} = 7$
Cofactor of $a_{ii}$	$= C_{ii} = (-1)^{i+j} M_{ii}$	
y y	y	
$C_{11} = -15,  C_{12} = 0$ $C_{21} = 11,  C_{22} = 0$ $C_{31} = 12,  C_{32} = 0$	$= 10,   C_{13} = 5$ = -10,   C_{23} = -7 = -10,   C_{33} = -4	
NOTE: Matrix of cofactors,	$C = \begin{bmatrix} -15 & 10 & 5\\ 11 & -10 & -7\\ 12 & -10 & -7 \end{bmatrix}$	,

CLASS: I B.B.A. COURSE CODE: 18BAU202

NOTE: Determinant of matrix of Cofactors by the method of Cofactors

 $det(A) = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$  $det(A) = a_{21}C_{21} + a_{22}C_{22} + a_{23}C_{23}$  $det(A) = a_{31}C_{31} + a_{32}C_{32} + a_{33}C_{33}$ 

The above equations can be used to check that the cofactors are found correctly as the values of determinants found must be equal, we open matrix from any row or column.

Example: 2 .Find the determinant of the matrix A by method of cofactors,

$$A = \begin{bmatrix} 3 & 4 & -1 \\ 1 & 0 & 3 \\ 2 & 5 & -4 \end{bmatrix}$$

**Solution:** : Using the cofactors found in the last example

$$det(A) = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$
  
= 3(-15)+4(10)+(-1)(5)  
=-45 + 40 - 5 = -10

**NOTE: 3.**We can find determinant by opening matrix from second or third row or first column, the value of the determinant will be same

$$det(A) = a_{21}C_{21} + a_{22}C_{22} + a_{23}C_{23}$$
  
=(1)(11)+0(-10)+3(-7)=11 - 21 = -10

$$det(A) = a_{31}C_{31} + a_{32}C_{32} + a_{33}C_{33}$$
  
= 2(12) +5(-10)+(-4)(-4) = 24 - 50 + 16 = -10

**NOTE : 4**. Determinant of A can be obtained by multiplying any row or any column of matrix A with the corresponding row or column of the matrix of cofactors.

**NOTE: 5.** Determinant of matrix  $\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ 

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IBATCH-2018-2021

$$\det \mathbf{A} = \begin{array}{cc} a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

**Inverse by method of Cofactors:** 

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \det \mathbf{A} \neq \mathbf{0}.$$

Find Matrix of cofactors

$$\mathbf{C} = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

Step : 2. Find Adjoint of matrix A, adj(A)

 $\mathbf{Adj}(\mathbf{A}) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^{T}$ 

Step: 3.

If A is an invertible matrix,  $det(A) \neq 0$ , then

$$A^{-1} = \frac{1}{\det A} [adj(A)]$$

**Example: 3**. Find A<sup>-1</sup> of matrix A

$$A = \begin{bmatrix} 2 & 0 & 3 \\ 0 & 3 & 2 \\ -2 & 0 & -4 \end{bmatrix}$$
 by the method of cofactors.

CLASS: I B.B.A. COURSE CODE: 18BAU202

Solution: Cofactors of the matrix A are

$$C_{11} = \begin{vmatrix} 3 & 2 \\ 0 & -4 \end{vmatrix} = -12, C_{12} = -\begin{vmatrix} 0 & 2 \\ -2 & -4 \end{vmatrix} = -4, C_{13} = \begin{vmatrix} 0 & 3 \\ -2 & 0 \end{vmatrix} = 6$$
  
$$C_{21} = -\begin{vmatrix} 0 & 3 \\ 0 & 4 \end{vmatrix} = 0, \quad C_{22} = \begin{vmatrix} 2 & 3 \\ -2 & -4 \end{vmatrix} = -2, \quad C_{23} = -\begin{vmatrix} 2 & 0 \\ -2 & 0 \end{vmatrix} = 0,$$
  
$$C_{31} = \begin{vmatrix} 0 & 3 \\ 3 & 2 \end{vmatrix} = -9, \quad C_{32} = -\begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} = -4, \quad C_{33} = \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix} = 6$$

Matrix of cofactors, 
$$C = \begin{bmatrix} -12 & -4 & 6 \\ 0 & -2 & 0 \\ -9 & -4 & 6 \end{bmatrix}$$

Adjoint of matrix A, 
$$adj(A) = \begin{bmatrix} -12 & 0 & -9 \\ -4 & -2 & -4 \\ 6 & 0 & 6 \end{bmatrix}$$

$$det(A) = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$
  
= 2(-12)+0(-4)+3(6)  
=-24 + 18 = -6 \ne 0

Inverse of matrix A is

$$A^{-1} = \frac{1}{\det A} [adj(A)] = \frac{1}{-6} \begin{bmatrix} -12 & 0 & -9 \\ -4 & -2 & -4 \\ 6 & 0 & 6 \end{bmatrix}$$

**NOTE :** If we can find  $A^{-1}$ , then solution of linear system AX = B is  $X = A^{-1}B$ 

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B.B.A.	COURSE NAME:BUSI	NESS MATHEMATICS AND STATISTICS		
COURSE CODE: 18BAU202	UNIT: I	BATCH-2018-2021		

### **Adjoint:**

The adjoint is the transpose of the cofactors  $(c^{T})$ 

To find the cofactors you find the 'minor' of each element (and use the alternate signs as shown above).

If: A = $\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$ Matrix of cofactors =	The minors of each element + $\begin{vmatrix} e \\ b \end{vmatrix} = \begin{vmatrix} d & f \\ c & i \end{vmatrix} + \begin{vmatrix} d & e \\ c & b \end{vmatrix}$	
	$ + \begin{vmatrix} e \\ h \end{vmatrix} \begin{bmatrix} f \\ i \end{vmatrix} - \begin{vmatrix} d & f \\ g & i \end{vmatrix} + \begin{vmatrix} d & e \\ g & h \end{vmatrix} $ $ - \begin{vmatrix} b & c \\ h & i \end{vmatrix} + \begin{vmatrix} a & c \\ g & i \end{vmatrix} - \begin{vmatrix} a & b \\ g & h \end{vmatrix} $ $ + \begin{vmatrix} b & c \\ e & f \end{vmatrix} - \begin{vmatrix} a & c \\ d & f \end{vmatrix} + \begin{vmatrix} a & b \\ d & e \end{vmatrix} $	

Nb. To find the adjoint we then need to transpose this matrix.

**Determinant**:  $|A| = +2\begin{vmatrix} 3 & 2 \\ 1 & 1 \end{vmatrix} - 1\begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix} + 5\begin{vmatrix} 3 & 3 \\ 4 & 1 \end{vmatrix}$ = 2(1) - 1(-5) + 5(-9)= -38  $\begin{bmatrix} + \begin{vmatrix} 3 & 2 \\ 1 & 1 \end{vmatrix} - \begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix} + \begin{vmatrix} 3 & 3 \\ 4 & 1 \end{vmatrix} = \begin{bmatrix} 1 & 5 & -9 \\ 4 & -18 & 2 \\ -13 & 11 & 3 \end{bmatrix}$  $= \begin{bmatrix} 1 & 5 & -9 \\ 4 & -18 & 2 \\ -13 & 11 & 3 \end{bmatrix}$ **Cofactors** =  $\begin{bmatrix} 1 & 4 & -13 \\ 5 & -18 & 11 \end{bmatrix}$  $Adjoint(C^T) :=$ Inverse:  $A^{-1} = \frac{1}{|A|} x adj = -\frac{1}{38} \begin{bmatrix} 1 & 4 & -13 \\ 5 & -18 & 11 \\ -9 & 2 & 3 \end{bmatrix}$ 

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### Simple Interest

The price to be paid for the use of a certain amount of money (called principal) for a certain period is known as Interest. The interest is payable yearly, half-yearly, quarterly or monthly.

The sum of the principal and interest due at any time, is called the Amount at that time.

The rate of interest is the interest charged on one unit of principal for one year and is denoted by i. If the principal is  $\overline{\mathbf{x}}$  100 then the interest charged for one year is usually called the amount of interest per annum, and is denoted by r ( = Pi).

e.g. if the principal is ₹ 100 and the interest ₹ 3, then we say usually that the rate of interest is 3 percent per annum (or r = 3 %)

Here  $i = \frac{3}{100} = 0.03$  (i.e. interest for 1 rupee for one year).

Simple interest is calculated always on the original principal for the total period for which the sum (principal) is used.

Let P be the principal (original)

n be the number of years for which the principal is used

r be the rate of interest p.a.

I be the amount of interest

i be the rate of interest per unit (i.e. interest on Re. 1 for one year)]

Now I = P.i.n, where  $i = \frac{r}{100}$ 

Amount A = P + I = P + P. i. n = P (1 + i.n) i.e. A = P (1 + n.i)

### SOLVED EXAMPLES :

1 Amit deposited ₹ 1200 to a bank at 9% interest p.a. find the total interest that he will get at the end of 3 years.

Here P = 1200, 
$$i = \frac{9}{100} = 0.09$$
, n = 3, l = ?

I = P. i.n = 1200 x 0.09 x 3 = 324.

Amit will get ₹ 324 as interest.

Sumit borrowed ₹ 7500 at 14.5% p.a. for  $2\frac{1}{2}$  years. Find the amount he had to pay after that 2.

### period.

P = 7500, 
$$i = \frac{14.5}{100} = 0.145$$
,  $n = 2\frac{1}{2} = 2.5$ , A = ?

A = P (1+in) = 7500 (1+0.145 x 2.5) = 7500 (1+0.3625)

= 7500 × 1.3625 = 10218.75

Reqd. amount = ₹ 10218.75.

COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### 3. Find the simple interest on ₹ 5600 at 12% p.a. from July 15 to September 26, 2013.

Time = number of days from July 15 to Sept. 26

= 16 (July) + 31 (Aug.) + 26 (Sept.)= 73 days.

P = 5600, 
$$i = \frac{12}{100} = 0.12$$
,  $n = \frac{73}{365}$  yr.  $= \frac{1}{5}$  yr.  
S.I. = P. i.n. = 5600 x 0.12 x  $\frac{1}{5} = 134.4$ 

To find Principle :

CLASS: I B.B.A.

**COURSE CODE: 18BAU202** 

### 4. What sum of money will amount to ₹ 1380 in 3 years at 5% p.a. simple interest?

Here A = 1380, n = 3, i = 
$$\frac{5}{100}$$
 = 0.05, P = ?  
From A = P (1 + 0.05x3) or, 1380 = P (1+0.15)  
Or, 1380 = P (1.15) or, P =  $\frac{1380}{1.15}$  = 1200

Problems to find rate % :

At what rate percent will a sum, become double of itself in  $5\frac{1}{2}$  years at simple interest?  $A = 2P, P = Principal, n = 5\frac{1}{2}, i = ?$   $A = P(1 + ni) \text{ or, } 2P = P(1 + \frac{11}{2} i)$ or,  $2 = 1 + \frac{11}{2} i \text{ or, } i = \frac{2}{11}$ or,  $n = \frac{2}{2} \times 100 = 18.18 \text{ (approx)}$ : Read, rate = 18.18%.

$$11^{-11}$$
  $11^{-11}$   $100^{-10.10}$  (approx), kequ. tale

### Problems to find time :

### 6 In how many years will a sum be double of itself at 10% p.a. simple interest.

A = 2P, P = Principal, 
$$i = \frac{10}{100} = 0.10$$
, n= ?  
A = P (1 + ni) or, 2P = P [1 + n(.10)] or, 2 = 1 + n (.10)  
or, n (.10) = 1 or, n =  $\frac{1}{.10} = 10$   $\therefore$  Read. time = 10 years.

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

## COMPOUND INTEREST

Interest as soon as it is due after a certain period, is added to the principal and the interest for the succeeding period is based upon the principal and interest added together. Hence the principal does not remain same, but increases at the end of each interest period.

A year is generally taken as the interest-period, but in most cases it may be half-year or quarter-year.

### Symbols :

Let P be the Principal (original)

A be the amount

i be the Interest on Re. 1 for 1 year

n be the Number of years (interest period).

### Formula : A = P(1 + i)<sup>n</sup> .... (i)

Cor.1. In formula (i) since P amount to A in years, P may be said to be present value of the sum A due in n years.

 $\mathsf{P} = \frac{\mathsf{A}}{(1+i)^n} = \mathsf{A}(1+i)^{-n}$ 

Cor.2. Formula (1) may be written as follows by using logarithm :

 $\log A = \log P + n \log (1 + i)$ 

### FEW FORMULAE :

Compound Interest may be paid half-yearly, quarterly, monthly instead of a year. In these cases difference in formulae are shown below :

(Taken P = principal, A = amount, T = total interest, i = interest on Re. 1 for 1 year, n = number of years.)

	Time	Amount	I = A - P
(i) Annual		A= P(1 + I) <sup>n</sup>	$I = P \{(1 + i)^n - 1\}$
(iii)	Half-Yearly	$A = P 1 + \frac{i}{2}^{2n}$	$I = P$ $1 + \frac{i}{2}^{2n} - 1$
(111)	Quarterly	$A = P  1 + \frac{i}{4}  4n$	$I = P + \frac{i}{4} + \frac{i}{4} - 1$
gene	eral if C.I. is paid p tim	es in a year, then $A = P 1 + \frac{i}{p}$	pn
:Le	t P = <b>₹ 1</b> 000, r = 5% i.e.	i = 0.05, n = 24 yrs.	

If interest is payable yearly the  $A = 1000 (1 + 0.5)^{24}$ 

CLASS: I B.B.A.	<u>A ACADEMY OF HIGHER E</u> COURSE NAME:BUSINE	ESS MATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: I	BATCH-2018-2021
	0.05 <sup>2×24</sup>	
If int. is payable half-yearly the	he A = $\frac{1000}{1 + \frac{0.00}{2}}$	
	0.05 <sup>4×24</sup>	
If int. is payable quarterly the	$A = 1000 + \frac{4}{4}$	
Note. or r = 100 I = interest p	er hundred.	
If r = 6% then If, however i =	0.02 then, r = 100 × 0.02 =	2%.
SOLVED EXAMPLES		
Find the compound intere	est on ₹ 1,000 for 4 years at	5% p.a.
Here P = ₹ 1000, n = 4, i = 0.0	05, A = ?	
We have $A = P(1 + i)^n$		
A = 1000 (1+0.05)4		
Or log A = log 1000 + 4log (	1 + 0.05) = 3 + 4 log (1.05)	= 3 + 4 (0.0212) = 3 + 0.0848 = 3.0848
A = antilog 3.0848 = 1215		
C.I. = ₹ 1215 - ₹ 1000 = ₹ 215	5	
2. In what time will a sum of	f money double itself at 59	% p.a. C.I.
Here, P = P, A = 2P, i = 0.05,	n = ?	
$A = P (1 + i)^n$ or, $2P = P(1 + 0)^n$	0.05) <sup>n</sup> = P (1.05) <sup>n</sup>	
or, 2 = (1.05) <sup>n</sup> or log 2 = n log	g 1.05	
$\therefore n = \frac{\log 2}{\log 1.05} = \frac{0.3010}{0.0212} = 14.2$	2 years (Approx)	
		erest on a sum put out for 5 years at 3%
was ₹ 46.80. Find the sum.		

# **KARPAGAM ACADEMY OF HIGHER EDUCATION** CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 BATCH-2018-2021 **UNIT: I** Let P = 100, i = .03, n = 5. From A = P $(1 + i)^n$ , $A = 100 (1 + .03)^{5} = 100 (1.03)^{5}$ $\log A = \log 100 + 5 \log (1.03) = 2 + 5 (.0128) = 2 + .0640 = 2.0640$ ∴ A = antilog 2.0640 = 115.9 ∴ C.I. = 115.9 - 100 = 15.9 Again S.I. = 3 × 5 = 15. ∴ difference 15.9 - 15 = 0.9 Capital $x = 100 \times \frac{46.80}{0.9} = 5,200$ Diff. 0.9 100 46.80 Х ∴ original sum = ₹ 5,200. 4 What is the present value of ₹1,000 due in 2 years at 5% compound interest, according as the interest is paid (a) yearly, (b) half-yearly? (a) Here A = ₹ 1,000, i = $\frac{5}{100}$ = 0.05, n = 2, P = ? $A = P (1 + i)^n \text{ or } 1000 = P (1 + .05)^2 = P (1.05)^2$ $\therefore P = \frac{1000}{(1.05)^2} = \frac{1000}{1.1025} = 907.03$ .: Present value = ₹ 907.03 (b) Interest per unit per half-year $\frac{1}{2} \times 0.05 = 0.025$ From A = P $1 + \frac{1}{2}^{2n}$ we find. $1,000 = P + \frac{0.05}{2} = P + (1 + .025)^4 = P + (1.025)^4$ or, $P = \frac{1000}{(1.025)^4}$ $\therefore \log P = \log 1000 - 4\log (1.025) = 3 - 4 (0.0107) = 3 - 0.0428 = 2.9572$ ∴ P = antilog 2.9572 = 906.1. Hence the present amount =₹906.10

### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 BATCH-2018-2021 **UNIT: I**

### Present value of an annuity :

Definition: Present value of an annuity is the sum of the present values of all payments (or instalments) made at successive annuity periods.

### Formula :

CLASS: I B.B.A.

(i) The present value V of an annuity P to continue for n years is given by

$$V = \frac{P}{i} \left\{ 1 - (1+i)^{-n} \right\}$$
 Where i = interest per rupee per annum.

(ii) The Present value V of an annuity P payable half-yearly, then

$$V = \frac{2P}{i} \quad 1 - 1 + \frac{i}{2}^{-2n}$$

(ii) The Present value V of an annuity P payable quarterly, then

$$V = \frac{4P}{1} \quad 1 - 1 + \frac{1}{4}^{-4}$$

### SOLVED EXAMPLES :

A man decides to deposit ₹ 20,000 at the end of each year in a bank which pays 10% p.a. compound interest. If the instalments are allowed to accumulate, what will be the total accumulation at the end of 9 years? Solution :

Let ₹ A be the total accumulation at the end of 9 years. Then we have

$$A = \frac{P}{i} \left\{ \left( 1 + i \right)^n - 1 \right\}$$

Here P = ₹ 20,000, i =  $\frac{10}{100}$  = 0.1, n = 9 years.

$$\therefore A = \frac{20,000}{0.1} \left\{ (1+.1)^{\circ} - 1 \right\} = 2,00,000 \left\{ (1.1)^{\circ} - 1 \right\} = 2,00,000 (2.3579 - 1)$$

= 2,00,000 x 1.3579 = ₹ 2,71,590

... The required total accumulation =₹2,71,590.

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### Nominal and Effective Interest

An interest rate takes two forms: nominal interest rate and effective interest rate. The nominal interest rate does not take into account the compounding period. The effective interest rate does take the compounding period into account and thus is a more accurate measure of interest charges.

A statement that the "interest rate is 10%" means that interest is 10% per year, compounded annually. In this case, the nominal annual interest rate is 10%, and the effective annual interest rate is also 10%. However, if compounding is more frequent than once per year, then the effective interest rate will be greater than 10%. The more often compounding occurs, the higher the effective interest rate.

The relationship between nominal annual and effective annual interest rates is:

 $i_a = [1 + (r/m)]^m - 1$ 

where " $i_a$ " is the effective annual interest rate, "r" is the nominal annual interest rate, and "m" is the number of compounding periods per year.

Example: A credit card company charges 21% interest per year, compounded monthly. What effective annual interest rate does the company charge?

 $\begin{aligned} r &= 0.21 \text{ per year} \\ m &= 12 \text{ months per year} \\ i_a &= \left[ 1 + (.21 / 12) \right]^{12} - 1 \\ &= \left[ 1 + 0.0175 \right]^{12} - 1 \\ &= (1.0175)^{12} - 1 = 1.2314 - 1 \\ &= 0.2314 = 23.14\% \end{aligned}$ 

It may be desired to find the effective interest rate for a period other than annual. In this case, adjust the period for "r" and "m" as needed. For example, if the effective interest rate per semi annual period (every 6 months) is desired, then

r = nominal interest rate per 6 months

m = number of compounding periods per 6 months

and the effective interest rate, isa, per semi-annual period, is:

 $i_{sa} = [1 + (r / m)]^m - 1$ 

2. If a lender charges 12% interest, compounded monthly, what is the effective interest rate per quarter?

Hint: m = number of compounding periods per quarter

Let i = effective interest rate per quarter.

### KARPAGAM ACADEMY OF HIGHER EDUCATION CLASS: I B.B.A.

COURSE CODE: 18BAU202

### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS **UNIT: I** BATCH-2018-2021

Choose an answer by clicking on one of the letters below, or click on "Review topic" if needed.

A i =  $[1 + (0.12/3)]^3 - 1 = (1.04)^3 - 1 = 0.1249 = 12.49\%$ **B** i =  $[1 + 0.03]^{12} - 1 = (1.03)^{12} - 1 = 0.4258 = 42.58\%$  $\underline{C}$  i =  $[1 + (0.03 / 3)]^3 - 1 = (1.01)^3 - 1 = 0.0303 = 3.03\%$  $Di = [1 + (0.03 / 12)]^3 - 1 = (1.0025)^3 - 1 = 0.0075 = 0.75\%$ 

### Continuous Compounding

Single payment formulas for continuous compounding are determined by taking the limit of compound interest formulas as m approaches infinity, where m is the number of compounding periods per year. Here "e" is the exponential constant (sometimes called Euler's number).

With continuous compounding at nominal annual interest rate r (time-unit, e.g. year) and n is the number of time units we have:

 $F = P e^{r n} F/P$ 

 $\mathbf{P} = \mathbf{F} \mathbf{e}^{-\mathbf{r} \mathbf{n}} \mathbf{P} / \mathbf{F}$ 

 $i_a = e^r - 1$  Actual interest rate for the time unit

Example 1: If \$100 is invested at 8% interest per year, compounded continuously, how much will be in the account after 5 years?

P = \$100

r = 8%

n = 5 years  $F = P e^{r n} = (\$100) e^{(.08)(5)}$ 

 $=(\$100) e^{0.4} = (\$100)(1.4918) = \$149.18$ 

Example 2: If \$100 is invested at 0.667% interest per month, compounded continuously, how much will be in the account after 5 years?

P =\$100

r = 0.666667%

n = 5 years \* 12 months

 $F = P e^{r n} = (\$100) e^{(.006666667)(60)}$ 

 $=(\$100) e^{0.4} = (\$100)(1.4918) = \$149.18$ 

Prepared by : Dr. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

Page 29/31

### CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: I BATCH-2018-2021

### POSSIBLE QUESTIONS 2 MARK QUESTIONS

- 1. What is an upper triangular matrix? Give example.
- 2. Write the formula for calculates Simple interest and compound interest?
- 3. Find the simple interest on the sum of Rs.6,000 at 10% p.a. for 3 years?
- 4. Find the inverse of  $A = \begin{pmatrix} 5 & 5 \\ 8 & 6 \end{pmatrix}$
- 5. What is singular matrix and non singular matrix?

6. If 
$$A = \begin{bmatrix} 3 & 5 \\ 1 & 9 \end{bmatrix}$$
 then show that  $(A')' = A$   
7. Evaluate  $A = \begin{bmatrix} 6 & 4 & 2 \\ 7 & 1 & 3 \\ 0 & -1 & 6 \end{bmatrix}$   
8. If  $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$  and  $B = \begin{bmatrix} 5 & 4 \\ 3 & 4 \end{bmatrix}$ , then find (A+B), (A-B).

1. If 
$$A = \begin{bmatrix} 1 & 0 & -2 \\ 2 & 2 & 4 \\ 0 & 0 & 2 \end{bmatrix}$$
, Find the inverse of A.

2. Mr. Ravi borrows Rs.20000 at 4% compound interest and agrees to pay both the principal and the interest in 10 equal installments at the end of each year. Find the amount of these installments.

3. If 
$$A = \begin{bmatrix} 2 & 3 & 5 \\ 4 & 7 & 9 \\ 1 & 6 & 4 \end{bmatrix}$$
 and  $B = \begin{bmatrix} 3 & 1 & 2 \\ 4 & 2 & 5 \\ 6 & -2 & 7 \end{bmatrix}$ , Show that  $5(A + B) = 5A + 5B$ .

- 4. Mr. Puthiyanayagam has two daughters A and B aged 10 <sup>1</sup>/<sub>2</sub> and 16 years. He has
- Rs.180, 000 with him now but he wants that both of them should get an equal amount when they are 18 years old. How he should divide the money if it were to be deposited with his friend who gives only 5 % p.a.
- 5. Verify that  $B^T A^T = (AB)^T$ , when  $A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 2 \\ 2 & 0 \\ -1 & 1 \end{bmatrix}$

6. i) If a term deposit of Rs.4000 earns an interest of Rs.2500 in 50 months find the rate of interest.

ii) Find the present value of Rs.695 due 2 years from now receiving simple interest at 5% per annum.

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B.B.A. COURSE CODE: 18BAU202	COURSE NAME:BUS UNIT: I	INESS MATHEMATICS AND STATISTICS BATCH-2018-2021		
7. If A= $\begin{bmatrix} 1 & -1 & 1 \\ 2 & 3 & 0 \\ 18 & 2 & 10 \end{bmatrix}$ , find t	the inverse of the given mat	rix A.		
interest being payable h ii). Find the simple intere	alf yearly. est on the sum of Rs.3000 at			
10. A person has two daughters	A and b aged 13 and 16 yea get an equal amount when t	-5I = 0.hence deduce the inverse of A. ars. He has Rs.40, 000 with him now but hey are 20 years old. How he should divide ompound interest p.		
11. Find the effective rate of intention find the effective rate when intention		te of 12% p.a., compounded monthly. Further, ously.		

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
A rectangular arrangement of numbers in rows and columns is called	set	matrix	order	sub matrix			matrix
A scalar matrix in which all the leading diagonal elements are unity is called-	square matrix	unit matrix	null matrix	Identity matrix			unit matrix
A diagonal matrix in which all the diagonal elements are equal is called	diagonal matrix	scalar matrix	unit matrix	null matrix			scalar matrix
[ 3 8 9 -2 ] is a row matrix of order	4 x 1	1x 4	1x 1	4 x 4			1x 4
A square matrix such that A' = A is called	symmetric	skew symmetric	hermitiIan	scalar			symmetric
An association of a real numbers with a matrix is known as	real matrix	complex matrix	null matrix	unit matrix			real matrix
The number of elements in an m x n matrix is	mn	n	m <sup>2</sup>	n <sup>2</sup>			mn
If in a square matrix A, $a_{ij} = 0$ for i>j, then it is called a matrix.	lower triangular	upper triangular	diagonal	scalar matrix			upper triangular
$(A^{T})^{T} =$	A <sup>-1</sup>	$\mathbf{A}^{\mathrm{T}}$	А	1 / A			А
A matrix A is said to be Orthogonal matrix if	$AA^{T} = I$	$AA^{T}=0$	$A=A^{T}$	A = 1/ A			$AA^{T} = I$
The addition of two matrices are possible only when they are	of same order	of any order	scalar matrices	unit matrices			of same order
A square matrix is said to be singular if its determinant is	0	1	2	-1			0
If A is non singular ,its inverse is	Adj A /  A	A  / Adj A	1/Adj A	1/ A			Adj A /  A

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
The condition for existing of A <sup>-1</sup> is	A is non singula	A is singular	A <sup>T</sup> is singular	A square matrix			A is non singular
in a square matrix A, $a_{ij} = 0$ for $i < j$ , then it is called a matrix.	lower triangular	upper triangular	diagonal	triangular			lower triangular
A Square matrix such that $A' = -A$ is called	symmetric	skew symmetric	hermit ion	scalar			skew symmetric
The sum of the diagonal elements in a matrix A is the	Trace of A	unit of A	Transpose of A	inverse of A			Trace of A
If a square matrix A of order n is of the form $A^n = 0$ , then it is an matrix matrix $A$ of order n is of the form $A^n = 0$ , then it is an matrix $A$ of order n is of the form $A^n = 0$ , then it is an	Identity	Idempotent	Nilpotent	Orthogonal			Nilpotent
If a square matrix A of order n is of the form $A^n = A$ , then it is an m	Identity	Idempotent	Nilpotent	Orthogonal			Idempotent
$(AB)^{-1} =$	$A^{-1}$	B <sup>-1</sup>	$A^{-1}B^{-1}$	$\mathrm{B}^{-1}\mathrm{A}^{-1}$			$B^{-1}A^{-1}$
I*A =	А	0	Identity matrix	Zero matrix			А
An identity matrix is also found as a matrix	Scalar	Diagonal	triangular	scalar and diagon	al		scalar and diagonal
In a 3x3 square matrix the minor and the cofactor of the element $a_{23}$ have	Same sign and sa	same sign and dif	Opposite sign and	opposite sign and different values		es	Opposite sign and sar
Every matrix A and its transpose A <sup>T</sup> have determinant value	Different	same	one as the negativ	one as the multiple of other			same
The determinant value of the unit matrix of order 2 is	1	0	-1	2			1
If A is singular ,its inverse is	null matrix	does not exist	1/Adj A	1/  <b>A</b>			does not exists

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
A rectangular matrix will not possesses	inverse	cofactor	determinant	transpose			inverse
For any square matrix (adj A).A = A. (adj A) is	A  .I	A	1 /  A	$A^{T}$			A  .I
For any two square matrices A and B (adj AB) =	(adj A). (adj B)	(adj B). (adj A)	(adj BA)	(adj B +adj A)			(adj B). (adj A)
The formula for computing simple Interest is	pnr / 100	p+n+r /100	p-n-r /100	p+n+r			pnr / 100
Simple interest will be an income for	lender	borrower	customer	creditor			lender
The simple interest on Rs. 5000 at 10 % for 3 years is	Rs.500	Rs.1000	Rs.1500	Rs.2000			Rs.1500
If the simple interest for 2 years at 12 % is Rs. 3000 the principal is	20,000	3,000	12,500	10,000			12,500
The formula for compound interest is	PNR /100	$P(1+r/100)^{n} - P$	A/(1+i)n	CI /(I+I)n			$P(1+r/100)^{n}-P$
The compound interest on Rs. 1000 at 12 % for one year is	Rs.150	Rs.175	Rs.120	Rs.126			Rs.120
The formula for calculating principal under compound interest. is	A /(1+i) <sup>n</sup>	$P(1+i)^n$	Pni	CI /(1+i) <sup>n</sup>			A /(1+i) <sup>n</sup>
If the principal isRs.5000 ,Interest is Rs.100 then amount A=	Rs.5100	Rs.4900	Rs.5000	Rs.6000			Rs.5100
If Rs.6,000 amounts to Rs.8,940 then the interest is	2940	Rs.3450	Rs.4560	Rs.3540			2940
The simple interest on Rs.6,000 at 10% for 2 years is	Rs.1200	Rs.1000	Rs.2000	Rs.1250			Rs.1200

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
The formula for simple interest is	Pni	P-n-r/100	P+n+r/100	Pn			Pni
The formula for calculating amount under compound interest is	$P(1+r/100)^{n}$	P(1+r/100)n-1	Pni	Р			P(1+r/100) <sup>n</sup>
Compound interest is more than	Principal	amount	simple interest	compound interes	t		simple interest
Formula for rate of interest under compound interest is	100[(A/P) <sup>1/n</sup> -1]	P(1+A/100)n-1	P(1+I)n	P(1+I)			100[(A/P) <sup>1/n</sup> -1]
In compound interest the formula for period is	log A-log P/log(	log A-log P/(1+r	log A+log P/(1+r/	A+P			log A-log P/log(1+r/
Compound interest for Rs.2,500 for 4 years at 8% per annum is	Rs.801.32	Rs.701.22	Rs.601.42	Rs.901.22			Rs.901.22
The effective rate of compound interest is	100[(1-r/100m) <sup>m</sup>	100[(1+r/100m)	$P(1+i)^n$	Р			$100[(1+r/100m)^{m}-1]$
In the effective rate of interest, a unit of time m is quarterly means m=	4	2	3	1			4
In the effective rate of interest, a unit of time m is half yearly means m=	4	2	3	1			2
Interest is compounded continuously means m is	large	small	middle	first			large
When interest is compounded continuously A is	Pe <sup>in</sup>	Pe-in	- Pe <sup>in</sup>	Р			Pe <sup>in</sup>
When interest is compounded continuously P is	Pe <sup>in</sup>	Pe-in	- Pe <sup>in</sup>	Р			Pe-in
When interest is compounded continuously, effective rate of interest S is	Pe <sup>in</sup>	A/ e <sup>in</sup>	- Pe <sup>in</sup>	$100(e^{i}-1)$			100(e <sup>i</sup> -1)

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
In what time a sum of Rs.1234 amount to Rs.5678 at 8% p.a. compound inter	18.72 years	19.27 years	19.72 years	17.95 years			19.27 years
Under compound interest, interest earns	interest	principal	amount	discount			interest
The simple interest on a sum of Rs.2000 for two years at 5% p.a. is	Rs.800	Rs.700	Rs.1000	Rs.1200			Rs.1200
If the simple interest is Rs 150 ,number of years is 6 and the rate if interest is	Rs. 500	Rs. 300	Rs.1000	Rs. 500			Rs. 500
If the simple interest is Rs 72, the rate if interest is 4 % p.a the principal is	3	4	5	1			3
Compound interest is	Amount - princip	Amount + princip	principal –Amou	Amount x principa	al		Amount - principal
The amount of money deposited or borrowed is called the	principal	interest	amount	discount			principal
The extra money which we get or which we pay is called	principal	interest	amount	discount			interest
The simple interest on a sum of Rs.1225 for 4 years at 11% p.a. is	Rs.539	Rs.700	Rs.100	Rs.120			Rs.539

ne value

100)

.]

# **KARPAGAM ACADEMY OF HIGHER EDUCATION** CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 UNIT: II BATCH-2018-2021 <u>UNIT – II</u> **SYLLABUS Differential Calculus** Mathematical functions and their types - linear, quadratic, polynomial; Concepts of limit and continuity of a function; Concept of differentiation; Rules of differentiation - simple standard forms. Applications of differentiation - elasticity of demand and supply; Maxima and Minima of functions (involving second or third order derivatives) relating to cost, revenue and profit. Prepared by : Dr. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE Page 1/21

CLASS: I B.B.A. COURSE CODE: 18BAU202

#### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: II

BATCH-2018-2021

### **Concept of Functions**

Let A and B be any two non-empty sets. Then a function 'ff' is a rule or law which associates each element of 'A' to a unique element of set 'B'.

#### Notation:

(i) A function is usually denoted by small letters, i.e. f,g,hf,g,h etc. and Greek letters,

i.e.  $\alpha, \beta, \gamma, \phi, \psi \alpha, \beta, \gamma, \phi, \psi$ etc.

(ii) If 'ff' is a function from 'A' to 'B' then we write  $f:A \rightarrow Bf:A \rightarrow B$ .

#### **Ordered Pair:**

Let 'a' and 'b' be any two elements then an element (a,b)(a,b) is called an ordered pair.

#### **Constant Function:**

Let 'A' and 'B' be any two non-empty sets, then a function 'ff' from 'A' to 'B' is called a constant function if and only if the range of 'ff' is a singleton.

#### **Algebraic Function:**

A function defined by an algebraic expression is called an algebraic function.

e.g.  $f(x)=x_2+3x+6f(x)=x_2+3x+6$ 

#### **Polynomial Function:**

A function of the form  $P(x) = a_m x_n + a_{n-1} x_{n-1} + \dots + a_{1x} + a_{0} P(x) = a_m x_n + a_{n-1} x_{n-1} + \dots + a_{1x} + a_{0} P(x)$ 

where 'n' is a positive integer and  $a_n, a_{n-1}, \dots, a_1, a_{0}a_{n-1}, \dots, a_{1,a_0}a_{n-1}$  are real numbers is called a polynomial function of degree 'n'.

#### **Linear Function:**

A polynomial function with degree 'tt' is called a linear function. The most general form of a linear function is

f(x)=ax+bf(x)=ax+b

#### **Quadratic Function:**

A polynomial function with degree '2' is called a quadratic function. The most general form of a quadratic equation is  $f(x)=ax_2+bx+c_1(x)=ax_2+bx+c_2(x)=ax_2+bx$ 

#### **Cubic Function:**

A polynomial function with degree '3' is called a cubic function. The most general form of a cubic 

#### **Identity Function:**

Let  $f:A \rightarrow Bf:A \rightarrow B$  be a function then 'ff' is called an identity function if  $f(x)=x, \forall x \in Af(x)=x, \forall x \in A$ . **Rational Function:** 

A function R(x)R(x) defined by R(x)=P(x)Q(x)R(x)=P(x)Q(x), where both P(x)P(x) and Q(x)Q(x) are polynomial functions is called a rational function.

	KARPAGAM ACADEMY OF HIGHER EDUCAT	ΓΙΟΝ
CLASS: I B.B.A. COURSE CODE: 18E	COURSE NAME:BUSINESS MATHEM BAU202 UNIT: II	ATICS AND STATISTICS BATCH-2018-2021
function. Exponential Fundamential A function in whe e.g. (i) $f(x)=axf$	= sinxf(x) = sin x, f(x) = cosxf(x) = cos x etc., then f nction: ich the variable appears as an exponent (power) is cal i(x)=ax (ii) f(x)=3xf(x)=3x. Logarithmic Function: n which the variable appears as an argument of a logarithm	led an exponential function
	e.g. $f(x) = loga(x)f(x) = loga(x)$ .	
	LIMIT AND CONTINUIT	ΓY
Limit of a Func	tion at Infinity	
Definition Let $> 0$ such that when	et f(x) be a function defined on R. $\lim_{x \to \infty} f(x) = \ell$ mean en x > X, $ f(x) - \ell  < \varepsilon$ .	s that for any $\epsilon > 0$ , there exists X
Theorem	UNIQUENESS of Limit Value	
	If $\lim_{x\to\infty} f(x) = a$ and $\lim_{x\to\infty} f(x) = b$ , then $a = b$ .	
	$x \rightarrow \infty$ $x \rightarrow \infty$	
Theorem	Rules of Operations on Limits	
	If $\lim_{x\to\infty} f(x)$ and $\lim_{x\to\infty} g(x)$ exist, then	
	(a) $\lim_{x \to \infty} [f(x) \pm g(x)] = \lim_{x \to \infty} f(x) \pm \lim_{x \to \infty} f(x) = \lim_{x \to \infty$	$\max_{\sigma \to \infty} g(x)$
	(b) $\lim_{x \to \infty} f(x)g(x) = \lim_{x \to \infty} f(x) \cdot \lim_{x \to \infty} g(x)$	x)
	(c) $\lim_{x \to \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \to \infty} f(x)}{\lim_{x \to \infty} g(x)} \text{ if } \lim_{x \to \infty} g(x) =$	≠ 0.

c) 
$$\lim_{x\to\infty}\frac{f(x)}{g(x)} = \frac{\lim_{x\to\infty}f(x)}{\lim_{x\to\infty}g(x)} \text{ if } \lim_{x\to\infty}g(x) \neq 0.$$

CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS							
COURSE CODE: 18BAU202		UNIT: II	BATCH-2018-2021				
	(d)	For any constant k, $\lim_{x \to \infty} [kf(x)] = k \lim_{x \to \infty} f(x)$ .					
	(e)	For any positive integer n,					
		(i) $\lim_{x \to \infty} [f(x)]^n = [\lim_{x \to \infty} f(x)]^n$					
		(ii) $\lim_{x \to \infty} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \to \infty} f(x)}$					
		DERIVATIVES					

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}$$

provided that the limit exists.

If we denote y = f(x), then f(a) is called the derivative of f, with respect to (the independent variable) x, at the point x = a.

Recall that the value of this limit is, if it exists, is the slope of the line tangent to the curve y = f(x) at the point x = a. As well, it also represents the instantaneous rate of change, with respect to x, of the function f ata. Therefore, a positive f'(a) means that the function f is *increasing* at a, while a negative f'(a) means that f is *decreasing* at a. If f(a) = 0, then f is neither increasing nor decreasing at a.

Equivalently, the derivative can be stated as

$$f'(a) = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

*ex.* Let  $f(t) = t^5 + 6t$ , find f'(a).

Prepared by : Dr. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

Page 4/21

### KARPAGAM ACADEMY OF HIGHER EDUCATION CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 BATCH-2018-2021 **UNIT: II** $f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \to 0} \frac{[(a+h)^5 + 6(a+h)] - [a^5 + 6a]}{h}$ $=\lim_{h\to 0}\frac{[a^5+5a^4h+10a^3h^2+10a^2h^3+5ah^4+h^5+6a+6h]-[a^5+6a]}{h}$ $=\lim_{h\to 0}\frac{5a^4h+10a^3h^2+10a^2h^3+5ah^4+h^5+6h}{h}$ $=\lim_{h\to 0}\frac{h(5a^4+10a^3h+10a^2h^2+5ah^3+h^4+6)}{h}$ $= \lim (5a^4 + 10a^3h + 10a^2h^2 + 5ah^3 + h^4 + 6) = 5a^4 + 6$ ex. Let $f(x) = \sqrt{4x^2 + 5}$ , find f(a). Write an equation of the line tangent to y = f(x)

when a = 1.

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \to 0} \frac{\sqrt{4(a+h)^2 + 5} - \sqrt{4a^2 + 5}}{h}$$
$$= \lim_{h \to 0} \frac{\sqrt{4(a+h)^2 + 5} - \sqrt{4a^2 + 5}}{h} \cdot \frac{\sqrt{4(a+h)^2 + 5} + \sqrt{4a^2 + 5}}{\sqrt{4(a+h)^2 + 5} + \sqrt{4a^2 + 5}}$$

$$=\lim_{h\to 0}\frac{4(a+h)^2+5-(4a^2+5)}{h(\sqrt{4(a+h)^2+5}+\sqrt{4a^2+5})}=\lim_{h\to 0}\frac{4a^2+8ah+4h^2+5-4a^2-5}{h(\sqrt{4(a+h)^2+5}+\sqrt{4a^2+5})}$$

CLASS: I B.B.A. COURSE CODE: 18BAU202

#### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: II BATCH-2018-2021

$$=\lim_{h\to 0}\frac{8ah+4h^2}{h\left(\sqrt{4(a+h)^2+5}+\sqrt{4a^2+5}\right)}=\lim_{h\to 0}\frac{8a+4h}{\sqrt{4(a+h)^2+5}+\sqrt{4a^2+5}}$$

$$=\frac{8a+4(0)}{\sqrt{4(a+0)^2+5}+\sqrt{4a^2+5}}=\frac{8a}{2\sqrt{4a^2+5}}=\frac{4a}{\sqrt{4a^2+5}}$$

At a = 1, the point on the curve, (a, f(a)) is (1, 3), and the slope of the tangent line is  $f'(1) = \frac{4}{3}$ .

The equation of the line is, in point-slope form, therefore

$$y-3=\frac{4}{3}(x-1)$$

or, in slope-intercept form,

$$y = \frac{4}{3}x + \frac{5}{3}$$
.  
ex. Let  $f(x) = \frac{1}{\sqrt{x^2 + 3}}$ , find  $f(a)$ .

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \to 0} \frac{\frac{1}{\sqrt{(a+h)^2 + 3}} - \frac{1}{\sqrt{a^2 + 3}}}{h}$$

$$=\lim_{h \to 0} \frac{\frac{1}{\sqrt{(a+h)^2 + 3}} - \frac{1}{\sqrt{a^2 + 3}}}{h} \cdot \frac{\sqrt{(a+h)^2 + 3}\sqrt{a^2 + 3}}{\sqrt{(a+h)^2 + 3}\sqrt{a^2 + 3}}$$

Prepared by : Dr. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

Page 6/21

KARPAGAM ACADEMY OF HIGHER EDUCATION           CLASS: I B.B.A.         COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS						
COURSE CODE: 18BAU202	UNIT: II	BATCH-2018-2021				
$= \lim_{h \to 0} \frac{\sqrt{a^2 + 3} - \sqrt{(a + h)^2}}{h(\sqrt{(a + h)^2 + 3})(\sqrt{a + h)^2}}$	$(a)^2 + 3$ $(a^2 + 3)$					
$= \lim_{h \to 0} \frac{\sqrt{a^2 + 3} - \sqrt{(a + h)^2}}{h(\sqrt{(a + h)^2 + 3})(\sqrt{a^2 + 3})}$	$\frac{(a)^2 + 3}{(a^2 + 3)} \cdot \frac{\sqrt{a^2 + 3} + \sqrt{(a + h)^2 + 3}}{\sqrt{a^2 + 3} + \sqrt{(a + h)^2 + 3}}$					
$= \lim_{h \to 0} \frac{(a^2 + 3)}{h(\sqrt{(a+h)^2 + 3})(\sqrt{a^2 + 3})}$	$\frac{(-(a^{2}+2ah+h^{2}+3))}{\sqrt{a^{2}+3}(\sqrt{a^{2}+3}+\sqrt{(a+h)^{2}+3})}$					
$=\lim_{h\to 0}\frac{1}{h(\sqrt{(a+h)^2+3})(\sqrt{a+b^2+3})}$	$\frac{-2ah-h^2}{\sqrt{a^2+3}}(\sqrt{a^2+3}+\sqrt{(a+h)^2+3})$					
$=\lim_{h\to 0}\frac{1}{(\sqrt{(a+h)^2+3})(\sqrt{a})}$	$\frac{-2a-h}{a^2+3}(\sqrt{a^2+3}+\sqrt{(a+h)^2+3}) =$	$=\frac{-2a}{(\sqrt{a^2+3})(\sqrt{a^2+3})(2\sqrt{a^2+3})}$				
$=\frac{-a}{(a^2+3)^{3/2}}$	Resig Differentiation Form					

#### **Basic Differentiation Formulas**

Suppose f and g are differentiable functions, c is any real number, then

- 1.  $\frac{d}{dx}(c) = 0$ 2.  $\frac{d}{dx}[f(x)+g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$ 3.  $\frac{d}{dx}[f(x) - g(x)] = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$
- 4.  $\frac{d}{dx}[cf(x)] = c \cdot \frac{d}{dx}f(x)$

Page 7/21

COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS

CLASS: I B.B.A. COURSE CODE: 18BAU202

#### UNIT: II

BATCH-2018-2021

5. 
$$\frac{d}{dx}[f(x)g(x)] = f(x)\frac{d}{dx}[g(x)] + g(x)\frac{d}{dx}[f(x)]$$

6. 
$$\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{g(x)\frac{d}{dx}[f(x)] - f(x)\frac{d}{dx}[g(x)]}{[g(x)]^2}$$

Or, in short-hand notations:

1. (c)' = 0

2. 
$$[f(x) + g(x)]' = f'(x) + g'(x)$$

- 3. [f(x) g(x)]' = f'(x) g'(x)
- 4. [cf(x)]' = cf'(x)
- 5. [f(x)g(x)]' = f(x)g'(x) + g(x)f'(x)

6. 
$$\left[\frac{f(x)}{g(x)}\right] = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$$

The Power Rule: For any real number *n*,

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

For n = 1, this means that  $\frac{d}{dx}(x) = 1 \cdot x^{1-1} = x^0 = 1$ . And if n = 0, then  $\frac{d}{dx}(x^0) = \frac{d}{dx}(1) = 0 \cdot x^{0-1} = 0$ , which is consistent with the constant rule of differentiation (rule #1 above).

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IIBATCH-2018-2021

ex. The instantaneous rate of change of a line

Suppose f(x) = mx + b, where *m* and *b* are constants, then

$$f'(x) = (mx)' + (b)' = m(x)' + (b)' = m(1) + 0 = m$$

Therefore, any linear function has a constant derivative equals to the slope of its graph, which is a line of slope m. It says that the instantaneous rate of change of a linear function is constant, and that the tangent line to the graph of a line is always the line itself (because the tangent line has the same slope as the line, and they obviously contain one common point, therefore they have the same equation and are therefore the same line).

ex. Differentiate  $y = 2t^3 - t^{\pi} + t^{-2} + 9$   $y' = 2(t^3)' - (t^{\pi})' + (t^{-2})' + (9)' = 2(3t^2) - \pi t^{\pi - 1} + (-2t^{-3}) + 0$   $= 6t^2 - \pi t^{\pi - 1} - 2t^{-3}$ ex. Differentiate  $s(t) = 5\sqrt{t} - \frac{2}{\sqrt{t}} + 4\sqrt[3]{t}$ 

This would be easier to do if we first rewrite s(t) in terms of powers of x.

$$s(t) = 5t^{1/2} - 2t^{-1/2} + 4t^{1/3}$$
, then

$$s'(t) = 5\left(\frac{1}{2}t^{(\frac{1}{2}-1)}\right) - 2\left(\frac{-1}{2}t^{(\frac{-1}{2}-1)}\right) + 4\left(\frac{1}{3}t^{(\frac{1}{3}-1)}\right) = \frac{5}{2}t^{-\frac{1}{2}} + t^{-\frac{3}{2}} + \frac{4}{3}t^{-\frac{2}{3}}$$

KARPA CLASS: I B.B.A. COURSE CODE: 18BAU202	AGAM ACADEMY OF HIC COURSE NAME:BUSIN UNIT: II		AND STATISTICS BATCH-2018-2021
ex. Differentiate $f(x)$	$) = \frac{x^2 + 3x + 2}{x^2 - 3x + 2}$		
$f'(x) = \frac{(x^2 - 3x + 2)(x)}{(x^2 - 3x + 2)(x)}$	$\frac{x^2 + 3x + 2)' - (x^2 + 3x + 2)'}{(x^2 - 3x + 2)^2}$	$2)(x^2-3x+2)'$	
$=\frac{(x^2-3x+2)(2x+3)}{(x^2-3)(x^2-3)(x^2-3)(x^2-3)}$	$\frac{(x^2 + 3x + 2)(2x - 3)}{(-3x + 2)^2}$		
ex. Differentiate $g(x)$	$) = \frac{2x^3 + 3x^2 - x + 5}{x^2}$		
The easiest way to do	this is to rewrite $g(x)$ as		
$g(x) = 2x + 3 - \frac{1}{x} + \frac{5}{x^2}$	, then		
g'(x) = 2(x)' + (3)' - (3)'	$(x^{-1})' + 5(x^{-2})' = 2 + 0 - (-1)^{-1}$	$-x^{-2})+5(-2x^{-3})$	
$= 2 + x^{-2} - 10x^{-3} = 2$	$+\frac{1}{w^2}-\frac{10}{w^3}$		
<i>ex.</i> Differentiate $y =$			
Simplify first: $y = x^{5/2}$	$-5x^{3/2}+2x^{1/2}$ .		
$y' = \frac{5}{2}x^{3/2} - 5\left(\frac{3}{2}x^{1/2}\right) $	$+2\left(\frac{1}{2}x^{-1/2}\right) = \frac{5}{2}x^{3/2} - \frac{15}{2}$	$x^{1/2} + x^{-1/2}$	
	×		

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IIBATCH-2018-2021

The longer way to do this is by using the product rule:

$$y' = \sqrt{x} (x^2 - 5x + 2)' + (x^2 - 5x + 2)(\sqrt{x})' = \sqrt{x} (2x - 5) + (x^2 - 5x + 2)(\frac{1}{2}x^{-1/2})$$

$$=x^{1/2}(2x-5)+(x^2-5x+2)(\frac{1}{2}x^{-1/2})=2x^{3/2}-5x^{1/2}+\frac{1}{2}x^{3/2}-\frac{5}{2}x^{1/2}+x^{-1/2}$$

$$=\frac{5}{2}x^{3/2}-\frac{15}{2}x^{1/2}+x^{-1/2}$$

#### Derivatives of logarithmic and exponential functions

**Formula**. The exponential rule. For the function of the type  $f(x) = a^x$ , where a is a constant, then  $f'(x) = a^x \ln a$ . Derivative of  $f(x) = e^x$  is  $f'(x) = e^x$ .

Example 1.Find  $\frac{dy}{dx}$ . a.  $f(x) = 3x^{20} + 10^x$  b.  $f(x) = e^x + e^{5x}$ Solution.a.  $\frac{df}{dx} = 60x^{19} + 10^x \ln 10$ , b.  $\frac{df}{dx} = e^x + 5e^{5x}$ Formula. For the logarithmic function  $f(x) = \ln x$ ,  $f'(x) = \frac{1}{x}$ . Derivative of  $f(x) = \log_a x$  is  $f'(x) = \frac{1}{x \ln a}$ . Example 2.Find  $\frac{dy}{dx}$ . a.  $f(x) = 3\ln x + 10$  b.  $f(x) = \log x + 3$ Solution.a.  $\frac{df}{dx} = \frac{3}{x}$ , b.  $\frac{df}{dx} = \frac{1}{x \ln 10}$ Example 1.Find  $\frac{dy}{dx}$  or y'. a.  $3y^2 - 5xy + 9x - 2 = 0$  b.  $y^2 = \log x + 3$ Solution.a.  $6yy' - 5y - 5xy' + 9 = 0 \Rightarrow y' = \frac{5y - 9}{6y - 5x}$ b.  $2yy' = \frac{1}{x \ln 10} \Rightarrow y' = \frac{1}{2xy \ln 10}$ 

## KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IIBATCH-2018-2021

*Example 2.*Find the slope of the tangent line to the curve  $3xy - 2x^2 = 7$  at (1, 3). Also compute the second derivative.

Solution.  $3xy'+3y-4x = 0 \Rightarrow y' = \frac{4x-3y}{3x} = \frac{4(1)-3(3)}{3(1)} = -\frac{5}{3}$ For the second derivative we consider again 3xy'+3y-4x = 0Taking derivative we find  $3xy''+3y'+3y'-4 = 0 \Rightarrow y'' = \frac{4-6y'}{3x} = \frac{6y-4x}{3x^2} = \frac{14}{3}$ 

#### **DEMAND AND SUPPLY**

#### The market and its economic agents

Purpose of this lesson: to study the behaviour of people as they interact with one another in markets.

Market: a group of buyers and sellers of a particular good (or service).

Demand: represents the behaviour of buyers.

Supply: represents the behaviour of sellers.

Main assumption that we will use for the time being: markets are competitive.

<u>Characteristics</u> of a competitive market:

- a) Goods offered for sale in one market are all the same.
- b) Buyers and sellers are so numerous, that they individually cannot influence the market price. We say that both buyers and sellers are price takers.

Not all markets are competitive. If sellers are few and individually can influence the market price, then we say that markets are not competitive (<u>oligopoly</u>, <u>monopoly</u>).

#### The concepts of demand and supply

To be specific let us concentrate on one particular market: for example, the market for butter.

COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 UNIT: II BATCH-2018-2021

#### Demand

CLASS: I B.B.A.

Factors that determine the quantity demanded of butter:

- The price of the good. The higher the price, the lower the quantity demanded. This is the Law of a) demand.
- b) <u>Income</u>. Normally, the richer people are, the more of a good they will buy. Let us call income *m*.
- c) Prices of related goods.

x and y substitutes. Think of butter (x) and margarine (y).

 $p_{y} \downarrow \Rightarrow y \uparrow \Rightarrow x \downarrow$ ; Therefore  $p_{y}$  and x positively related.

x and y complements. Think of butter (x) and bread (y)

 $p_v \downarrow \Rightarrow y \uparrow \Rightarrow x \uparrow$ ; Therefore  $p_v$  and x negatively related.

d) Tastes. Some people like butter, others like olive oil.

#### The demand curve

Suppose that we give a mathematical form to the relationship between butter (x) and the factors that determine the quantity demanded of butter.

$$x = 10 - 4p_x + 0.005m + 2p_y$$

Discuss signs.

Normally we do not want to work with so many variables. The ones we are interested in depend on the problem at hand. Suppose we are only interested in the price and quantity of butter, and that we are given

the values of the other variables  $m = 3,000 \in; p_y = 0.5 \in$ . Then,

 $x = 10 - 4p_x + 0.005(3,000) + 2(0.5)$ 

 $x = (10 + 15 + 1) - 4p_{r}$ 

$$x = 26 - 4p_x$$

## KARPAGAM ACADEMY OF HIGHER EDUCATION CLASS: I B.B.A. COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 UNIT: II BATCH-2018-2021

The expression in the box is the <u>demand curve</u> (or the <u>demand function</u>). Suppose we have a market with three buyers: Ana, Víctor and Pilar. Each has the following demand curve:

Ana:  $x_A = 10 - 2p_x$ Víctor:  $x_V = 20 - 4p_x$ Pilar:  $x_P = 15 - 3p_x$ 

We obtain the aggregate demand curve by adding horizontally (adding the x's).

$$x = x_A + x_V + x_P$$
  
= (10-2p<sub>x</sub>)+(20-4p<sub>x</sub>)+(15-3p<sub>x</sub>)  
= 45-9p<sub>x</sub>

Therefore, the aggregate demand curve is

$$x = 45 - 9p_x$$

2. A consumer buys 100 units of a product when the price is  $\in 1$ . When the price is increased to  $\in 1.25$  the consumer buys 80 units. Calculate the Price Elasticity of demand for this consumer.


$\frac{P1 + P2}{Q1 + Q2}$	×	$\frac{\Delta Q}{\Delta P}$
$\frac{1+1.25}{100+80}$	x	<u>-20</u> +0.25
= -1		

Demand for this good is unitary elastic as the answer is equal to 1

#### MAXIMA AND MINIMA

You are aware that in any transaction the total amount paid increases with the number of items purchased. Consider a function as f(x) = 2x + 1, x > 0. Let the function f(x) represent the amount required for purchasing 'x' number of items.


The graph of the function y = f(x) = 2x + 1, x > 0 for different values of x is shown in



A look at the graph of the above function prompts us to believe that the function is increasing for positive values of x i.e., for x>0 Can you think of another example in which value of the function decreases when x increases ? Any such relation would be relationship between time and manpower/person(s) involved. You have learnt that they are inversely proportional. In other words we can say that time required to complete a certain work increases when number of: persons (manpower) involved decreases and vice-versa. Consider such similar function as

$$g(x) = \frac{2}{x}, x > 0$$

The values of the function g (x) for different values of x are plotted



CLASS: I B.B.A. COURSE CODE: 18BAU202

#### COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS UNIT: II BATCH-2018-2021


All these examples, despite the diversity of the variables involved, have one thing in common : function is either increasing or decreasing.

#### INCREASING AND DECREASING FUNCTIONS

Let a function f(x) be defined over the closed interval [a,b].

Let  $x_1, x_2 \in [a, b]$ , then the function f(x) is said to be an increasing function in the given interval. if  $f(x_2) \ge f(x_1)$  whenever  $x_2 > x_1$ . It is said to be strictly increasing if  $f(x_2) > f(x_1)$  for all.  $x_2 > x_1, x_1, x_2 \in [a, b]$ .

In Fig. 25.3, sin x increases from -1 to +1 as x increases from  $-\frac{\pi}{2}$  to  $+\frac{\pi}{2}$ .




Note : A function is said to be an increasing function in an interval if f(x + h) > f(x) for all x belonging to the interval when h is positive.

A function f(x) defined over the closed interval [a, b] is said to be a decreasing function in the given interval, if  $f(x_2) \le f(x_1)$ , whenever  $x_2 > x_1$ ,  $x_1, x_2 \in [a, b]$ . It is said to be strictly decreasing if  $f(x_1) > f(x_2)$  for all  $x_2 > x_1$ ,  $x_1, x_2 \in [a, b]$ .

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: IIBATCH-2018-2021

In Fig. 25.4,  $\cos x$  decreases from 1 to -1 as x increases from 0 to  $\pi$ .





Note: A function is said to be a decreasing in an internal if f(x+h) < f(x) for all x belonging to the interval when h is positive.

#### Example

Prove that the function f(x) = 4x + 7 is monotonic for all values of  $x \in \mathbb{R}$ .

**Solution :** Consider two values of x say  $x_1, x_2 \in \mathbb{R}$ 

such that

 $x_2 > x_1$  .....(1)

Multiplying both sides of (1) by 4, we have  $4x_2 > 4x_1$ 

Adding 7 to both sides of (2), to get

 $4x_2 + 7 > 4x_1 + 7$  $f(x_2) > f(x_1)$ 

We have

Prepared by : Dr. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

.....(2)

CLASS: I B.B.A. URSE CODE: 18		ME:BUSINESS MATHEMATI UNIT: II	BATCH-2018-20
Thus, we find f	$f(x_2) > f(x_1)$ wheneve	$x_{1} = x_{1}$	
Hence the giver	function $f(x) = 4x + 7$	7 is monotonic function. (mono	otonically increasing).
Example			
Show that			
f(x) = x	2		
is a strictly decr	reasing function for all x	< 0.	
Solution : Con	sider any two values of a	x say x1, x2 such that	<i>©</i> .
	$x_2 > x_1$ ,	$x_1, x_2 < 0$	(i)
Order of the ine	equality reverses when it i	is multiplied by a negative num	ber. Now multiplying (i)
by x <sub>2</sub> , we hav	re		
	$x_2 \cdot x_2 < x_1 \cdot x_2$	2	
or,	$x_2^2 < x_1 x_2$		(ii)
Now multiplyin	ng (i) by x <sub>1</sub> , we have		
	$x_1 \cdot x_2 < x_1 \cdot x_1$	1	
or,	$x_1 x_2 < x_1^2$		(iii)
From (ii) and (i	ii), we have		
	$x_2^2 < x_1 x_2 < x_1$	2	
OT,	$x_{1}^{2} < x_{1}^{2}$		
	2 1	)	
or,	$f(x_2) < f(x_1)$	)	
1 mus, from (1)	and (iv), we have for		
	$x_2 > x_1,$	×	
	$f(x_2) < f(x_1)$	)	

CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 BATCH-2018-2021 **UNIT: II** 

#### Example

Find the interval in which  $f(x) = 2x^3 - 3x^2 - 12x + 6$  is increasing or decreasing.  $f(x) = 2x^3 - 3x^2 - 12x + 6$ Solution :

)

$$f'(x) = 6x^{2} - 6x - 12$$
$$= 6(x^{2} - x - 2)$$
$$= 6(x - 2)(x + 1)$$

For f(x) to be increasing function of x,

i.e.

f'(x) > 06(x-2)(x+1) > 0 or, (x-2)(x+1) > 0

Since the product of two factors is positive, this implies either both are positive or both are negative.

Either

OT.

x - 2 > 0 and x + 1 > 0x - 2 < 0 and x + 1 < 0or i.e. x < 2 and x < 1i.e. x > 2 and x > -1x > 2 implies x > -1x < -1 implies x < 2x > 2∴ x <-1 ÷.,

Hence f(x) is increasing for  $x \ge 2$  or  $x \le -1$ .

Now, for f (x) to be decreasing,

 $f'(x) \le 0$ 6(x-2)(x+1) < 0(x-2)(x+1) < 0Or,

Since the product of two factors is negative, only one of them can be negative, the other positive. Therefore.

		or	
	-2 > 0 and x	+ 1< 0	x - 2 < 0 and $x + 1 > 0$
i.e. x	>2 and x <-	1	i.e. $x \le 2$ and $x \ge -1$
T	here is no such	possibility	This can be put in this form
th	at x > 2 and at	the same time	
х	<-1		$-1 \le x \le 2$
∴ The fu	nction is decre	asing in $-1 \le x \le 2$ .	
Example			
Determine		or which the function	$(x) = \frac{x}{x^2 + 1}$ is increasing or decreasing. $(x) = \frac{x}{x^2 + 1}$
Solution	: f '(x):	$=\frac{\left(x^{2}+1\right)\frac{dx}{dx}-x\frac{d}{dx}\left(x^{2}+1\right)^{2}}{\left(x^{2}+1\right)^{2}}$	<u> </u>
		$=\frac{(x^{2}+1) - x (2x)}{(x^{2}+1)^{2}}$	
		$=\frac{1-x^2}{\left(x^2+1\right)^2}$	
ä	f'x =	$\frac{(1-x)(1+x)}{\left(x^2+1\right)^2}$	
As $(x^2 +$	$(-1)^2$ is positive	for all real x.	
Therefore	e, if -1 < x <0	(1 - x) is positive and (	(1+x) is positive, so f '(x) > 0;

:. If 0 < x < 1, (1 + x) is positive and (1+x) is positive, so f f '(x) > 0;

COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS

COURSE CODE: 18BAU202

CLASS: I B.B.A.

UNIT: II

BATCH-2018-2021

#### **POSSIBLE QUESTIONS 2 MARK QUESTIONS**

1. What are the types of Functions?

2. Define derivative.

3. If  $f(x)=(x^2-7)^2$ , find f'(x).

4. Find the derivative of  $x^3-3x^2+4x+3$ 

5. Define limit of a function.

6. Find  $\lim_{x \to 1} \frac{x^3 + 1}{2x^2 + 5x = 3}$ 

7. Define elasticity of supply.

8. Define elasticity of demand.

9. Find  $\frac{dy}{dx}$ , if  $y = (2x + 5)^3$ 

#### **8 MARK QUESTIONS**

- 1.A firm sells a product at Rs.3 per unit. The total cost of the firm for producing x units is given by  $C=20+0.6x+0.001x^2$ . How many units should be made to achieve maximum profit? Verify that the condition for a maximum is satisfied.
- 2. If the demand law is  $x = \frac{20}{p+1}$  find the elasticity of demand at the point when p=3.
- **3.** Find the elasticity of supply from the function p = -2+5x.
- 4. Find  $\frac{dy}{dx}$  i)  $x^2 + y^2 = 1$  ii)  $xy = c^2$

5.If  $f(x) = \frac{x^3 - 2x^2 + 50}{x^2}$  find f'(5) and f'(10). 6.Differentiate the following with respect to x (i)  $x^3 - 3x^2 + 4x + 3$  (ii)  $x^5 + 3\log x - 4e^x$ 

7.Differentiate the following with respect to x

(i) 
$$y = (x^2 + 5)(3x + 1)$$
 (ii)  $y = \frac{3x^2}{4x - 1}$   
8.Findlim <sub>$x \to 0$</sub>   $\frac{(1+x)^4 - 1}{(1+x)^2 - 1}$   
9. Find  $\lim_{x \to 0} \frac{1 - \sqrt{1-x^2}}{x^2}$   
10. Find  $\lim_{x \to 3} \frac{x^2 + x - 12}{x^2 - x - 6}$ 

- 11. If the demand law is  $x = \frac{20}{p+1}$ , find the elasticity of demand at the point when p=3.
- 12. Find for what values of x, the following expression is maximum and minimum respectively.  $2x^3 - 21x^2 + 36x - 20$  find also the maximum and the minimum values.

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
Quantities which take the same values throughout particular investigation are called	Constants	Variables	Functions	Integers			Constants
Quantities which change that is different values in an investigation are called	Constants	Variables	Functions	Integers			Variables
The gross profit ddepends on the sales and the volume of a sphere depends on its	Circle	Square	Radius	Diameter			Radius
Two variables x and y are said to be function of each other when the values of one of them on those of ther. other	Independent	propotional	depend	perpendicular			depend
If corresponding to each value of x, there is only one value of y then y is calledValued	Single	many	two	three			Single
If corresponding to each value of x, there existsmore than one value of y then y is calledValued	Single	many	two	three			many
The relation between two variables can be expressed in the form $y=f(x)$ , y is	implicit	explicit	odd	even			explicit
The relation between two variables can be expressed in the formf(x,y)=0, y is	implicit	explicit	odd	even			implicit
If f(-x)=f(x) then f(x) is called	odd	even	implicit	explicit			even
If $f(-x)=-f(x)$ then $f(x)$ is called	odd	even	implicit	explicit			odd
y=a0+a1x+anxn is called function.	odd	even	implicit	polynomial			polynomial
Limit x tends to a f(x)=1 is function.	limit	even	implicit	polynomial			limit
Limit h tends to 0 f(a-h)=11 is limit	Right hand	Left hand	finite	infinite			Left hand

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
Limit h tends to 0 f(a+h)=11 is limit	Right hand	Left hand	finite	infinite			Right hand
If left hand limit = right hand limit then limit	does not exists	zero	exists	one			exists
If the limit is in indeterminant form then apply the method of	Induction	partial fraction	Integration	factorization			factorization
If the limit is in indeterminant form then apply the method of	rationalisation	Induction	partial fraction	Integration			rationalisation
If the limit is in indeterminant form then apply the method of	Induction	partial fraction	Substitution	Integration			Substitution
If the limit is in indeterminant form then apply the method of	Induction	partial fraction	Integration	L'Hospitals			L'Hospitals
If the limit is in indeterminant form then apply the method of	Induction	partial fraction	Integration	Infinite Limit			Infinite Limit
The rate of change of the object is called	Derivative	Induction	Integration	Velocity			Derivative
The process of finding derivatives are called	Integration	Differentiatio n	Induction	Acceleration			Differentiation
Derivative of exponential is	algebra	exponential	logerthms	function			exponential
Derivative of constant is	one	two	three	zero			zero
One of the rule of differentiation is	addition	commutative	assiciative	closure			addition
One of the rule of differentiation is	commutative	assiciative	closure	Difference			Difference

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
One of the rule of differentiation is	commutative	Product	closure	assiciative			Product
One of the rule of differentiation is	closure	assiciative	Quotient	commutative			Quotient
If x is function of t and y is function of t then it is in form.	closure	assiciative	Quotient	Parametric			Parametric
Process of finding second derivative ,third derivative etc is called	Initial	final	Successive	Single			Successive
One of the application of Differentiation is	Boundary	Elasticity	Simple Interest	Compound Interest			Elasticity
If the graph of the function rises then the function is function	Decreasing	Increasing	rising	falling			Increasing
If the graph of the function fallss then the function is function	Decreasing	Increasing	rising	falling			Decreasing
If the first dereivative zero and the second derivative less than zero then function is	Minima	Maxima	Optima	Extrima			Maxima
If the first dereivative zero and the second derivative greater than zero then function is	Minima	Maxima	Optima	Extrima			Minima

CLASS: I B.B.A. COURSE CODE: 18BAU202

#### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V BATCH-2018-2021

#### <u>UNIT – V</u>

#### **SYLLABUS**

#### Time-based Data: Index Numbers and Time-Series Analysis

Meaning and uses of index numbers; Construction of index numbers: Aggregative and average of relatives – simple and weighted, Tests of adequacy of index numbers, Construction of consumer price indices. Components of time series; additive and multiplicative models; Trend analysis: Finding trend by moving average method and Fitting of linear trend line using principle of least squares

KARPAGAM ACADEMY OF HIGHER EDUCATION						
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	IATHEMATICS AND STATISTICS				
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021				

#### **Index Numbers**

#### Introduction:

An index number is a statistical device for comparing the general level of magnitude of a group of related variables in two or more situation. If we want to compare the price level of 2000 with what it was in 1990, we shall have to consider a group of variables such as price of wheat, rice, vegetables, cloth, house rent etc., If the changes are in the same ratio and the same direction, we face no difficulty to find out the general price level. But practically, if we think changes in different variables are different and that too, upward or downward, then the price is quoted in different units i.e milk for litre, rice or wheat for kilogram, rent for square feet, etc

#### Uses of Index numbers

Index numbers are indispensable tools of economic and business analysis. They are particular useful in measuring relative changes. Their uses can be appreciated by the following points.

- 1. They measure the relative change.
- 2. They are of better comparison.
- 3. They are good guides.
- 4. They are economic barometers.
- 5. They are the pulse of the economy.
- 6. They compare the wage adjuster.
- 7. They compare the standard of living.
- 8. They are a special type of averages.
- 9. They provide guidelines to policy.
- 10. To measure the purchasing power of money.

KARPAGAM ACADEMY OF HIGHER EDUCATION					
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	IATHEMATICS AND STATISTICS			
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021			

Notation: For any index number, two time periods are needed for comparison. These are called the Base period and the Current period. The period of the year which is used as a basis for comparison is called the base year and the other is the current year. The various notations used are as given below:

 $P_1$  = Price of current year $P_0$  = Price of base year $q_1$  = Quantity of current year $q_0$  = Quantity of base year

#### 10.4 Problems in the construction of index numbers


No index number is an all purpose index number. Hence, there are many problems involved in the construction of index numbers, which are to be tackled by an economist or statistician. They are

- 1. Purpose of the index numbers
- 2. Selection of base period
- 3. Selection of items
- 4. Selection of source of data
- Collection of data
- 6. Selection of average
- 7. System of weighting

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS			
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021	

## 10.5 Method of construction of index numbers:

Index numbers may be constructed by various methods as shown below:



## 10.5.1 Simple Aggregate Index Number

This is the simplest method of construction of index numbers. The price of the different commodities of the current year are added and the sum is divided by the sum of the prices of those commodities by 100. Symbolically,

Simple aggregate price index =  $P_{01} = \frac{\sum p_1}{\sum p_0} \times 100$ 

Where ,  $\Sigma p_1$  = total prices for the current year  $\Sigma p_0$  = Total prices for the base year

## Example 1:

Calculate index numbers from the following data by simple aggregate method taking prices of 2000 as base.

## KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B.B.A. COURSE CODE: 18BAU202 COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V BATCH-2018-2021

Commodity	Price pe (in Ruj	
	2000	2004
A	80	95
В	50	60
С	90	100
D	30	45

#### Solution:

Commodity	Price pe (in Ru	
	2000	2004
А	(P <sub>0</sub> ) 80	(P <sub>1</sub> ) 95
B	50	60
С	90	100
D	30	45
Total	250	300

Simple aggregate Price index = 
$$P_{01} = \frac{\sum p_1}{\sum p_0} \times 100$$
  
=  $\frac{300}{250} \times 100$  =

## 10.5.2 Simple Average Price Relative index:

In this method, first calculate the price relative for the various commodities and then average of these relative is obtained by using arithmetic mean and geometric mean. When arithmetic mean is used for average of price relative, the formula for computing the index is

Prepared by: Dr. S. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

120

KARPAG	AM ACADEMY OF HIGHER ED	DUCATION
S: I B.B.A.	COURSE NAME: BUSINESS	MATHEMATICS AND STATISTICS
	TINITT, X7	DATCH 2040 2024

CLASS: I B.B.A. COURSE CODE: 18BAU202

UNIT: V BATCH-2018-2021

Simple average of price relative by arithmetic mean

$$P_{01} = \frac{\sum \left(\frac{p_1}{p_0} \times 100\right)}{n}$$

 $P_1 = Prices of current year$ 

 $P_0 =$  Prices of base year

n = Number of items or commodities

when geometric mean is used for average of price relative, the formula for obtaining the index is

Simple average of price relative by geometric Mean

$$P_{01} = \text{Antilog} \left( \frac{\sum \log(\frac{p_1}{p_0} \times 100)}{n} \right)$$

#### Example 2:

From the following data, construct an index for 1998 taking 1997 as base by the average of price relative using (a) arithmetic mean and (b) Geometric mean

Commodity	Price in 1997	Price in 1998
A	50	70
В	40	60
С	80	100
D	20	30

#### Solution:

#### (a) Price relative index number using arithmetic mean

Commodity	Price in 1997 (P <sub>0</sub> )	Price in 1998 (P <sub>1</sub> )	$\frac{p_1}{p_0} \times 100$
A	50	70	140
В	40	60	150
С	80	100	125
D	20	30	150
		Total	565

KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS			
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021	

Simple average of price relative by arithmetic mean

$$\mathbf{P}_{01} = \frac{\sum \left(\frac{p_1}{p_0} \times 100\right)}{n}$$

 $P_1 = Prices of current year$ 

 $P_0 =$  Prices of base year

n = Number of items or commodities

when geometric mean is used for average of price relative, the formula for obtaining the index is

Simple average of price relative by geometric Mean

$$P_{01} = \text{Antilog} \left( \frac{\sum \log(\frac{p_1}{p_0} \times 100)}{n} \right)$$

#### Example 2:

From the following data, construct an index for 1998 taking 1997 as base by the average of price relative using (a) arithmetic mean and (b) Geometric mean

Commodity	Price in 1997	Price in 1998
A	50	70
B	40	60
С	80	100
D	20	30

Solution:

#### (a) Price relative index number using arithmetic mean

Commodity	Price in 1997 (P <sub>0</sub> )	Price in 1998 (P <sub>1</sub> )	$\frac{p_1}{p_0} \times 100$
A	50	70	140
В	40	60	150
С	80	100	125
D	20	30	150
		Total	565

KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	ATHEMATICS AND STATISTICS		
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021		

Simple average of price relative by arithmetic mean

$$P_{01} = \frac{\sum \left(\frac{p_1}{p_0} \times 100\right)}{n}$$

 $P_1 = Prices of current year$ 

 $P_0 =$  Prices of base year

n = Number of items or commodities

when geometric mean is used for average of price relative, the formula for obtaining the index is

Simple average of price relative by geometric Mean

$$P_{01} = \text{Antilog} \left( \frac{\sum \log(\frac{p_1}{p_0} \times 100)}{n} \right)$$

#### Example 2:

From the following data, construct an index for 1998 taking 1997 as base by the average of price relative using (a) arithmetic mean and (b) Geometric mean

Commodity	Price in 1997	Price in 1998
A	50	70
B	40	60
С	80	100
D	20	30

#### Solution:

### (a) Price relative index number using arithmetic mean

Commodity	Price in 1997 (P <sub>0</sub> )	Price in 1998 (P <sub>1</sub> )	$\frac{p_1}{p_0} \times 100$
A	50	70	140
В	40	60	150
С	80	100	125
D	20	30	150
		Total	565

#### KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B.B.A. COURSE CODE: 18BAU202

Simple average of price relative index = (P<sub>01</sub>) =  $\sum \left( \frac{P_1}{P_0} \times 100 \right)$ 

$$= \frac{565}{4} = 141.25$$

#### (b) Price relative index number using Geometric Mean

Commodity	Price in 1997 (P <sub>0</sub> )	Price in 1998 (P <sub>1</sub> )	$\frac{p_1}{p_0}\times 100$	$\log(\frac{p_1}{p_0} \times 100)$
A	50	70	140	2.1461
В	40	60	150	2.1761
С	80	100	125	2.0969
D	20	30	1.50	2.1761
			Total	8.5952

Simple average of price Relative index

$$\Sigma \log \left[ \frac{p_1}{p_o} x 100 \right]$$

$$(P_{01}) = \text{Antilog} \qquad \begin{array}{c} n \\ 8.5952 \\ 4 \\ = \text{Antilog} \quad \begin{bmatrix} 2.1488 \end{bmatrix} = 140.9 \end{array}$$

#### 10.5.3 Weighted aggregate index numbers

In order to attribute appropriate importance to each of the items used in an aggregate index number some reasonable weights must be used. There are various methods of assigning weights and consequently a large number of formulae for constructing index numbers have been devised of which some of the most important ones are

- 1. Laspeyre's method
- 2. Paasche's method
- 3. Fisher's ideal Method
- 4. Bowley's Method
- 5. Marshall- Edgeworth method
- 6. Kelly's Method

KARPAG	AM ACADEMY OF HIGHER EDU	CATION
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	ATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021

#### 1. Laspeyre's method:

The Laspeyres price index is a weighted aggregate price index, where the weights are determined by quantities in the based period and is given by

Laspeyre's price index = 
$$P_{01}^{T} = \frac{\Sigma p_1 q_0}{\Sigma p_0 q_0} \times 100$$

#### 2. Paasche's method

The Paasche's price index is a weighted aggregate price index in which the weight are determined by the quantities in the current year. The formulae for constructing the index is

Paasche's price index number =  $P_{01}^{P} = \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1} \times 100$ 

Where

 $P_0 =$  Price for the base year  $P_1 =$  Price for the current year

 $q_0$  – Quantity for the base year  $q_1$  – Quantity for the current year **3. Fisher's ideal Method** 

Fisher's Price index number is the geometric mean of the Laspeyres and Paasche indices Symbolically

Fisher's ideal index number = 
$$P_{01}^{F} = \sqrt{L \times P}$$
  
=  $\sqrt{\frac{\Sigma p_1 q_0}{\Sigma p_0 q_0}} \times \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1} \times 100$ 

It is known as ideal index number because

(a) It is based on the geometric mean

(b) It is based on the current year as well as the base year

(c) It conform certain tests of consistency

(d) It is free from bias.

## 4. Bowley's Method:

Bowley's price index number is the arithmetic mean of Laspeyre's and Paasche's method. Symbolically

Bowley's price index number = 
$$P_{01}^{B} = \frac{L+P}{2}$$
  
=  $\frac{1}{2} \left[ \frac{\Sigma p_1 q_0}{\Sigma p_0 q_0} + \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1} \right] \times 100$ 

#### 5. Marshall- Edgeworth method

This method also both the current year as well as base year prices and quantities are considered. The formula for constructing the index is

Marshall Edgeworth price index = 
$$P_{01}^{ME} = \frac{\Sigma(q_0 + q_1)p_1}{\Sigma(q_0 + q_1)p_0} \times 100$$
  
=  $\frac{\Sigma p_1 q_0 + \Sigma p_1 q_1}{\Sigma p_0 q_0 + \Sigma p_0 q_1} \times 100$ 

#### 6. Kelly's Method

Kelly has suggested the following formula for constructing the index number

Kelly's Price index number = 
$$P_{01}^{k} = \frac{\Sigma p_1 q}{\Sigma p_0 q} \times 100$$
  
Where  $= q = \frac{q_0 + q_1}{2}$ 

Where 
$$= q = \frac{10}{2}$$

Here the average of the quantities of two years is used as weights

#### Example 3:

Construct price index number from the following data by applying

- 1. Laspeyere's Method
- 2. Paasche's Method
- 3. Fisher's ideal Method

Commenditor	20	00	2001		
Commodity	Price	Qty	Price	Qty	
A	2	8	4	5	
В	5	12	6	10	
С	4	15	5	12	
D	2	18	4	20	

#### Solution:

Commodity	$\mathbf{p}_0$	q <sub>0</sub>	p <sub>1</sub>	q	P <sub>0</sub> q <sub>0</sub>	Poq1	$\mathbf{p}_1\mathbf{q}_0$	$\mathbf{p}_1 \mathbf{q}_1$
A	2	8	4	5	16	10	32	20
В	5	12	6	10	60	50	72	60
С	4	15	5	12	60	48	75	60
D	2	18	4	20	36	40	72	80
	5 E-				172	148	251	220

CLASS: I B.B.A. COUR DURSE CODE: 18BAU202	SE NAME: BUSINES UNIT: V	5S MATHEMATICS AND STATISTICS BATCH-2018-2021
CONCL CODE. 100/10202		DATCH 2010 2021
Laspeyre's price index =	$= \mathbf{P}_{01}^{L} = \frac{\Sigma \mathbf{p}_1 \mathbf{q}_0}{\Sigma \mathbf{p}_0 \mathbf{q}_0}$	× 100
	$=\frac{251}{172}$	100 = 145.93
Paasche price index numb	$er = P_{01}^{p} = \frac{\Sigma p}{\Sigma p}$	$\frac{q_1}{q_1} \times 100$
	$=\frac{2}{1}$	$\frac{20}{48} \times 100$
	= 14	18.7
Fisher's ideal index nu	mber = $\sqrt{L \times L}$	P
	$=\sqrt{(145)}$	.9)×(148.7)
	$=\sqrt{2169}$	95.33
	= 147.3	
Or		
Fisher's ideal index nu	umber = $\sqrt{\frac{\Sigma p}{\Sigma p}}$	$\frac{p_1 q_0}{p_0 q_0} \times \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1} \times 100$

leal index number = 
$$\sqrt{\frac{27140}{\Sigma p_0 q_0}} \times \frac{27141}{\Sigma p_0 q_1} \times 100$$
  
=  $\sqrt{\frac{251}{172}} \times \frac{220}{148} \times 100$   
=  $\sqrt{(1.459)} \times (1.487) \times 100$   
=  $\sqrt{2.170} \times 100$   
=  $1.473 \times 100 = 147.3$ 

## Interpretation:

The results can be interpreted as follows:

If 100 rupees were used in the base year to buy the given commodities, we have to use Rs 145.90 in the current year to buy the same amount of the commodities as per the Laspeyre's formula. Other values give similar meaning.

CLASS: I B.B.A. JRSE CODE: 18BAU202	COURSE NAME: BUSINES UNIT: V	S MATHEMATICS AND STATISTICS BATCH-2018-2021
When the spec weighted index num	ber is calculated by the	for each commodity, the formula.
Weighted Average of	of Price Relative index =	$=\frac{\Sigma pw}{\Sigma w}$
Where $w = the weighted weigh$	tht of the commodity e relative index	
$=\frac{\mathbf{p}_1}{\mathbf{p}_0}\times 10$	00	
then the formula is		the weight i.e. $W=P_0q_0$
Weighted Average of	of Price Relative index =	
This is nothing but I When the weights an		$= \frac{\Sigma p_1 q_0}{\Sigma p_0 q_0} \times 100$ formula is
Weighted Average of	of Price Relative index =	$\Sigma p_0 q_1$
		$=\frac{\Sigma p_1 q_1}{\Sigma p_0 q_1} \times 100$
This is nothing but	Paasche' s Formula.	
	Kumar, Asst Prof, Department of M	Aathematics, KAHE Page 13

### KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B.B.A. COURSE CODE: 18BAU202

#### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V BATCH-2018-2021

### Example 6:

Compute the weighted index number for the following data.

Commodity	Prie	ce	Weight
	Current year	Base year	
A	5	4	60
В	3	2	50
С	2	1	30

## Solution:

Commodity	P <sub>1</sub>	P <sub>0</sub>	W	$P = \frac{p_1}{p_0} \times 100$	PW
Α	5	4	60	125	7500
В	3	2	50	150	7500
С	2	1	30	200	6000
			140		21000

Weighted Average of Price Relative index =  $\frac{\Sigma pw}{\Sigma w}$ 

$$=\frac{21000}{140}$$
  
= 150

## 10.7 Tests of Consistency of index numbers:

Several formulae have been studied for the construction of index number. The question arises as to which formula is appropriate to a given problems. A number of tests been developed and the important among these are

- 1. Unit test
- 2. Time Reversal test
- 3. Factor Reversal test

KARPA	GAM ACADEMY OF HIGHER EDU	CATION
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	ATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021

## 1. Unit test:

The unit test requires that the formula for constructing an index should be independent of the units in which prices and quantities are quoted. Except for the simple aggregate index (unweighted), all other formulae discussed in this chapter satisfy this test.

## 2. Time Reversal test:

Time Reversal test is a test to determine whether a given method will work both ways in time, forward and backward. In the words of Fisher, "the formula for calculating the index number should be such that it gives the same ratio between one point of comparison and the other, no matter which of the two is taken as base". Symbolically, the following relation should be satisfied.

 $P_{01} \times P_{10} = 1$ 

Where  $P_{01}$  is the index for time '1' as time '0' as base and  $P_{10}$  is the index for time '0' as time '1' as base. If the product is not unity, there is said to be a time bias is the method. Fisher's ideal index satisfies the time reversal test.

$$P_{01} = \sqrt{\frac{\Sigma p_1 q_0}{\Sigma p_0 q_0}} \times \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1}$$

$$P_{10} = \sqrt{\frac{\Sigma p_0 q_1}{\Sigma p_1 q_1}} \times \frac{\Sigma p_0 q_0}{\Sigma p_1 q_0}$$

$$p_{10} = \sqrt{\frac{\Sigma p_1 q_1}{\Sigma p_1 q_1}} \times \frac{\Sigma p_0 q_0}{\Sigma p_1 q_0}$$

Then 
$$P_{01} \times P_{10} = \sqrt{\frac{2P_1 q_0}{\Sigma p_0 q_0}} \times \frac{2P_1 q_1}{\Sigma p_0 q_1} \times \frac{2P_0 q_1}{\Sigma p_1 q_1} \times \frac{2P_0 q_0}{\Sigma p_1 q_0}$$

$$=\sqrt{1} = 1$$

Therefore Fisher ideal index satisfies the time reversal test.

## 3. Factor Reversal test:

Another test suggested by Fisher is known s factor reversal test. It holds that the product of a price index and the quantity index should be equal to the corresponding value index. In the words of Fisher, "Just as each formula should permit the interchange of the two times without giving inconsistent results, so it ought to permit interchanging the prices and quantities without giving inconsistent result, ie, the two results multiplied together should give the true value ratio.

In other word, if  $P_{01}$  represent the changes in price in the current year and  $Q_{01}$  represent the changes in quantity in the current year, then

$$\mathbf{P}_{01} \times \mathbf{Q}_{01} = -\frac{\Sigma \mathbf{p}_1 \mathbf{q}_1}{\Sigma \mathbf{p}_0 \mathbf{q}_0}$$

Thus based on this test, if the product is not equal to the value ratio, there is an error in one or both of the index number. The Factor reversal test is satisfied by the Fisher's ideal index.

CLASS: I B.B.A.	AM ACADEMY OF HIGHER EDUCA COURSE NAME: BUSINESS MAT	
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021
ie. P <sub>01</sub> =	$\sqrt{\frac{\Sigma p_1 q_0}{\Sigma p_0 q_0}} \times \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1}$	
Q <sub>01</sub> =	$\sqrt{\frac{\Sigma q_1 p_0}{\Sigma q_0 p_0}} \times \frac{\Sigma q_1 p_1}{\Sigma q_0 p_1}$	
Then $P_{01} \times Q_{01} =$	$\sqrt{\frac{\Sigma \mathbf{p}_1 \mathbf{q}_0}{\Sigma \mathbf{p}_0 \mathbf{q}_0}} \times \frac{\Sigma \mathbf{p}_1 \mathbf{q}_1}{\Sigma \mathbf{p}_0 \mathbf{q}_1} \times \frac{\Sigma \mathbf{q}_1 \mathbf{p}_1}{\Sigma \mathbf{q}_0 \mathbf{p}_1}$	$\frac{p_0}{p_0} \times \frac{\Sigma \mathbf{q}_1 \mathbf{p}_1}{\Sigma \mathbf{q}_0 \mathbf{p}_1}$
=	$\sqrt{\left(\frac{\Sigma p_1 q_1}{\Sigma p_0 q_0}\right)^2}$	
=	$\frac{\Sigma \mathbf{p}_1 \mathbf{q}_1}{\Sigma \mathbf{p}_0 \mathbf{q}_0}$	

Since  $P_{01} \times Q_{01} = \frac{\Sigma p_1 q_1}{\Sigma p_0 q_0}$ , the factor reversal test is satisfied by

the Fisher's ideal index.

## Example 8:

Construct Fisher's ideal index for the Following data. Test whether it satisfies time reversal test and factor reversal test.

	Base	year	Current yea		
Commodity	Quantity	Price	Quantity	Price	
A	12	10	15	12	
В	15	7	20	5	
С	5	5	8	9	

## Solution:

Commodity	$\mathbf{q}_0$	p <sub>0</sub>	$\mathbf{q}_1$	$\mathbf{p}_1$	P <sub>o</sub> q <sub>o</sub>	$p_0q_1$	$p_1q_0$	$p_1q_1$
A	12	10	15	12	120	150	144	180
B	15	7	20	5	105	140	75	100
С	5	5	8	9	25	40	45	72
					250	330	264	352

KARPAGA CLASS: I B.B.A.	M ACADEMY OF HIGHER	REDUCATION ESS MATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021
Fisher ideal index m	umber P <sub>01</sub> <sup>F</sup> = $\sqrt{\frac{\Sigma p}{\Sigma p}}$ = $\sqrt{\frac{264}{250}} \times$	$\frac{p_{1}q_{0}}{p_{0}q_{0}} \times \frac{\Sigma p_{1}q_{1}}{\Sigma p_{0}q_{1}} \times 100$ $\frac{352}{330} \times 100$
	$=\sqrt{(1.056)}$	)×(1.067) ×100
	$=\sqrt{1.127}$	×100
	= 1.062 ×	100 = 106.2
Time Reversal test	t:	
Time Revers	sal test is satisfied v	when $P_{01} \times P_{10} = 1$
$\mathbf{P}_{01} = \sqrt{\frac{\Sigma \mathbf{p}_1}{\Sigma \mathbf{p}_0}}$	$\frac{\mathbf{q}_0}{\mathbf{q}_0} \times \frac{\mathbf{\Sigma}\mathbf{p}_1\mathbf{q}_1}{\mathbf{\Sigma}\mathbf{p}_0\mathbf{q}_1}$	
$=\sqrt{\frac{26}{25}}$	$\frac{54}{50} \times \frac{352}{330}$	
	$\frac{\Sigma_0 q_1}{p_1 q_1} \times \frac{\Sigma p_0 q_0}{\Sigma p_1 q_0}$	
1. 2	$\frac{330}{352} \times \frac{250}{264}$	
	$\sqrt{\frac{264}{250}} \times \frac{352}{330} \times \frac{33}{35}$ $\sqrt{1}$	$\frac{0}{2} \times \frac{250}{264}$
=		
Hence Fisher idea	al index satisfy the	time reversal test.

LASS: I B.B.A.		<b>S MATHEMATICS AND STATISTICS</b>
RSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021
Factor Reversal	test:	
Factor Reversal	test is satisfied when P <sub>01</sub>	$\times \mathbf{Q}_{01} = \frac{\Sigma \mathbf{p}_1 \mathbf{q}_1}{\Sigma \mathbf{p}_0 \mathbf{q}_0}$
Now $P_{01} = $	$\frac{\Sigma p_1 q_0}{\Sigma p_0 q_0} \times \frac{\Sigma p_1 q_1}{\Sigma p_0 q_1}$	
=	$\frac{264}{250} \times \frac{352}{330}$	
F		
$Q_{01} = \sqrt{\frac{2}{2}}$	$\frac{\Sigma q_1 p_0}{\Sigma q_0 p_0} \times \frac{\Sigma q_1 p_1}{\Sigma q_0 p_1}$	
=	$\frac{330}{250} \times \frac{352}{264}$	
	264 352 330	352
Then $P_{01} \times Q_{01} =$	$= \sqrt{\frac{264}{250}} \times \frac{352}{330} \times \frac{330}{250} \times$	264
	$=\sqrt{\left(\frac{352}{250}\right)^2}$	
-	$=\frac{352}{250}$	
_	250	
$=\frac{\Sigma p_1 q_1}{\Sigma p_0 q_0}$		
-P 0 10		

Prepared by: Dr. S. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

Page 19/48

## 10.8 Consumer Price Index

Consumer Price index is also called the cost of living index. It represent the average change over time in the prices paid by the ultimate consumer of a specified basket of goods and services. A change in the price level affects the costs of living of different classes of people differently. The general index number fails to reveal this. So there is the need to construct consumer price index. People consume different types of commodities. People's consumption habit is also different from man to man, place to place and class to class i.e richer class, middle class and poor class.

The scope of consumer price is necessary, to specify the population group covered. For example, working class, poor class, middle class, richer class, etc and the geographical areas must be covered as urban, rural, town, city etc.

## Use of Consumer Price index

The consumer price indices are of great significance and is given below.

- 1. This is very useful in wage negotiations, wage contracts and dearness allowance adjustment in many countries.
- At government level, the index numbers are used for wage policy, price policy, rent control, taxation and general economic policies.
- 3. Change in the purchasing power of money and real income can be measured.
- 4. Index numbers are also used for analysing market price for particular kinds of goods and services.

## Method of Constructing Consumer price index:

There are two methods of constructing consumer price index. They are

- 1. Aggregate Expenditure method (or) Aggregate method.
- 2. Family Budget method (or) Method of Weighted Relative method.

## KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: VBATCH-2018-2021

1. Aggregate Expenditure method:

This method is based upon the Laspeyre's method. It is widely used. The quantities of commodities consumed by a particular group in the base year are the weight.

The formula is Consumer Price Index number =  $\frac{\Sigma p_1 q_0}{\Sigma p_0 q_0} \times 100$ 

2. Family Budget method or Method of Weighted Relatives:

This method is estimated an aggregate expenditure of an average family on various items and it is weighted. The formula is

Consumer Price index number =  $\frac{\Sigma pw}{\Sigma w}$ 

Where  $P = \frac{p_1}{p_0} \times 100$  for each item.  $w = value weight (i.e) p_0 q_0$ 

"Weighted average price relative method" which we have studied before and "Family Budget method" are the same for finding out consumer price index.

#### Example 9:

Construct the consumer price index number for 1996 on the basis of 1993 from the following data using Aggregate expenditure method.

	Pr	ice in
Quantity consumed	1993	1996
100	8	12
25	6	7
10	5	8
20	15	18
		Quantity consumed 1993

Solution:

Commodity	q <sub>0</sub>	p <sub>0</sub>	<b>p</b> <sub>1</sub>	$p_0 q_0$	$p_1q_0$
A	100	8	12	800	1200
В	25	6	7	150	175
С	10	5	8	50	80
D	20	15	18	300	360
			Total	1300	1815

Consumer price index by Aggregate expenditure method

#### KARPAGAM ACADEMY OF HIGHER EDUCATION

CLASS: I B.B.A. COURSE CODE: 18BAU202

## COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V BATCH-2018-2021

$$= \frac{\Sigma p_1 q_0}{\Sigma p_0 q_0} \times 100$$
$$= \frac{1815}{1300} \times 100 = 139.6$$

#### Example 10:

Calculate consumer price index by using Family Budget method for year 1993 with 1990 as base year from the following data.

		Price in		
Items	Weights	1990	1993	
Food	35	(Rs.) 150	(Rs.) 140	
Rent	20	75	90	
Clothing	10	25	30	
Fuel and lighting	15	50	60	
Miscellaneous	20	60	80	

#### Solution:

Items	W	Po	P <sub>1</sub>	P =	PW
				$\frac{p_1}{p_0} \times 100$	
Food	35	150	140	93.33	3266.55
Rent	20	75	90	120.00	2400.00
Clothing	10	25	30	120.00	1200.00
Fuel and lighting	15	50	60	120.00	1800.00
Miscellaneous	20	60	80	133.33	2666.60
	100				11333.15

Consumer price index by Family Budget method =  $\frac{\Sigma pw}{\Sigma w}$ 

$$= \frac{11333.15}{100}$$
$$= 113.33$$

KARPAGAM ACADEMY OF HIGHER EDUCATION					
CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS					
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021			

## Introduction:


Arrangement of statistical data in chronological order ie., in accordance with occurrence of time, is known as "Time Series". Such series have a unique important place in the field of Economic and Business statistics. An economist is interested in estimating the likely population in the coming year so that proper planning can be carried out with regard to food supply, job for the people etc. Similarly, a business man is interested in finding out his likely sales in the near future, so that the businessman could adjust his production accordingly and avoid the possibility of inadequate production to meet the demand. In this connection one usually deal with statistical data, which are collected, observed or recorded at successive intervals of time. Such data are generally referred to as ' time series'.

## Definition:

According to Mooris Hamburg "A time series is a set of statistical observations arranged in chronological order"

Ya-Lun- chou defining the time series as "A time series may be defined as a collection of readings belonging to different time periods, of some economic variable or composite of variables. A time series is a set of observations of a variable usually at equal intervals of time. Here time may be yearly, monthly, weekly, daily or even hourly usually at equal intervals of time.

Hourly temperature reading, daily sales, monthly production are examples of time series. Number of factors affect the observations of time series continuously, some with equal intervals of time and others are erratic studying, interpreting analyzing the factors is called Analysis of Time Series.



Regular

In time series analysis, it is assumed that there is a multiplicative relationship between these four components. Symbolically,

 $Y = T \times S \times C \times I$ 

Where Y denotes the result of the four elements; T = Trend; S = Seasonal component; C = Cyclical components; I = Irregular component

In the multiplicative model it is assumed that the four components are due to different causes but they are not necessarily independent and they can affect one another.

Another approach is to treat each observation of a time series as the sum of these four components. Symbolically

 $\mathbf{Y} = \mathbf{T} + \mathbf{S} + \mathbf{C} + \mathbf{I}$ 

The additive model assumes that all the components of the time series are independent of one another.

1) Secular Trend or Long - Term movement or simply Trend

- 2) Seasonal Variation
- 3) Cyclical Variations
- 4) Irregular or erratic or random movements(fluctuations)

	CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS						
C	COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021				

## 8.2.2 Methods of Measuring Trend:

Trend is measured by the following mathematical methods.

- 1. Graphical method
- 2. Method of Semi-averages
- 3. Method of moving averages
- 4. Method of Least Squares

## Method of Moving Averages:

This method is very simple. It is based on Arithmetic mean. Theses means are calculated from overlapping groups of successive

time series data. Each moving average is based on values covering a fixed time interval, called "period of moving average" and is shown against the center of the interval.

The method of 'odd period of moving average is as follows.

(3 or 5). The moving averages for three years is  $\frac{a+b+c}{3}$ ,

$$\frac{b+c+d}{3}, \frac{c+d+e}{3}$$
 etc

The formula for five yearly moving average is  $\frac{a+b+c+d+c}{5}$ 

$$\frac{b+c+d+e+f}{5}, \frac{c+d+e+f+g}{5}$$
 etc.

	KARPAGAM ACADEMY OF HIGHER EDUCATION					
	CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS					
•	COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021			

Steps for calculating odd number of years.

- Find the value of three years total, place the value against the second year.
- Leave the first value and add the next three years value (ie 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> years value) and put it against 3<sup>rd</sup> year.
- 3. Continue this process until the last year's value taken.
- 4. Each total is divided by three and placed in the next column.

These are the trend values by the method of moving averages

## Example 4 :

Calculate the three yearly average of the following data.

Year	1975	1976	1977	1978	1979	1980
Production in (tones)	50	36	43	45	39	38

Year	1981	1982	1983	1984
Production in (tones)	33	42	41	34

KARPAGAM ACADEMY OF HIGHER EDUCATION									
CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS									
COURSE CODE: 18BAU202	UNIT: V BATCH-2018-2021								
Solution:	Solution								
	Production	3 vears	3 years moving						

Year	(in tones)	moving total	average as Trend values
1975	50	18 <u>-</u> 1	<b>1</b> 17
1976	36	129	43.0
1977	43	124	41.3
1978	45	127	42.3
1979	39	122	40.7
1980	38	110	36.7
1981	33	113	37.7
1982	42	116	38.7
1983	41	117	39.0
1984	34	-	-

## **Even Period of Moving Averages:**

When the moving period is even, the middle period of each set of values lies between the two time points. So we must center the moving averages.

KARPAG	KARPAGAM ACADEMY OF HIGHER EDUCATION				
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	ATHEMATICS AND STATISTICS			
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021			

The steps are

- Find the total for first 4 years and place it against the middle of the 2<sup>nd</sup> and 3<sup>rd</sup> year in the third column.
- Leave the first year value, and find the total of next four-year and place it between the 3<sup>rd</sup> and 4<sup>th</sup> year.
- 3. Continue this process until the last value is taken.
- Next, compute the total of the first two four year totals and place it against the 3<sup>rd</sup> year in the fourth column.
- 5. Leave the first four years total and find the total of the next two four years' totals and place it against the fourth year.
- 6. This process is continued till the last two four years' total is taken into account.
- 7. Divide this total by 8 (Since it is the total of 8 years) and put it in the fifth column.

These are the trend values.

## Example 5 :

The production of Tea in India is given as follows. Calculate the Four-yearly moving averages

Year	1993	1994	1995	1996	1997	1998
Production (tones)	464	515	518	467	502	540

Year	1999	2000	2001	2002
Production (tones)	557	571	586	612

KARPA	GAM ACADEMY OF HIGHER EDU	CATION
CLASS: I B.B.A.	COURSE NAME: BUSINESS M	ATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021

## Solution:

Year	Production (in tones)	4 years Moving total	Total of Two four years	Trend Values
1993	464		-	-
		-		
1994	515			
		1964		
1995	518	5	3966	495.8
		2002		
1996	467		4029	503.6
		2027		
1997	502		4093	511.6
		2066		
1998	540		4236	529.5
		2170		
1999	557	6	4424	553.0
		2254		
2000	571		4580	572.5
		2326		
2001	586			
		-0		
2002	612			

KARPAG	KARPAGAM ACADEMY OF HIGHER EDUCATION			
CLASS: I B.B.A. COURSE NAME: BUSINESS MATHEMATICS AND STATISTIC				
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021		

## 8.3 Method of Least Square:

This method is widely used. It plays an important role in finding the trend values of economic and business time series. It helps for forecasting and predicting the future values. The trend line by this method is called the line of best fit.

The equation of the trend line is y = a + bx, where the constants *a* and *b* are to be estimated so as to minimize the sum of the squares of the difference between the given values of *y* and the estimate values of y *by* using the equation. The constants can be obtained by solving two normal equations.

$$\Sigma y - na + b\Sigma x \qquad (1)$$
  

$$\Sigma xy = a\Sigma x + b\Sigma x^2 \qquad (2)$$

Here x represent time point and y are observed values. 'n' is the number of pair-values.

## When odd number of years are given

Step 1: Writing given years in column 1 and the corresponding sales or production etc in column 2.

	AM ACADEMY OF HI		
CLASS: I B.B.A.		USINES	S MATHEMATICS AND STATISTICS
OURSE CODE: 18BAU202	UNIT: V		BATCH-2018-2021
Step 2. Write in col	umn 3 start with (	) 1 2	against column 1 and
denote it as		(a + a #	ugunot corunni i unu
Step 3: Take the mid		A	
Step 4: Find the dev			rite in column 4
Step 5: Find u <sup>2</sup> valu			
Step 6: Column 6 gi			
Now the normal equ	50 ST		
$\Sigma y = na + b\Sigma$	Cu -	(1)	where u = X-A
$\Sigma uy = a\Sigma u + l$	$b\Sigma u^2$	(2)	
Since $\Sigma u = 0$ , F	From equation (1)		
a = -	$\Sigma y$		
<i>a</i> – -	n		
From equation (2)			
$\Sigma uy = b\Sigma$	$\Sigma u^2$		
$\therefore b = \frac{\Sigma}{\Sigma}$	Luy		
$\therefore b = \frac{1}{\Sigma}$	$u^2$		
The fitted straight	t line is		
	+bu = a + b (X)	- A)	

KARPAG	AM ACADEMY OF HIGHER EDUC	ATION
CLASS: I B.B.A.	COURSE NAME: BUSINESS MA	THEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021
Example 6:		

### Example 6:

For the following data, find the trend values by using the method of Least squares

Year	1990	1991	1992	1993	1994
Production (in tones)	50	55	45	52	54


Estimate the production for the year 1996

#### Solution:

Year (x)	Production (y)	X= x -1990	u = X-A $= X-2$	u <sup>2</sup>	uy	Trend values
1990	50	0	-2	4	-100	50.2
1991	55	1	-1	1	-55	50.7
1992	45	2 A	0	0	0	51.2
1993	52	3	1	1	52	51.7
1994	54	4	2	4	108	52.2
Total	256			10	5	

Where A is an assumed value

The equation of straight line is Y = a + bX = a + bu, where u = X - 2the normal equations are  $\Sigma y = na + b\Sigma u.....(1)$   $\Sigma uy = a\Sigma u + b\Sigma u^{2} ...(2)$ since  $\Sigma u = 0$  from(1)  $\Sigma y = na$   $a = \frac{\Sigma y}{n} = \frac{256}{5} = 51.2$ From equation (2)  $\Sigma uy = b\Sigma u^{2}$  5 = 10b  $b = \frac{5}{10} = 0.5$ 



Then  $u = \frac{X - A}{1/2} = 2$  (X-A). The other steps are as given in the odd number of years.

## KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: VBATCH-2018-2021

## Example 7:

Fit a straight line trend by the method of least squares for the following data.

Year	1983	1984	1985	1986	1987	1988
Sales		· ^	с		8	5
(Rs. in lakhs)	3	8	7	9	11	14

Also estimate the sales for the year 1991

Solution:


Year (x)	Sales (y)	X = x-1983	u =2X-5	u2	иу	Trend values
1983	3	0	-5	25	-15	3.97
1984	8	1	-3	9	-24	5.85
1985	7	2	-1	1	-7	7.73
1986	9	3	1	1	9	9.61
1987	11	4	3	9	33	11.49
1988	14	5	5	25	70	13.37
Total	52	\$ *	0	70	66	3

$$u = \frac{X - A}{1/2}$$
  
= 2 (X - 2.5) = 2X - 5  
The straight line equation is  
 $y = a + bX = a + bu$   
The normal equations are  
 $\Sigma y = na \dots(1)$   
 $\Sigma uy = b\Sigma u^2 \dots(2)$ 

From (1) 52 = 6a

KARPA	GAM ACADEMY OF	HIGHER EDUCATION	
CLASS: I B.B.A.			TICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT:	V	BATCH-2018-2021
From (2) $66 = 7$ b = -1 b = -1 The fitted straight y = a + 1 y = 8.6 y = 8.6	8.67 0 b <u>66</u> 70 0.94 at line equation is <i>bu</i> 7+0.94(2X-5) 7 + 1.88X - 4.7 7 + 1.88X are 3.97 7.73		
A - 4, y -		<i>A</i> – 3, <i>y</i> – 13.37	

UNIT: V	BATCH-2018-2021
year 1991 is; put 2	X = x - 1983
	= 1991 - 1983 = 8
	$y = 3.97 + 1.88 \times 8$
	= 19.01 lakhs
show clearly the tre	end line.
	-



## **Parabolic Trend Model**

The curvilinear relationship for estimating the value of a dependent variable y from an independent variable x might take the form

 $y^{2} = a + bx + cx^{2}$ . This trend line is called the *parabola*.

For a non-linear equation  $y^{2} = a + bx + cx^{2}$ , the values of constants *a*, *b*, and *c* can be determined by solving three normal equations.

 $\Sigma y = na + b\Sigma x + c\Sigma x2$   $\Sigma xy = a\Sigma x + b\Sigma x2 + c\Sigma x3$  $\Sigma x2y = a\Sigma x2 + b\Sigma x3 + c\Sigma x4$ 

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: VBATCH-2018-2021

When the data can be coded so that  $\Sigma x = 0$  and  $\Sigma x 3 = 0$ , two term in the above expressions drop out and we have

 $\Sigma y = na + c\Sigma x2$   $\Sigma xy = b\Sigma x2$  $\Sigma x2y = a\Sigma x2 + c\Sigma x4$ 

To find the exact estimated value of the variable *y*, the values of constants *a*, *b*, and *c* need to be calculated. The values of these constants can becalculated by using the following shortest method:

$$a = \frac{\sum y - c \sum x^2}{n}; b = \frac{\sum xy}{\sum x^2} andc = \frac{n \sum x^2 y - \sum x^2 \sum y}{n \sum x^4 - (\sum x^2)^2}$$

**Example :**The prices of a commodity during 1999-2004 are given below. Fit a parabola to these data. Estimate

the price of the commodity for the year 2005.

Year	Price	Year	Price	
1999	100	2002	140	
2000	107	2003	181	
2001	128	2004	192	

**Solution:** To fit a parabola  $y^{2} = a + bx + cx^{2}$ , the calculations to determine the values of constants *a*, *b*, and *c* are shown in table

	CLASS: I B.B.A. URSE CODE: 18BAU202		COURSE NAME: BUSINESS UNIT: V			MATHEMATICS AND STATIS BATCH-2018-2021		
Calculatio	ons for Para	ıbola Tr	end I	line				
Year	Time	Price	x <sup>2</sup>	x <sup>3</sup>	x <sup>4</sup>	xy	x <sup>2</sup> y	Trend
	Scale	(y)						Values
	(x)							
						(y^)		
1999	-2	100	4	-8	16	-200	400	97.72
2000	-1	107	1	-1	1	-107	107	110.34
2001	0	128	0	0	0	0	0	126.68
2002	1	140	1	1	1	140	140	146.50
2003	2	181	4	8	16	362	724	169.88
2004	3	192	9	27	81	576	1728	196.82
	3	848	19	27	115	771	3099	847.94

(iv) 694 = 35b + 35c

Eliminating a from eqns. (ii) and (iii), we get

(v) 5352 = 280b + 168c

Solving eqns. (iv) and (v) for b and c we get b = 18.04 and c = 1.78.

Substituting values of *b* and *c* in eqn. (i), we get a = 126.68.

Hence, the required non-linear trend line becomes

 $y = 126.68 + 18.04x + 1.78x^2$ 

<b>KARPAGAM ACADEMY</b>	<b>OF HIGHER EDUCATION</b>

CLASS: I B.B.A. COURSE CODE: 18BAU202

## COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V BATCH-2018-2021

## **Conversion of Trend Equation**

Trend equation depends on:

- 1. The origin of time reference
- 2. The units of time viz., weekly, monthly, yearly etc.
- 3. The time series value relate to annual figures or monthly averages.

Trend values can be computed by the straight line method. For simplicity and convenience, the following methods are also used. They are

- 1. Shifting of origin
- 2. Conversion of annual trend equation to monthly trend equation when the y values are in annual totals and are given as monthly averages

## Conversion of annual trend values to monthly trend equation

We calculate monthly tend equation from annual trend equation. When Y unit are annual total and X units are in year then the annual trend equation can be converted into monthly trend equation i.e.; we divide 'a' by 12 and 'b' by 144.  $Y = \frac{a}{12} + \frac{b}{144}X$ . If the time unit of X in the trend equation represents only 6 months, then 'b' is divided by 72 i.e.,

$$Y = \frac{a}{12} + \frac{b}{72}X$$

The annual trend equation trend equation is Y = a + b X

The monthly trend equation is  $Y = \frac{a}{12} + \frac{b}{144}X$ 

## **Example:**

The trend of annual production of a company described by the following equation

 $Y_c = 18 + 0.6X$ 

Origin 1988; X-unit = 1 Year; Y-unit = annual production. Convert the equation to a monthly trend equation.

## Solution:

Monthly trend equation:

$$Y_c = \frac{18}{12} + \frac{0.6}{144} X$$
  
= 1.50 + 0.0041 X

KARPAGAM	ACADEMY	<b>OF HIGHER</b>	EDUCATION

CLASS: I B.B.A. COURSE CODE: 18BAU202

### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V BATCH-2018-2021

### **Exponential Trend Model**

When the given values of dependent variable y from approximately a geometric progression while the corresponding independent variable x values form an arithmetic progression, the relationship between variables x and y is given by an exponential function, and the best fitting curve is said to describe the *exponential trend*. Data from the fields of biology, banking, and

economics frequently exhibit such a trend. For example, growth of bacteria, money accumulating at compound interest, sales or earnings over a short period, and so on, follow exponential growth.

The characteristics property of this law is that the rate of growth, that is, the rate of change of y with respect to x is proportional to the values of the function. The following function has this property.

y = abcx, a > 0

The letter b is a fixed constant, usually either 10 or e, where a is a constant to be determined from the data.

To assume that the law of growth will continue is usually unwarranted, so only short range predictions can be made with any considerable degree or reliability.

If we take logarithms (with base 10) of both sides of the above equation, we obtain Log  $y = \log a + (c \log b) x (7.2)$ 

For b = 10, log b = 1, but for b = e, log b = 0.4343 (approx.). In either case, this equation is of the form y' = c + dx

Where  $y' = \log y$ ,  $c = \log a$ , and  $d = c \log b$ .

Equation (7.2) represents a straight line. A method of fitting an exponential trend line to a set of observed values of y is to fit a straight trend line to the logarithms of the y-values.

# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 18BAU202UNIT: VBATCH-2018-2021

In order to find out the values of constants *a* and *b* in the exponential function, the two normal equations to be solved are  $\sum \log y = n \log a + \log b \Sigma x$ 

 $\sum x \log y = \log a \sum x + \log b \sum x^2$ 

When the data is coded so that  $\Sigma x = 0$ , the two normal equations become  $\Sigma \log y = n \log a$  or  $\log a = \frac{1}{n} \Sigma \log y$ and  $\Sigma x \log y = \log b \Sigma x^2$  or  $\log b = \frac{\Sigma x \log y}{\Sigma x^2}$ 

Coding is easily done with time-series data by simply designating the center of the time period as x = 0, and have equal number of plus and minus period on each side which sum to zero.

### Example :

The sales (Rs. In million) of a company for the years 1995 to 1999 are:Year : 19951996199719981999Sales :1.64.513.840.2125.0Find the exponential trend for the given data and estimate the sales for 2002.

### Solution:

computational time can be reduced by coding the data. For this consider u = x-3. The necessary computations are shown in table

CLASS: I B.B.A.		OURSE		INESS MATH	EMATICS AND	
URSE CODE: 18	BAU202		UNIT: V		BATCH-20	18-2021
Fitting the E	xponential Trer	nd Lin	e			
Year Time	$u=x-3 u^2$	Sale	s y Log	$g y u \log u$	og y	
Period :	x					
1995 1	-2	4	1.60	0.2041	-0.4082	
1996 2	-1	1	4.50	0.6532	-0.6532	
1997 3	0	0	13.80	1.1390	0	
1998 4	1	1	40.20	1.6042	1.6042	
1999 5	2	4	125.00	2.0969	4.1938	

 $\log a = \frac{1}{n} \sum \log y = \frac{1}{5} (5.6983) = 1.1397$ 

 $\log b = \frac{\sum u \log y}{\sum u^2} = \frac{4.73 \ 66}{10} = 0.4737x$ Therefore log y= log a + (x+3) log b = 1.1397 + 0.4737x For sales during 2002, x = 3, and we obtain log y= 1.1397 + 0.4737 (3) = 2.5608 y = antilog (2.5608) = 363.80

### Seasonal Variations:

Seasonal Variations are fluctuations within a year during the season. The factors that cause seasonal variation are

- i) Climate and weather condition.
- ii) Customs and traditional habits.

For example the sale of ice-creams increase in summer, the umbrella sales increase in rainy season, sales of woolen clothes increase in winter season and agricultural production depends upon the monsoon etc.,

CLASS: I B.B.A.	COURSE NAME: BUSINESS M	ATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021

Secondly in marriage season the price of gold will increase, sale of crackers and new clothes increase in festival times.

So seasonal variations are of great importance to businessmen, producers and sellers for planning the future. The main objective of the measurement of seasonal variations is to study their effect and isolate them from the trend.

#### Measurement of seasonal variation:

The following are some of the methods more popularly used for measuring the seasonal variations.

- Method of simple averages.
- 2. Ratio to trend method.
- 3. Ratio to moving average method.
- 4. Link relative method

### Method of simple averages

The steps for calculations:

- i) Arrange the data season wise
- ii) Compute the average for each season.
- iii) Calculate the grand average, which is the average of seasonal averages.
- iv) Obtain the seasonal indices by expressing each season as percentage of Grand average

The total of these indices would be 100n where 'n' is the number of seasons in the year.

#### Example

Find the seasonal variations by simple average method for the data given below.

	Quarter								
Year	I	II	III	IV					
1989	30	40	36	34					
1990	34	52	50	44					
1991	40	58	54	48					
1992	54	76	68	62					
1993	80	92	86	82					

0----

CLASS: I B.B.A.	<u>SAM ACADEMY OF HIGHER EDU</u> COURSE NAME: BUSINESS M	IATHEMATICS AND STATISTICS
COURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021

		Quarter		
Year	Ι	II	III	IV
1989	30	40	36	34
1990	34	52	50	44
1991	40	58	54	48
1992	54	76	68	62
1993	80	92	86	82
Total	238	318	294	270
Average	47.6	63.6	58.8	54
Seasonal Indices	85	113.6	105	96.4

Grand average =  $\frac{47.6 + 63.6 + 58.8 + 54}{47.6 + 63.6 + 58.8 + 54}$ 

$$= \frac{224}{4} = 56$$

Seasonal Index for

I quarter =  $\frac{First \ quarterly \ Average}{Grand \ Average} \times 100$ 

$$=\frac{47.6}{56} \times 100 = 85$$

Seasonal Index for

II quarter = 
$$\frac{Second \ quarterly \ Average}{Grand \ Average} \times 100$$

$$= \frac{63.6}{56} \times 100 = 113.6$$

Prepared by: Dr. S. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.B.A.COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS<br/>BATCH-2018-2021COURSE CODE: 18BAU202UNIT: VBATCH-2018-2021Seasonal Index forIII quarter $= \frac{Third quarterly Average}{Grand Average} \times 100$  $= \frac{58.8}{56} \times 100 = 105$ Seasonal Index forIV quarter $= \frac{Fourth quarterly Average}{Grand Average} \times 100$ 

$$=\frac{54}{56}\times 100 = 96.4$$

### **Ratio to Trend Method**

The steps for calculation:

(i)By applying the method of least squares, we can obtain the trend values.

(ii)We must divide the original data of time series for each season (month quarter) by the corresponding trend values and multiply these ratio by 10, i.e.

 $\frac{T \times S \times C \times I}{T} \times 100 = S \times C \times I \times 100$ . Thus the trend eliminated values are obtained. This percentage will include seasonal, cyclical and irregular fluctuation.

- (iii)In order to eliminate the irregular and cyclical movement, the seasonal figures are averaged with any one of the measures of central tendency, mean or median. Thu we obtain the indices for seasonal variation for different season.
- (iv)Thee indices are adjusted to a total of ,200 for monthly data or 400 for quarterly data by multiplying each index by a suitable factor

 $\left(\frac{1200}{\text{thesumofthe 12 monthlyvalues}} \text{Or} \frac{400}{\text{thesumofthe 4 quarterlyvalues}}\right)$ 

(thesumofthe 12 monthlyvalues) thesumofthe 4 quarterlyvalues/ Thus we get Seasonal Index.

Prepared by: Dr. S. Santhosh Kumar, Asst Prof, Department of Mathematics, KAHE

	KARPAGAM ACADEMY OF HIGHER EDUCATION						
	CLASS: I B.B.A.	COURSE NAME: BUSINESS MA	ATHEMATICS AND STATISTICS				
C	OURSE CODE: 18BAU202	UNIT: V	BATCH-2018-2021				

### **Ratio to Moving Average Method**

The steps for calculation:

(i)Calculate12 month moving average , which eliminates seasonal and irregular fluctuations and represents trend and cycle. i.e., T.C.

(ii)Express the original values a percentage of centres moving average values for all month, i.e.,

$$\frac{\text{Original Value}}{\text{Moving Average vlue}} \ge 100 = \frac{\text{TSCI}}{\text{TC}} \ge 100$$
$$= \text{SI} \ge 100$$

(iii)As in the above said two methods, arrange these percentages according to year and months. By averaging these percentages for each month, we can eliminate irregular factors. Mean or Median can be used for averaging.

(iv)The sum of thee indices should be 1,200 (400) for monthly or quarterly data. If it is not so, an adjustment is made to eliminate the discrepancy; i.e.,

$$\left(\frac{1200}{1200} \text{ or } \frac{400}{1200}\right)$$

(sum of monthly indices sum of quarterly indices)

This express the preliminary index as a percentage of their arithmetic mean.

CLASS: I B.B.A. COURSE CODE: 18BAU202

#### COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS UNIT: V

#### BATCH-2018-2021

#### **POSSIBLE QUESTION**

#### **2 MARK QUESTIONS**

1. Write any two characteristics of index numbers.

2..Find the trend, where  $Y_T = 140 + 20(X - 1981)$ , if x = 1979, 1980.

3. What are the types of secular trend?

4. Write the formula for laspeyre's and pasche's method.

5. What is moving average?

#### **8 MARK QUESTIONS**

1.Compute Price Index based on the simple average of price relatives by using Arithmetic Mean.

Commodity	Α	В	С	D	Е	F	G	Н
Price 1997 (in Rs.):	40	120	140	130	60	70	65	75
Price 1998 (in Rs.)	60	140	170	135	100	80	75	80

2.Draw a trend line by the method of semi averages.

Year	:1987	1988	1989	1990	1991	1992	1993	

Production : 90 110 130 150 100 150 200

**3.** Construct the Cost of Living Index number for the following data:

Item	<b>Base Year Price</b>	<b>Current Year Price</b>	Weight
Food	39	47	4
Fuel	8	12	1
Clothing	14	18	3
House rent	12	15	2
Miscellaneous	5	30	1

4. Fit a linear trend equation by the method of least squares and estimate the net profit in 2003.

Year	: 1995	1996	1997	1998	1999	2000	2001
Net profi	it : 32	36	44	37	71	72	109
(Rs.)							

5.Calculate the Price Index by Weighted G.M method for the following data.

Item	Α	В	С	D	Е
Weight	40	25	15	10	10
Price 2001 (in Rs.):	25	50	40	10	20
Price 2011 (in Rs.)	40	80	60	20	40

RSE COD	E: 18BAU202		UNIT: V	T: V BATCH-2018-2021					
6. Using Year:		oving averages 2 1983 198	determine the tren 4 1985 1986	d and short ter 1987 1988					
Production				557 571	586 612				
					e the net profit in 2003	3			
Year		996 1997	1998 1999		2001	).			
		36     44	37     71	72	109				
	evre's, Paasche	's and Fisher's	Price Index for th	e following da	ta:				
			2003		2004				
	Commodity	Price (Rs)	Quantity ('00)	Price (Rs)	Quantity ('00)				
ľ	Erasers	05	25	06	30				
	Pencils	10	05	15	04				
	Markers	03	40	02	50				
	Ball Pens	06	30	08	35	l			
	-	ex numbers by	(i) Simple G.M me	thod (ii) Weig	hted G.M method				
	owing data.								
	ltem	А		C	D E				
	Weight	40		15	10 10				
	Price in 1991(R Price in 2001(R	/		40 60	$   \begin{array}{cccc}     10 & 20 \\     20 & 40   \end{array} $				
		*							

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
A is an arrangement of statistical data in a chronological order.	forecasting	evaluation	comparison	all the above			all the above
Time series helps in	3	4	2	5			4
There are types of components of a time series.	multiplicative	secular	additive	cyclical			additive
The model assumes that the observed value is the sum of four component of time series	multiplicative	secular	additive	cyclical			multiplicative
The model assumes that the observed value is obtained by multiplying the trend by the rates of three other components	5	1	7	3			1
Seasonal variations repeat during a period of years.	depression in business	growth of population	weather and social customs	none of these			weather and social customs
The most important factor causing seasonal variations is	ratio to trend method	simple average method	ratio to moving average method	none of these			simple average method
If the trend is absent, the seasonal indices are known by	trend is not clear	the trend is linear	trend is not linear	none of these			trend is not linear
The trend can be found by the method of least squares if the	rate of growth is positive	growth rate is constant	growth is not constant	none of these			growth rate is constant
The trend is linear if	ratio to trend method	link relative	ratio to moving average	none of these			ratio to moving average
The most widely used method of measuring seasonal variations is	graphic method	method of least squares	semi average method	moving average method			method of least squares
Themay be used either to fit a straight line trend or a parabolic trend.	which year was selected as the origin?	what is the unit of time represented by X?	In what kind of units is Y being measured?	all the above			all the above
Whenever we fit any straight line trend by the least squares method, which things should be specified?	linear trend	secular trend	non-linear trend	none of the above			non-linear trend

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
The simplest example of the is the second degree parabola.	1	2	3	0			0
In second deree parabola when time origin is taken between two middle years $\sum X$ would be	straight line	non-linear curve	either a or b	none of these			either a or b
Trends may also be plotted on a semi-loarithmic chart in the form of a	1	2	3	4			2
How many types of trend are usually computed by logarithms?	exponential tends	growth curves	both a and b	none of these			both a and b
The types of trend usually computed by logarithms are	seasonal	secular	cyclical	irregular			seasonal
A variations repeat during a period of 1 year.	time series	data	correlation	index number			time series
The helps in forescating, evaluation and comparison.	non-linear	linear	clear	none of these			linear
The trend is if growth rate is constant.	seasonal	secular	cyclical	irregular			seasonal
The most important factor causing variations is weather and social customs.	ellipse	parabola	hyperbola	circle			parabola
The simplest example of the non-linear trend is the second degree	averages	percentages	economic activity	time series			time series
Index numbers are special type of	current	past	future	arbitrary			current
The base period should not be too distant from the	unit test	time reversal test	factor test and circular test	all the above			all the above
A good index number is one that satisfies	Index numbers	averages	time series	trend			Index numbers

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
help to calculate the real wages	A.M	G.M	H.M	both A.M and G.M			G.M
The best average to calculate index number is	simple A.M method	Kelly's method	Laspeyre's method	Fisher's method			Fisher's method
Current year quantity is used in	Base year quartiles	Current year quartiles	Both of them	Average of current and base year			Base year quartiles
Laspeyre's index is based on	Laspeyre's method	Paasche's method	Fisher's method	Bow ley's method			Fisher's method
Time Reversal test is satisfied by	8	90	111.11	110			90
If the price of a commodity is Rs.80 in the base year and Rs.72 in the current year, the Price index number is	current year	base year	future year	arbitrary year			base year
In Laspeyre's index number, importance is given to the quantity of	Bow ley's formula	Laspeyre's formula	Paasche's formula	Fisher's formula			Paasche's formula
The current year quantities are taken as weights in	P01 xP10 = 0	$P_{01} x P_{10} < 1$	$P_{01} x P_{10} = 1$	$P_{01} x P_{10} > 1$			$P_{01} x P_{10} = 1$
Time reversal test condition is	$P_{01}Q_{01} = a_{1}p_{1}q_{1}/a_{0}p_{0}q_{0}$	$\frac{P_{01}Q_{01}{$	$P_{01}Q_{01} > a p_1 q_1 / a p_0 q_0$	$P_{01}Q_{01} = ap_1q_1$			$P_{01}Q_{01} = a p_1 q_1 / a p_0 q_0$
Factor reversal test condition is =	an upward bias	a downward bias	either upward or downward bias	no bias			an upward bias
Paasche's index number is generally expected to have	å log P/N	åP/N	N/åP	åP/ log N			åP/N
The formula for unweighted averages of relatives' method by using A.M.,	arithmetic mean of Laspeyre's and Paasche's index	geometric mean of Laspeyre's and Paasche's index	median of Lasoeyre's and Paasche's index	all of the above			geometric mean of Laspeyre's and Paasche's index
Fisher's ideal index is	151	150	151.61	125.2			151.61

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
If $ap_0$ is 3100 and $ap_1$ is 4700 then $P_{01} =$	Laspeyre's formula	Paasche's formula	Fisher's formula	both a) and b)			Fisher's formula
$P = \sqrt{P_{01}}^{L} x P_{01}^{P}$ is	consumer	Laspeyre's	Paasche's	Fisher			consumer
Family budget method is a method to calculate price index.	median	geometric mean	arithmetic mean	mode			geometric mean
The best average in the construction of index number is	value index	price index	quantity index	quality index			quantity index
Commodities which show considerable price fluctuation could be best measured by a	factor reversal test	time reversal test	t-test	f-test			time reversal test
The circular test is an extension of the	weighted	fixed weighted	un weighted	fixed un weighted			weighted
Most frequently used index number formulae are	base year quantities	current year quantities	base year qualities	current year qualities			base year quantities
Laspeyre's index is based on	Kelly's	Walsch's	Fisher's price	Marshall- Edgeworth's			Walsch's
index number uses the geometric mean of the base year and current year quantities as weights.	Value index	Laspeayre's index	Paasche's index	Fisher ideal index			Value index
is the sum of the values of a given year divided by the sum of the values of the base year.	$(p_1 / p_{-0}) * 100$	$(p_1 / q_{-0}) * 100$	$(q_1 / p_{-1}) * 100$	$(p_1 / q_1) * 100$			$(p_1 / p_{-0}) * 100$
Formula for price relative or price index number of a commodity P is	Ideal	economic	special	commercial			Ideal
Fisher's formula is called index number formula	Laspeyre's method	Paasche's method	Fisher's method	both a) and b)			Fisher's method
Factor reversal test is satisfied by	current	base	average	calculated			current

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
The year for which index number is calculated is calledyear.	p <sub>1</sub>	p <sub>0</sub>	q <sub>0</sub>	<b>q</b> <sub>1</sub>			<b>p</b> <sub>1</sub>
Notation of price of a commodity in the current year is	Time series	Mean	Mode	Index number			Index number
are the pulse of an economy	100/Price index	Price index /100	Money wage/Price index *100	Price index *100			100/Price index
Purchasing power =	p <sub>1</sub>	<b>p</b> <sub>0</sub>	<b>q</b> <sub>0</sub>	<b>q</b> <sub>1</sub>			<b>p</b> <sub>0</sub>
Notation of price of a commodity in the base year is	p <sub>1</sub>	p <sub>0</sub>	q <sub>0</sub>	<b>q</b> <sub>1</sub>			q <sub>1</sub>
Notation of quantity of a commodity in the current year is	25%	10%	125%	35%			25%
If the price of a commodity is Rs.40 in the base year and Rs.50 in the current year, the Price has increased by	$P_{01} \times P_{12} \times P_{20} = 1$	$P_{12} \ x \ P_{20} = 1$	$P_{01} \ge P_{12} = 1$	$ \begin{array}{c} P_{01} x P_{12} & x P_{20} \\ = 0 \end{array} $			$P_{01} \ge P_{12} \ge P_{20} = 1$
By circular test	current year	base year	arbitrary year	previous year			base year
Link relative is a price or quantity relative with the condition that is the preceding year.	Consumer price	Consumer price index number	Consumer price index number	price index number			Consumer price index number
Cost of living index number is also known as	price	quantity	both	neither price nor quantity			price
In Factor reversal test $P_{01}$ gives the relative change in	price	quantity	both	neither price nor quantity			quantity
In Factor reversal test $Q_{01}$ gives the relative change in	Time reversal test	Factor reversal test	Fisher's test	both a and b			Time reversal test
satisfies the Kelly's test.	Time reversal test	Factor reversal test	Fisher's test	both a and b			both a and b

Questions	opt1	opt2	opt3	opt4	opt5	opt6	Answer
Fisher's index satisfy	price multiplied by quantity	quantity	both	price			price multiplied by quantity
In Factor reversal test $P_{01} \ge Q_{01}$ gives the relative change in	ratio	percentage	fraction	mean			percentage
Index number are expressed in	base year quantities	current year quantities	hase year dualities	current year qualities			current year quantities
Paasche index is based on	forecasting	evaluation	comparison	index numbers			Index numbers

CLASS: I BBA COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 **UNIT: III** 

BATCH-2018-2021

#### <u>UNIT – III</u>

#### **SYLLABUS**

#### **Uni-variate Analysis**

Measures of Central Tendency including arithmetic mean, geometric mean and harmonic mean: properties and applications; mode and median. Partition values - quartiles, deciles, and percentiles. Measures of Variation: absolute and relative. Range, quartile deviation and mean deviation; Variance and Standard deviation: calculation and properties.

#### Measures of Central Tendency:

In the study of a population with respect to one in which we are interested we may get a large number of observations. It is not possible to grasp any idea about the characteristic when we look at all the observations. So it is better to get one number for one group. That number must be a good representative one for all the observations to give a clear picture of that characteristic. Such representative number can be a central value for all these observations. This central value is called a measure of central tendency or an average or a measure of locations. There are five averages. Among them mean, median and mode are called simple averages and the other two averages geometric mean and harmonic mean are called special averages.

### Characteristics for a good or an ideal average :

The following properties should possess for an ideal average.

- 1. It should be rigidly defined.
- 2. It should be easy to understand and compute.
- 3. It should be based on all items in the data.
- 4. Its definition shall be in the form of a mathematical formula.
- It should be capable of further algebraic treatment.
- It should have sampling stability.
- 7. It should be capable of being used in further statistical computations or processing.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

### Arithmetic mean or mean :

Arithmetic mean or simply the mean of a variable is defined as the sum of the observations divided by the number of observations. If the variable x assumes n values  $x_1, x_2 ... x_n$  then the mean,  $\overline{x}$ , is given by

$$\overline{x} = \frac{x_1 + x_2 + x_3 + \dots + x_n}{n}$$
$$= \frac{1}{n} \sum_{i=1}^n x_i$$

This formula is for the ungrouped or raw data.

### Example 1 :

Calculate the mean for 2, 4, 6, 8, 10

### Solution:

$$\overline{x} = \frac{2+4+6+8+10}{5} = \frac{30}{5} = 6$$

### Short-Cut method :

Under this method an assumed or an arbitrary average (indicated by A) is used as the basis of calculation of deviations from individual values. The formula is

$$\overline{x} = A + \frac{\sum d}{n}$$

where, A = the assumed mean or any value in x

d = the deviation of each value from the assumed mean

### Example 2 :

A student's marks in 5 subjects are 75, 68, 80, 92, 56. Find his average mark.

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

Solution:

X	d=x-A	
75	7	
A 68	0	
80	12	
92	24	
56	-12	
Total	31	
$\Sigma J$		
$\overline{x} = A + \frac{\sum d}{n}$		
x = A +		
$x = A + \frac{n}{n}$		

### Grouped Data :

The mean for grouped data is obtained from the following formula:

$$\overline{x} = \frac{\sum fx}{N}$$

where x = the mid-point of individual class

f = the frequency of individual class

N = the sum of the frequencies or total frequencies.

### Short-cut method :

$$\overline{x} = A + \frac{\sum fd}{N} \times c$$
where  $d = \frac{x - A}{c}$   
 $A = any value in x$   
 $N = total frequency$   
 $c = width of the class interval$ 

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

### Example 3:

Given the following frequency distribution, calculate the arithmetic mean Marks : 64 63 62 61 60 59 Number of Students { : 8 18 12 9 7 6

#### Solution:

X	F	fx	d=x-A	fd	
64	8	512	2	16	
63	18	1134	1	18	
62	12	744	0	0	
61	9	549	-1	_9	
60	7	420	-2	-14	
59	6	354	-2 -3	-18	
	60	3713		- 7	1

Direct method

$$\overline{x} = \frac{\sum fx}{N} = \frac{3713}{60} = 61.88$$

Short-cut method

$$\overline{x} = A + \frac{\sum fd}{N} = 62 - \frac{7}{60} = 61.88$$

### Example 4 :

Following is the distribution of persons according to different income groups. Calculate arithmetic mean.

Income Rs(100)	0-10	10-20	20-30	30-40	40-50	50-60	60-70
Number of persons	6	8	10	12	7	4	3

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

### Solution:

mber of	Mid		<b>T</b> 1
	IVIIG	X - A	Fd
rsons (f)	Х	$d = \frac{c}{c}$	
6	5	-3	-18
8	15	-2	-16
10	25	-1	-10
12	A 35	0	0
7	45	1	7
4	55	2	8
3	65	3	9
50			-20
	8 10 12 7 4 3	6       5         8       15         10       25         12       A 35         7       45         4       55         3       65	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

## CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

Mean =  $\overline{x} = A + \frac{\sum fd}{N}$ =  $35 - \frac{20}{50} \times 10$ = 35 - 4= 31

### Merits and demerits of Arithmetic mean :

### Merits:

- 1. It is rigidly defined.
- 2. It is easy to understand and easy to calculate.
- 3. If the number of items is sufficiently large, it is more accurate and more reliable.
- It is a calculated value and is not based on its position in the series.
- 5. It is possible to calculate even if some of the details of the data are lacking.
- Of all averages, it is affected least by fluctuations of sampling.
- 7. It provides a good basis for comparison.

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

### Demerits:

- 1. It cannot be obtained by inspection nor located through a frequency graph.
- It cannot be in the study of qualitative phenomena not capable of numerical measurement i.e. Intelligence, beauty, honesty etc.,
- It can ignore any single item only at the risk of losing its accuracy.
- 4. It is affected very much by extreme values.
- 5. It cannot be calculated for open-end classes.
- 6. It may lead to fallacious conclusions, if the details of the data from which it is computed are not given.

### Median :

The median is that value of the variate which divides the group into two equal parts, one part comprising all values greater, and the other, all values less than median.

### Ungrouped or Raw data :

Arrange the given values in the increasing or decreasing order. If the number of values are odd, median is the middle value .If the number of values are even, median is the mean of middle two values.

By formula

Median = Md = 
$$\left(\frac{n+1}{2}\right)^{\text{th}}$$
 item.

### Example 11:

When odd number of values are given. Find median for the following data

25, 18, 27, 10, 8, 30, 42, 20, 53

### Solution:

Arranging the data in the increasing order 8, 10, 18, 20, 25, 27, 30, 42, 53

The middle value is the 5<sup>th</sup> item i.e., 25 is the median

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

Using formula

Md = 
$$\left(\frac{n+1}{2}\right)^{\text{th}}$$
 item.

$$=\left(\frac{9+1}{2}\right)^{\text{th}}$$
 item.

$$= \left(\frac{10}{2}\right)^{\text{th} \text{ item}}$$
$$= 5^{\text{th}} \text{ item}$$
$$= 25$$

### Example 12 :

When even number of values are given. Find median for the following data

5, 8, 12, 30, 18, 10, 2, 22

### Solution:

Arranging the data in the increasing order 2, 5, 8, 10, 12, 18, 22, 30

Here median is the mean of the middle two items (ie) mean of (10,12) ie

$$=\left(\frac{10+12}{2}\right) = 11$$

∴median = 11.

Using the formula

Median = 
$$\left(\frac{n+1}{2}\right)^{\text{th}}$$
 item.

#### CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

$$= \left(\frac{8+1}{2}\right)^{\text{th}} \text{ item.}$$

$$= \left(\frac{9}{2}\right)^{\text{th}} \text{ item } = 4.5^{\text{th}} \text{ item}$$

$$= 4^{\text{th}} \text{ item} + \left(\frac{1}{2}\right)(5^{\text{th}} \text{ item} - 4^{\text{th}} \text{ item})$$

$$= 10 + \left(\frac{1}{2}\right)[12 - 10]$$

$$= 10 + \left(\frac{1}{2}\right) \times 2$$

$$= 10 + 1$$

$$= 11$$

### Example 13:

The following table represents the marks obtained by a batch of 10 students in certain class tests in statistics and Accountancy.

Serial No	1	2	3	4	5	6	7	8	9	10
Marks (Statistics)	53	55	52	32	30	60	47	46	35	28
Marks (Accountancy)	57	45	24	31	25	84	43	80	32	72

Indicate in which subject is the level of knowledge higher ?

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

### Solution:

For such question, median is the most suitable measure of central tendency. The mark in the two subjects are first arranged in increasing order as follows:

Serial No	1	2	3	4	5	6	7	8	9	10
Marks in Statistics	28	30	32	35	46	47	52	53	55	60
Marks in Accountancy	24	25	31	32	43	45	57	72	80	84

Median =  $\left(\frac{n+1}{2}\right)^{\text{th}}$  item =  $\left(\frac{10+1}{2}\right)^{\text{th}}$  item =5.5<sup>th</sup> item

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

$$= \frac{Value \ of \ 5^{th} \ item + value \ of \ 6^{th} \ item}{2}$$
  
Md (Statistics) 
$$= \frac{46 + 47}{2} = 46.5$$
  
Md (Accountancy) 
$$= \frac{43 + 45}{2} = 44$$

There fore the level of knowledge in Statistics is higher than that in Accountancy.

### Grouped Data:

In a grouped distribution, values are associated with frequencies. Grouping can be in the form of a discrete frequency distribution or a continuous frequency distribution. Whatever may be the type of distribution, cumulative frequencies have to be calculated to know the total number of items.

### Cumulative frequency : (cf)

Cumulative frequency of each class is the sum of the frequency of the class and the frequencies of the pervious classes, ie adding the frequencies successively, so that the last cumulative frequency gives the total number of items.

### **Discrete Series:**

Step1: Find cumulative frequencies.

Step2: Find 
$$\left(\frac{N+1}{2}\right)$$

Step3: See in the cumulative frequencies the value just greater than

$$\left(\frac{N+1}{2}\right)$$

Step4: Then the corresponding value of x is median.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

### Example 14:


The following data pertaining to the number of members in a family. Find median size of the family.

Number of members x	1	2	3	4	5	6	7	8	9	10	11	12
Frequency F	1	3	5	6	10	13	9	5	3	2	2	1

### Solution:

X	f	cf
1	1	1
2	3	4
3	5	9
4	6	15
5	10	25
6	13	38
7	9	47
8	9 5	52
9	3	55
10	2	57
11	2	59
12	1	60
	60	

### Median = size



CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

$$= \text{size of}\left(\frac{60+1}{2}\right)^{\text{th}} \text{item}$$

 $= 30.5^{\text{th}}$  item

The cumulative frequencies just greater than 30.5 is 38.and the value of x corresponding to 38 is 6.Hence the median size is 6 members per family.

### **Continuous Series:**

The steps given below are followed for the calculation of median in continuous series.

Step1: Find cumulative frequencies.

Step2: Find  $\left(\frac{N}{2}\right)$ 

Step3: See in the cumulative frequency the value first greater than  $\left(\frac{N}{2}\right)$ , Then the corresponding class interval is called the Median

class. Then apply the formula

Median = 
$$l + \frac{\frac{N}{2} - m}{f} \times c$$

Where

l = Lower limit of the median class m = cumulative frequency preceding the median

c = width of the median class

f = frequency in the median class.

N=Total frequency.

### Example 15:

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

The following table gives the frequency distribution of 325 workers of a factory, according to their average monthly income in a certain year.

Income group (in Rs)	Number of workers
Below 100	1
100-150	20
150-200	42
200-250	55
250-300	62
300-350	45
350-400	30
400-450	25
450-500	15
500-550	18
550-600	10
600 and above	2
	325

## Calculate median income **Solution:**

· <b>u</b> .		
Income group	Number of	Cumulative
(Class-interval)	workers	frequency
	(Frequency)	c.f
Below 100	1	1
100-150	20	21
150-200	42	63
200-250	55	118
250-300	62	180
300-350	45	225
350-400	30	255
400-450	25	280
450-500	15	295
500-550	18	313
550-600	10	323
600 and above	2	325
	325	

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

$$\frac{N}{2} = \frac{325}{2} = 162.5$$

Here 
$$l = 250$$
, N = 325, f = 62, c = 50, m = 118  
Md =  $250 + \left(\frac{162.5 - 118}{62}\right) \times 50$   
=  $250 + 35.89$   
=  $285.89$ 

### Example 16:

Following are the daily wages of workers in a textile. Find the median.

Wages	Number of
( in Rs.)	workers
less than 100	5
less than 200	12
less than 300	20
less than 400	32
less than 500	40
less than 600	45
less than 700	52
less than 800	60
less than 900	68
less than 1000	75
_	

CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

### Solution :

We are given upper limit and less than cumulative frequencies. First find the class-intervals and the frequencies. Since the values are increasing by 100, hence the width of the class interval equal to 100.

Class		c.f	
0-100	) 5	5	
100-20	7753	12	
200-30	9399	20	
300- 4	00 12	32	
400-50	00 8	40	
500-60		45	
600-70		52	
700-80		60	
800-90		68	
900-10		75	
	75		
$\left(\frac{N}{2}\right) = \left(\frac{N}{2}\right)$	$\left(\frac{75}{2}\right) = 3^{\circ}$		
Md = i	$r + \left(\frac{\frac{N}{2}}{f}\right)$	$\binom{m}{-} \times \mathbf{c}$	
(37.5 - 32)			

$$= 400 + \left(\frac{37.5 - 32}{8}\right) \times 100 = 400 + 68.75 = 468.75$$

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

### Merits of Median :

- Median is not influenced by extreme values because it is a positional average.
- Median can be calculated in case of distribution with openend intervals.
- 3. Median can be located even if the data are incomplete.
- Median can be located even for qualitative factors such as ability, honesty etc.

### Demerits of Median :

- A slight change in the series may bring drastic change in median value.
- In case of even number of items or continuous series, median is an estimated value other than any value in the series.
- 3. It is not suitable for further mathematical treatment except its use in mean deviation.
- 4. It is not taken into account all the observations.

### Mode :

The mode refers to that value in a distribution, which occur most frequently. It is an actual value, which has the highest concentration of items in and around it.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

### Computation of the mode: Ungrouped or Raw Data:

For ungrouped data or a series of individual observations, mode is often found by mere inspection.

### Example 29:

2, 7, 10, 15, 10, 17, 8, 10, 2

 $\therefore$  Mode = M<sub>0</sub> = 10

In some cases the mode may be absent while in some cases there may be more than one mode.

### Example 30:

- 1. 12, 10, 15, 24, 30 (no mode)
- 2. 7, 10, 15, 12, 7, 14, 24, 10, 7, 20, 10
- : the modes are 7 and 10

### Grouped Data:

For Discrete distribution, see the highest frequency and corresponding value of X is mode.

### **Continuous distribution :**

See the highest frequency then the corresponding value of class interval is called the modal class. Then apply the formula.

Mode = 
$$M_0 = l + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times C$$

l = Lower limit of the model class

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

- $f_1 =$ frequency of the modal class
- $f_0$  = frequency of the class preceding the modal class
- f<sub>2</sub> = frequency of the class succeeding the modal class The above formula can also be written as

Mode = 
$$l + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \times c$$

### Example 31:

Calculate mode for the following :

f
5
14
40
91
150
87
60
38
15

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

### Solution:

The highest frequency is 150 and corresponding class interval is 200 - 250, which is the modal class.

Here l=200, f<sub>1</sub>=150, f<sub>0</sub>=91, f<sub>2</sub>=87, C=50

Mode = 
$$M_0 = 1 + \frac{I_1 - I_0}{2f_1 - f_0 - f_2} \times c$$
  
=  $200 + \frac{150 - 91}{2 \times 150 - 91 - 87} \times 50$   
=  $200 + \frac{2950}{2}$ 

$$122 = 200 + 24.18 = 224.18$$

### **Determination of Modal class :**

For a frequency distribution modal class corresponds to the maximum frequency. But in any one (or more) of the following cases

## CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

- i.If the maximum frequency is repeated
- ii. If the maximum frequency occurs in the beginning or at the end of the distribution
- iii. If there are irregularities in the distribution, the modal class is determined by the method of grouping.

### Steps for Calculation :

We prepare a grouping table with 6 columns

- 1. In column I, we write down the given frequencies.
- Column II is obtained by combining the frequencies two by two.
- 3. Leave the 1<sup>st</sup> frequency and combine the remaining frequencies two by two and write in column III
- 4. Column IV is obtained by combining the frequencies three by three.
- 5. Leave the 1st frequency and combine the remaining frequencies three by three and write in column V
- Leave the 1st and 2<sup>nd</sup> frequencies and combine the remaining frequencies three by three and write in column VI

Mark the highest frequency in each column. Then form an analysis table to find the modal class. After finding the modal class use the formula to calculate the modal value.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

## Example 32:

Calculate mode for the following frequency distribution.

Class	0-	5-	10-	15-	20-	25-	30-	35-
interval	5	10	15	20	25	30	35	40
Frequency	9	12	15	16	17	15	10	13

## **Grouping Table**

CI	f	2	3	4	5	6
0- 5	9	21				
5-10	12	-1	27	36		
10-15	15	31	21		43	
15-20	16	51	33			48
20-25	17	32	55	48		
25-30	15	32	25		42	38
30-35	10	23	25			
35-40	13	23				

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## Analysis Table

Columns	0-5	5-10	10-15	15-20	20-25	25-30	30-35	35-40
1					1			
2					1	1		
3				1	1			
4				1	1	1		
5		1	1	1				
6			1	1	1			
Total		1	2	4	5	2		

The maximum occurred corresponding to 20-25, and hence it is the modal class.

Mode = Mo = 
$$l + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times C$$

Here l = 20;  $\triangle_1 = f_1 - f_0 = 17 - 16 = 1$  $\triangle_2 = f_1 - f_2 = 17 - 15 = 2$  $\therefore M_0 = 20 + \frac{1}{1+2} \times 5$ = 20 + 1.67 = 21.67

## **MEASURES OF DISPERSION**

## Characteristics of a good measure of dispersion:

An ideal measure of dispersion is expected to possess the following properties

1.It should be rigidly defined

- 2. It should be based on all the items.
- 3. It should not be unduly affected by extreme items.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

4. It should lend itself for algebraic manipulation.5. It should be simple to understand and easy to calculate

## Absolute and Relative Measures :

There are two kinds of measures of dispersion, namely

1. Absolute measure of dispersion

2.Relative measure of dispersion.

The various absolute and relative measures of dispersion are listed below.

## Absolute measure

## **Relative measure**

- 1. Range1. Co-efficient of Range2. Quartile deviation2. Co-efficient of Quartile deviation3. Mean deviation3. Co-efficient of Mean deviation
- 4.Standard deviation 4.Co-efficient of variation

## 7.3 Range and coefficient of Range:

## 7.3.1 Range:

This is the simplest possible measure of dispersion and is defined as the difference between the largest and smallest values of the variable.

> In symbols, Range = L - S. Where L = Largest value. S = Smallest value.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

In individual observations and discrete series, L and S are easily identified. In continuous series, the following two methods are followed.

## Method 1:

L = Upper boundary of the highest class

S = Lower boundary of the lowest class.

## Method 2:

L = Mid value of the highest class.

S = Mid value of the lowest class.

## 7.3.2 Co-efficient of Range :

Co-efficient of Range =  $\frac{L-S}{L+S}$ 

## Example1:

Find the value of range and its co-efficient for the following data.

7, 9, 6, 8, 11, 10, 4

## Solution:

L=11, S = 4. Range = L - S = 11-4 = 7 Co-efficient of Range =  $\frac{L-S}{L+S}$ =  $\frac{11-4}{11+4}$ =  $\frac{7}{15}$  = 0.4667

## Example 2:

Calculate range and its co efficient from the following distribution.

Size:	60-63	63-66	66-69	69-72	72-75
Number:	5	18	42	27	8

## Solution:

L = Upper boundary of the highest class.= 75

Page 25/46

CLASS: I B.Com(CA) COURS	<b>3E NAME:BUSINESS M</b>	ATHEMATICS AND STATISTICS
COURSE CODE: 17CCU202	UNIT: III	BATCH-2017-2020

S = Lower boundary of the lowest class. = 60 Range = L - S = 75 - 60 = 15 Co-efficient of Range =  $\frac{L-S}{L+S}$ =  $\frac{75-60}{75+60}$ =  $\frac{15}{135}$  = 0.1111

7.3.3 Merits and Demerits of Range : Merits:

- 1. It is simple to understand.
- 2. It is easy to calculate.
- In certain types of problems like quality control, weather forecasts, share price analysis, et c., range is most widely used.

## Demerits:

- 1. It is very much affected by the extreme items.
- 2. It is based on only two extreme observations.
- 3. It cannot be calculated from open-end class intervals.
- 4. It is not suitable for mathematical treatment.
- 5. It is a very rarely used measure.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

# 7.6 Standard Deviation and Coefficient of variation:7.6.1 Standard Deviation :

Karl Pearson introduced the concept of standard deviation in 1893. It is the most important measure of dispersion and is widely used in many statistical formulae. Standard deviation is also called Root-Mean Square Deviation. The reason is that it is the square-root of the mean of the squared deviation from the arithmetic mean. It provides accurate result. Square of standard deviation is called Variance.

## Definition:

It is defined as the positive square-root of the arithmetic mean of the Square of the deviations of the given observation from their arithmetic mean.

The standard deviation is denoted by the Greek letter  $\sigma$  (sigma)

## 7.6.2 Calculation of Standard deviation-Individual Series :

There are two methods of calculating Standard deviation in an individual series.

- a) Deviations taken from Actual mean
- b) Deviation taken from Assumed mean

## a) Deviation taken from Actual mean:

This method is adopted when the mean is a whole number. **Steps:** 

- 1. Find out the actual mean of the series  $(\bar{x})$
- 2. Find out the deviation of each value from the mean  $(x = X \overline{X})$
- 3.Square the deviations and take the total of squared deviations  $\Sigma x^2$

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

4. Divide the total (  $\sum x^2$  ) by the number of observation

The square root of 
$$\left(\frac{\sum x^2}{n}\right)$$
 is standard deviation.

Thus 
$$\sigma = \sqrt{\left(\frac{\Sigma x^2}{n}\right)}$$
 or  $\sqrt{\frac{\Sigma(x-\overline{x})}{n}}$ 

## b) Deviations taken from assumed mean:

This method is adopted when the arithmetic mean is fractional value.

Taking deviations from fractional value would be a very difficult and tedious task. To save time and labour, We apply short –cut method; deviations are taken from an assumed mean. The formula is:

$$\sigma = \sqrt{\frac{\sum d^2}{N} - \left(\frac{\sum d}{N}\right)^2}$$

Where d-stands for the deviation from assumed mean = (X-A)

## Steps:

- 1. Assume any one of the item in the series as an average (A)
- Find out the deviations from the assumed mean; i.e., X-A denoted by d and also the total of the deviations ∑d
- Square the deviations; i.e., d<sup>2</sup> and add up the squares of deviations, i.e, ∑d<sup>2</sup>
- 4. Then substitute the values in the following formula:

CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

$$\sigma = \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2}$$

**Note:** We can also use the simplified formula for standard deviation.

$$\sigma = \frac{1}{n} \sqrt{n \sum d^2 - (\sum d)^2}$$

For the frequency distribution

$$\sigma = \frac{c}{N} \sqrt{N \sum f d^2 - (\sum f d)^2}$$

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## Example 9:

Calculate the standard deviation from the following data.

14, 22, 9, 15, 20, 17, 12, 11

## Solution:

Deviations from actual mean.

Values (X)	X - X	$(X - \overline{X})^2$
14	-1	1
22		49
9	-6	36
15	0	0
20	5	25
17	2	4
12	7 -6 0 5 2 -3 -4	4 9
11	-4	16
120	89	140
$\overline{X} = \frac{120}{8} = 15$ $\overline{5} = \sqrt{\frac{\Sigma(x - \overline{x})^2}{n}}$ $= \sqrt{\frac{140}{8}}$		
$=\sqrt{17.5} = 4.18$		

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

#### Example 10:

The table below gives the marks obtained by 10 students in statistics. Calculate standard deviation.

Student N	Nos :	1	2	3	4	5	6	7	8	9	10
Marks	:	43	48	65	57	31	60	37	48	78	59

Solution:	(Deviations	from assumed	mean)
Solution.	Deviations	nom assumed	mcan)

Nos.	Marks (x)	d=X-A (A=57)	$d^2$
1	43	-14	196
2	48	-9	81
3	65	8	64
4	57	0	0
3 4 5	31	-26	676
6	60	3	9
7	37	-20	400
8	48	-20 -9	81
9	78	21	441
10	59	2	4
n = 10		∑d=-44	$\Sigma d^2 = 1952$

$$\sigma = \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} = \sqrt{\frac{1952}{10} - \left(\frac{-44}{10}\right)^2} = \sqrt{195.2 - 19.36} = \sqrt{175.84} = 13.26$$

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## 7.6.3 Calculation of standard deviation: Discrete Series:

There are three methods for calculating standard deviation in discrete series:

(a) Actual mean methods

(b) Assumed mean method

(c) Step-deviation method.

## (a) Actual mean method:

## Steps:

- 1. Calculate the mean of the series.
- 2. Find deviations for various items from the means i.e.,

 $\mathbf{x} \cdot \mathbf{x} = \mathbf{d}$ .

- Square the deviations (= d<sup>2</sup>) and multiply by the respective frequencies(f) we get fd<sup>2</sup>
- 4. Total to product  $(\sum fd^2)$  Then apply the formula:

 $\sum fd^2$ 

CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

## (b) Assumed mean method:

Here deviation are taken not from an actual mean but from an assumed mean. Also this method is used, if the given variable values are not in equal intervals.

## Steps:

- Assume any one of the items in the series as an assumed mean and denoted by A.
- 2. Find out the deviations from assumed mean, i.e, X-A and denote it by d.
- Multiply these deviations by the respective frequencies and get the ∑fd
- 4. Square the deviations  $(d^2)$ .
- Multiply the squared deviations (d<sup>2)</sup> by the respective frequencies (f) and get ∑fd<sup>2</sup>.
- 6. Substitute the values in the following formula:

$$\sigma = \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2}$$

Where 
$$d = X - A$$
,  $N = \sum f$ .

## Example 11:

Calci	ilate Sta	andard o	leviatio	n from	the follo	owing d	ata.		
<b>X</b> :	20	22	25	31	35	40	42	45	
f:	5	12	15	20	25	14	10	6	

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## Solution:

Deviations from assumed mean

х	f	d = x - A	$d^2$	fd	fd <sup>2</sup>
		(A = 31)			
20	5	-11	121	-55	605
22	12	-9	81	-108	972
25	15	-6	36	-90	540
31	20	0	0	0	0
35	25	4	16	100	400
40	14	9	81	126	1134
42	10	11	121	110	1210
45	6	14	196	84	1176
	N=107			∑fd=167	$\Sigma fd^2$
					=6037

$$\sigma = \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2}$$
$$= \sqrt{\frac{6037}{107} - \left(\frac{167}{107}\right)^2}$$
$$= \sqrt{56.42 - 2.44}$$

$$=\sqrt{53.98} = 7.35$$

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## 7.6.4 Calculation of Standard Deviation –Continuous series:

In the continuous series the method of calculating standard deviation is almost the same as in a discrete series. But in a continuous series, mid-values of the class intervals are to be found out. The step- deviation method is widely used.

The formula is,

$$\sigma = \sqrt{\frac{\sum {fd'}^2}{N} - \left(\frac{\sum {fd'}}{N}\right)^2} \quad \times \, C$$

$$d = \frac{m-A}{C}$$
, C- Class interval.

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## Steps:

- 1. Find out the mid-value of each class.
- 2. Assume the center value as an assumed mean and denote it by A

3. Find out d = 
$$\frac{m-A}{C}$$

- 4. Multiply the deviations d by the respective frequencies and get  $\Sigma fd$
- 5. Square the deviations and get d  $^{2}$
- 6.Multiply the squared deviations (d  $^2$ ) by the respective frequencies and get  $\Sigma fd^{-2}$
- 7. Substituting the values in the following formula to get the standard deviation

$$\sigma = \sqrt{\frac{\sum {fd'}^2}{N} - \left(\frac{\sum {fd'}}{N}\right)^2} \quad \times C$$

## Example 13:

The daily temperature recorded in a city in Russia in a year is given below.

Tempe	erature C <sup>0</sup>	No. of days
-40	to -30	10
-30	to -20	18
-20	to -10	30
-10	to 0	42
0	to 10	65
10	to 20	180
20	to 30	20
		365

Calculate Standard Deviation.

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

#### Solution:

Temperature	Mid value (m)	No. of days f	$\frac{d}{\frac{m-(-5^n)}{10^n}}$	fd	fd <sup>2</sup>
-40 to -30	-35	10	-3	-30	90
-30 to -20	-25	18	-2	-36	72
-20 to -10	-15	30	-1	-30	30
-10 to -0	-5	42	0	0	0
0 to 10	5	65	1	65	65
10 to 20	15	180	2	360	720
20 to 30	25	20	3	60	180
		N=365		$\sum_{\substack{\text{fd} = 389}} fd =$	$\sum fd^2$ =1157

$$\sigma = \sqrt{\frac{\sum fd'^2}{N} - \left(\frac{\sum fd'}{N}\right)^2} \times C$$
  
=  $\sqrt{\frac{1157}{365}} - \left(\frac{389}{365}\right)^2 \times 10$   
=  $\sqrt{3.1699} - 1.1358 \times 10$   
=  $\sqrt{2.0341} \times 10$   
=  $1.4262 \times 10$   
=  $14.26^\circ c$ 

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

## 7.6.6 Merits and Demerits of Standard Deviation: Merits:

- 1. It is rigidly defined and its value is always definite and based on all the observations and the actual signs of deviations are used.
- 2. As it is based on arithmetic mean, it has all the merits of arithmetic mean.
- It is the most important and widely used measure of dispersion.
- 4. It is possible for further algebraic treatment.
- 5. It is less affected by the fluctuations of sampling and hence stable.
- 6. It is the basis for measuring the coefficient of correlation and sampling.

## Demerits:

- 1. It is not easy to understand and it is difficult to calculate.
- 2. It gives more weight to extreme values because the values are squared up.
- 3. As it is an absolute measure of variability, it cannot be used for the purpose of comparison.

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

## 7.6.7 Coefficient of Variation :

The Standard deviation is an absolute measure of dispersion. It is expressed in terms of units in which the original figures are collected and stated. The standard deviation of heights of students cannot be compared with the standard deviation of weights of students, as both are expressed in different units, i.e heights in centimeter and weights in kilograms. Therefore the standard deviation must be converted into a relative measure of dispersion for the purpose of comparison. The relative measure is known as the coefficient of variation.

The coefficient of variation is obtained by dividing the standard deviation by the mean and multiply it by 100. symbolically,

Coefficient of variation (C.V) =  $\frac{\sigma}{\overline{\mathbf{v}}} \times 100$ 

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

If we want to compare the variability of two or more series, we can use C.V. The series or groups of data for which the C.V. is greater indicate that the group is more variable, less stable, less uniform, less consistent or less homogeneous. If the C.V. is less, it indicates that the group is less variable, more stable, more uniform, more consistent or more homogeneous.

## Example 15:

In two factories A and B located in the same industrial area, the average weekly wages (in rupees) and the standard deviations are as follows:

Factory	Average	Standard Deviation	No. of workers
A	34.5	5	476
В	28.5	4.5	524

- 1. Which factory A or B pays out a larger amount as weekly wages?
- 2. Which factory A or B has greater variability in individual wages?

## Solution:

Given  $N_1 = 476$ ,  $\overline{X}_1 = 34.5$ ,  $\sigma_1 = 5$ 

CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IIIBATCH-2017-2020

$$N_2 = 524, \ \overline{X}_2 = 28.5, \ \sigma_2 = 4.5$$

1. Total wages paid by factory A =  $34.5 \times 476$ = Rs.16.422 Total wages paid by factory B =  $28.5 \times 524$ = Rs.14,934.

Therefore factory A pays out larger amount as weekly wages.

2. C.V. of distribution of weekly wages of factory A and B are

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

$$C.V.(A) = \frac{\sigma_1}{\overline{X_1}} \times 100$$
$$= \frac{5}{34.5} \times 100$$
$$= 14.49$$
$$C.V(B) = \frac{\sigma_2}{\overline{X_2}} \times 100$$
$$= \frac{4.5}{28.5} \times 100$$
$$= 15.79$$

Factory B has greater variability in individual wages, since C.V. of factory B is greater than C.V of factory A

## Example 16:

Prices of a particular commodity in five years in two cities are given below:

Price in city A	Price in city B
20	10
22	20
19	18
23	12
16	15

Which city has more stable prices?

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: III BATCH-2017-2020

#### Solution:

Actual mean method

	City A			City B		]
Prices	Deviations	dx <sup>2</sup>	Prices	Deviations	dy <sup>2</sup>	
(X)	from X=20		(Y)	from $Y = 15$		
	dx			dy		
20	0	0	10	-5	25	
22	2	4	20	5	25	
19	-1	1	18	3	9	
23	3	9	12	-3	9	
16	-4	16	15	0	0	
∑x=100	$\sum dx=0$	$\sum dx^2 = 30$	∑y=75	∑dy=0	$\sum dy^2$	
			-		$\sum dy^2 = 68$	

Prepared by K.Aarthiya & R.Gayathri, Asst Prof, Department of Mathematics, KAHE

Page 43/46

CLASS: I B.Com(CA) COUI	SAM ACADEMY OF HIGHER RSE NAME:BUSINESS MATHI	EMATICS AND STATISTICS
RSE CODE: 17CCU202	UNIT: III	BATCH-2017-2020
City A: $\overline{2}$	$\bar{\zeta} = \frac{\Sigma x}{n} = \frac{100}{5} = 20$	
	$x = \sqrt{\frac{\Sigma(x - \overline{x})^2}{n}} = \sqrt{\frac{\Sigma d}{n}}$	$\frac{x^2}{x}$
	$=\sqrt{\frac{30}{5}} = \sqrt{6} = 2.45$	
C.V(x)	$=\frac{\sigma_x}{\overline{x}} \times 100$	
=	$=\frac{2.45}{20} \times 100$	
	20 = 12.25 %	
	$\overline{Z} = \frac{\Sigma y}{n} = \frac{75}{5} = 15$	
$\sigma_y = \sqrt{\frac{\Sigma}{2}}$	$\frac{(y-\overline{y})^2}{n} = \sqrt{\frac{\sum dy^2}{n}}$	
And the second se	$\sqrt{13.6}$ = 3.69	
C.V.(y) = $\frac{\sigma}{y}$	$\frac{y}{y} \ge 100$	
	$\frac{.69}{15}$ ×100	
	15 4.6 %	
2000 Bar 10 10 10 10	more stable prices that	n City B, because the
apafficient of meriati	on is loss in City A	

coefficient of variation is less in City A.

Prepared by K.Aarthiya & R.Gayathri, Asst Prof, Department of Mathematics, KAHE

Page 44/46

RSE CODE: 17CCU20	2	UNIT: III				BATCH	1-2017	-2020	
2 MARK QUESTIO	NS								
I.Find the Arithmetic I	Mean for th	e following d	ata.						
	50 42	95 46							
2.Find Median and Mo		0		0 15	10				
13 16 3.Write the relation bet	17 15	18 14 dand Daviatia		-	12				
4. Calculate the Range					1				
•	12 14	16 18		-					
f :	1 3	5							
5. Define Quartile Dev	iation?								
6. Find the Standard D			-						
240 260 290	245 255	288 27	2 26	53 277	25	1			
8 MARK QUESTIO	NS								
	oth 1 ''	1 reth				C 11			
I.Find median ,quartile		and 56 <sup>th</sup> perc 40-49 5							
Narks: 20-2 No. of stude		40-49 5		60-69 52	70-79 9	9 80 14	)-89 9	0-99	1
NO. OI Stude	ints. 7	11 24		)2	9	14		2	1
2. Calculate the Stand	ard Deviati	on and Coeff		• • •		0			
			cient of	v arianc	e(CV)	tor			
he following data.			cient of	Varianc	e (CV)	for			
X 0-	- 10 10	0 - 20 20	- 30	$\sqrt{30 - 40}$	40 -	- 50			
X 0- f	- 10 10 2	0 - 20 20 5	- 30		40 -				
X   0 -     f   0     3.Calculate the G.M ar	- 10 10 2 d H.M are	) - 20 20 5 9 shown below	- 30	$\frac{30-40}{3}$	40 -	- 50			
$\begin{array}{c c} X & 0 \\ \hline f \\ \hline \end{array}$ 3.Calculate the G.M ar <b>value</b>	- 10 10 2 d H.M are	0 - 20         20           5         0           shown below         0 - 10	- 30 ) -20 2	30 - 40 3	40-	- 50 1 40-5	50		
X   0 -     f   0     3.Calculate the G.M ar	- 10 10 2 d H.M are	0 - 20         20           5         0           shown below         0 - 10	- 30	$\frac{30-40}{3}$	40 -	- 50	50		
X     0 -       f     0       3.Calculate the G.M ar       value       frequency	- 10 10 2 d H.M are	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	- 30 9 - 20 2 2	$     \begin{array}{r}       30 - 40 \\       3     \end{array}     $ 30-30     20	40-	- 50 1 40-5	50		
X     0 -       f     0       3.Calculate the G.M ar       value       frequency       4.Find the Mean deviate	- 10 10 2 d H.M are	$\begin{array}{c cccc} \hline 0 - 20 & 20 \\ \hline 5 & 9 \\ \hline 6 & 9 \\ \hline 6 & 9 \\ \hline 7 & 9 $	- 30 - 30 - 20 2 2 2 2 2 2 2 2 2 2 2 2 2	30 - 40 3 20 - 30 20 lata.	40 - 30-40 6	- 50 1 40-5 4		80 - 89	90 - 99
X     0 -       f     -       3.Calculate the G.M ar       value       frequency       4.Find the Mean deviat       Marks	- 10 10 2 d H.M are	$\begin{array}{c cccc} \hline 0 - 20 & 20 \\ \hline 5 & 9 \\ \hline 6 & 9 \\ \hline 6 & 9 \\ \hline 7 & 9 $	- 30 - 30 - 20 2 2 2 2 2 2 2 2 2 2 2 2 2	$     \begin{array}{r}       30 - 40 \\       3     \end{array}     $ 30-30     20	40-		50 - 79 14	<u>80 - 89</u> 2	<u>90 - 99</u> 1
X     0 -       f     0       3.Calculate the G.M ar       value       frequency       4.Find the Mean deviate	- 10 10 2 d H.M are ions , Med 20 - 29	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	- 30 - 30 - 20 2 - 20 2 - 20 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2	$   \begin{array}{r}     30 - 40 \\     3   \end{array} $ 20-30 20 4ata. 50 - 59	40 - 30-40 6 60 - 6		- 79		
X       0 -         f       -         3.Calculate the G.M ar         value         frequency         4.Find the Mean deviat         Marks         No. of students         5. Calculate the Media	- 10 10 2 d H.M are ions , Med 20 - 29 7	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	- 30 - 30 - 20 2 - 20 2 - 20 2 - 20 2 - 20 2 - 20 - 2 - 20 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2	$   \begin{array}{r}     30 - 40 \\     3   \end{array} $ 20-30 20 lata. 50 - 59 32 Frequence	40 - 30-40 6 6 9 y Distri	- 50 1 40-5 59 70 ibution.	- 79 14	2	
X       0 -         f       -         3.Calculate the G.M ar         value         frequency         4.Find the Mean deviat         Marks         No. of students         5. Calculate the Mediat	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the foi ges (in Rs.	20 $20$ $5$ $20$ $5$ $9$ shown below $0 - 10$ $10$ $0 - 10$ $10$ $10$ $8$ $1$ ian of the foll $30 - 39$ $4$ $11$ $11$ $11$ llowing Cont $0 - 19$ $10 - 19$	- 30 - 30 - 20 2 2 2 2 2 2 2 2 2 2 2 2 2 2	$   \begin{array}{r}     30 - 40 \\     3   \end{array} $ 20-30 20 20 20 20 20 20 20 20 20 20 20 20 20	40 - 30-40 6 6 9 y Distri - 59	-50 1 40-5 40-5 59 70 59 70 50 - 79	- 79 14 80 -	2	
X       0 -         f       -         3.Calculate the G.M ar       - <b>value</b> -         frequency       -         4.Find the Mean deviate       -         Marks       -         No. of students       -         5. Calculate the Media       -         Wa       -         No.       -	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker	20 $20$ $5$ $20$ $5$ $9$ $10$ $10$ $0 - 10$ $10$ $8$ $1$ $10$ $10$ $8$ $1$ $10$ $30 - 39$ $4$ $11$ Illowing Cont $0 - 19$ $rs:$ $5$	- 30 - 30 - 20 2 2 - 20 2 2 - 20 2 - 20 - 49 24 - 20 - 20 - 20 	$   \begin{array}{r}     30 - 40 \\     3   \end{array} $ 20-30 20 lata. 50 - 59 32 Frequenc 39 40 0 3	40 - 30-40 6 6 9 y Distri - 59 85	-50 1 40-5 40-5 59 70 59 70 1000 100 1000 1000 1000 1000 1000 1000 1000 100	- 79 14 80 -	2 • 99 2	
X       0 -         f       0         3.Calculate the G.M ar         value         frequency         4.Find the Mean deviat         Marks         No. of students         5. Calculate the Media         Wa         No.         6. The income distribut	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker	20 $20$ $5$ $20$ $5$ $9$ $10$ $10$ $0 - 10$ $10$ $8$ $1$ $10$ $10$ $8$ $1$ $10$ $30 - 39$ $4$ $11$ Illowing Cont $0 - 19$ $rs:$ $5$	- 30 - 30 - 20 2 2 - 20 2 2 - 20 2 - 20 - 49 24 - 20 - 20 - 20 	$   \begin{array}{r}     30 - 40 \\     3   \end{array} $ 20-30 20 lata. 50 - 59 32 Frequenc 39 40 0 3	40 - 30-40 6 6 9 y Distri - 59 85	-50 1 40-5 40-5 59 70 59 70 1000 100 1000 1000 1000 1000 1000 1000 1000 100	- 79 14 80 -	2 • 99 2	
X       0 -         f       -         3.Calculate the G.M ar         value         frequency         4.Find the Mean deviat         Marks         No. of students         5. Calculate the Media         Wa         No.         6. The income distribut         coefficient.	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker on of 102 p	20 $20$ $5$ $9$ $5$ $9$ $5$ $9$ $10$ $10$ $8$ $1$ $10$ $10$ $8$ $1$ $10$ $10$ $30 - 39$ $4$ $11$ $10$ $0 - 19$ $11$ $0 - 19$ $11$ $10$ $10$ $10$ $11$ $10$ $11$ $10$ $11$ $10$ <	$ \begin{array}{c c} -30 \\ \hline 20 \\ 2 \\ \hline 20 \\ 2 \\ \hline 20 \\ 24 \\ \hline 20 \\ \hline 24 \\ \hline 20 \\ 20 \\ \hline 20 \\ 20 \\ \hline 20 \\ \hline 20 \\ 20 \\ \hline 20 \\ 20 \\ \hline 20 \\ 20 \\ 20 \\ 20 \\ 20 \\ 20 \\ 20 \\ 20 \\$	$   \begin{array}{r}     30 - 40 \\     3   \end{array} $ 20-30 20 lata. 50 - 59 32 Frequence 39 40 0 3 ulate the	40 - 30-40 6 6 9 y Distri - 59 0 35 quartil	-50 1 40-5 4 59 70 ibution. 60 - 79 20 e devia	- 79 14 80 - 11 tion ar	2 - 99 2 nd its	1
X       0 -         f       -         3.Calculate the G.M ar         value         frequency         4.Find the Mean deviate         Marks         No. of students         5. Calculate the Media         Wa         No.         6.The income distribut         coefficient.         Income (Rs.	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker on of 102 p	20 $20$ $5$ $20$ $5$ $9$ $5$ $10$ $0 - 10$ $10$ $8$ $1$ $10$ $30 - 39$ $4$ $11$ $10$ $10$ $30 - 39$ $4$ $11$ $10$ $0 - 19$ $rs:$ $5$ persons is given between the second s	$ \begin{array}{c c} -30 \\ \hline 20 \\ 2 \\ \hline 2 \\ \hline 2 \\ \hline 0 - 49 \\ \hline 24 \\ \hline 10 \\ \hline 20 \\ 20 \\ \hline 20 \\ 20 \\ \hline 20 \\ 20 \\ \hline 20 \\ 20 \\ 20 \\ 20 \\ 20 \\ 20 \\ 20 \\ 20 \\$	$     \begin{array}{r}       30 - 40 \\       3     \end{array}     $ 20-30     20     20 lata. 50 - 59 32 Frequence 39     40 0	40 - 30-40 6 6 9 y Distri - 59 9 guartil 00	-50 1 40-5 4 59 70 50 - 79 20 e devia 350	- 79 14 80 - 11 tion ar 400	2 • 99 2 nd its 0 70	1
X       0 -         f       0         3.Calculate the G.M ar         value         frequency         4.Find the Mean deviat         Marks         No. of students         5. Calculate the Media:         Wa         No.         6. The income distribut coefficient.         Income (Rs.         No.of p	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker on of 102 p ersons	20 $20$ $5$ $20$ $5$ $9$ $6$ $10$ $10$ $10$ $8$ $1$ $10$ $10$ $8$ $1$ $10$ $10$ $30 - 39$ $4$ $11$ $10$ llowing Cont $0 - 19$ $11$ $10$ $10$ $10$	$ \begin{array}{c c} -30 \\ \hline 20 \\ \hline 22 \\ \hline 2 \\ \hline 0 - 49 \\ \hline 24 \\ \hline 100 \\ \hline 20 \\ \hline 10 \\ \hline 10 \\ \hline 20 \\ \hline 10 \\ \hline 1$	$     \begin{array}{r}       30 - 40 \\       3     \end{array}     $ $       30 - 40 \\       3     \end{array}     $ $       30 - 40 \\       30 - 30 \\       32     \end{array}     $ $       \begin{array}{r}       30 - 40 \\       30 - 30 \\       30$	40 - 30-40 6 9 y Distri - 59 9 9 9 9 9 9 9 9 9 9 9 9 9	-50 1 40-5 40-5 4 59 70 1000 1000 1000 1000 1000 1000 1000 1000 1000 1000 1000 1000	- 79 14 80 - 11 tion ar	2 • 99 2 nd its 0 70	1
X       0 -         f       -         3.Calculate the G.M ar       - <b>arks</b> -         4.Find the Mean deviat       -         Marks       No. of students         5. Calculate the Media       Wa         No.       -         6. The income distribut coefficient.       -         Income (Rs.       No. of p         7.Calculate the Arithm       -	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker on of 102 j ersons etic Mean ,	20 $20$ $5$ $20$ $5$ $20$ $5$ $20$ $0 - 10$ $10$ $8$ $1$ $10$ $8$ $10$ $30 - 39$ $4$ $11$ $10$ $0 - 19$ $rs:$ $5$ persons is given berown is given berow	$ \begin{array}{c c} -30 \\ \hline -30 \\ \hline 20 \\ 2 \\ \hline 20 \\ 2 \\ \hline 20 \\ \hline 24 \\ \hline 20 \\ \hline 16 \\ \hline 16 \\ \hline node for \\ \hline \end{array} $	$     \begin{array}{r}       30 - 40 \\       3     \end{array}     $ 20-30     20     20 lata. 50 - 59 32 Frequence 39       40 0       30 ulate the 0       30 0       30 0       40 0       30 0       30	40 - 30-40 6 6 9 y Distri - 59 0 35 quartil 00 0 0 0 0 0 0	-50 1 40-5 4 59 70 50 - 79 20 e devia 350 21 lata.	- 79 14 80 - 11 tion ar 400 15	2 • 99 2 nd its 0 70	1
X       0 -         f       -         3.Calculate the G.M ar       - <b>arks</b> -         4.Find the Mean deviat       -         Marks       No. of students         5. Calculate the Media       Wa         No.       -         6. The income distribut coefficient.       -         Income (Rs.       No. of p         7.Calculate the Arithm       -	- 10 10 2 d H.M are ions , Med 20 - 29 7 n for the fo ges (in Rs. of Worker on of 102 p ersons	20 $20$ $5$ $20$ $5$ $10$ $0 - 10$ $10$ $8$ $1$ $10$ $30 - 39$ $4$ $11$ $10$ $10$ $30 - 39$ $4$ $11$ $10$ $10$ $10$ $10$ $10$ $10$ $10$ $10$ $10$ $10$ $10$ $11$ $162$	$ \begin{array}{c c} -30 \\ \hline 20 \\ \hline 22 \\ \hline 2 \\ \hline 0 - 49 \\ \hline 24 \\ \hline 100 \\ \hline 20 \\ \hline 10 \\ \hline 10 \\ \hline 20 \\ \hline 10 \\ \hline 1$	$     \begin{array}{r}       30 - 40 \\       3     \end{array}     $ $       30 - 40 \\       3     \end{array}     $ $       30 - 40 \\       30 - 30 \\       32     \end{array}     $ $       \begin{array}{r}       30 - 40 \\       30 - 30 \\       30$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	-50 1 40-5 4 59 70 50 - 79 20 e devia 350 21 lata.	- 79 14 80 - 11 tion ar 400	2 • 99 2 nd its 0 70	1

#### **KARPAGAM ACADEMY OF HIGHER EDUCATION** CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 BATCH-2017-2020 **UNIT: III** 8. Calculate the Coefficient of Variance for the following data. 9.Calculate the Median for the following. 40 - 50 50 - 6060 - 70 70 - 80 80 - 90 Hourly Wages (in Rs.) 90 - 100 **Number of Employees** 10. The following data give the details about salaries (in thousands of rupees) of seven employees randomly selected from a Pharmaceutical Company. Serial No. Salary per Annum ( '000) Calculate the Standard Deviation and Coefficient of variance of the given data. 11. i) Calculate the range and its coefficient for the following data. ii) Calculate the standard deviation for the following data. X:20f : 5 12. Calculate the Coefficient of Variation for the following data. Х F

#### CLASS: I BBA COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 18BAU202 UNIT: IV BATCH-2018-2021

#### <u>UNIT – IV</u>

#### SYLLABUS

#### **Bi-variate Analysis**

Simple Linear Correlation Analysis: Meaning, and measurement. Karl Pearson's co-efficient and Spearman's rank correlation Simple Linear Regression Analysis: Regression equations and estimation. Relationship between correlation and regression coefficients

# CORRELATION

#### Introduction:

The term correlation is used by a common man without knowing that he is making use of the term correlation. For example when parents advice their children to work hard so that they may get good marks, they are correlating good marks with hard work.

Thus Correlation refers to the relationship of two variables or more. (e-g) relation between height of father and son, yield and rainfall, wage and price index, share and debentures etc.

#### Definitions:

- Correlation Analysis attempts to determine the degree of relationship between variables- Ya-Kun-Chou.
- Correlation is an analysis of the covariation between two or more variables.- A.M.Tuttle.

Correlation expresses the inter-dependence of two sets of variables upon each other. One variable may be called as (subject)

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

independent and the other relative variable (dependent). Relative variable is measured in terms of subject.

## Uses of correlation:


- 1. It is used in physical and social sciences.
- It is useful for economists to study the relationship between variables like price, quantity etc. Businessmen estimates costs, sales, price etc. using correlation.
- It is helpful in measuring the degree of relationship between the variables like income and expenditure, price and supply, supply and demand etc.
- 4. Sampling error can be calculated.
- 5. It is the basis for the concept of regression.

## Scatter Diagram:

It is the simplest method of studying the relationship between two variables diagrammatically. One variable is represented along the horizontal axis and the second variable along the vertical axis. For each pair of observations of two variables, we put a dot in the plane. There are as many dots in the plane as the number of paired observations of two variables. The direction of dots shows the scatter or concentration of various points. This will show the type of correlation.


1. If all the plotted points form a straight line from lower left hand corner to the upper right hand corner then there is

Perfect positive correlation. We denote this as r = +1




CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

- 1. If all the plotted dots lie on a straight line falling from upper left hand corner to lower right hand corner, there is a perfect negative correlation between the two variables. In this case the coefficient of correlation takes the value r = -1.
- If the plotted points in the plane form a band and they show a rising trend from the lower left hand corner to the upper right hand corner the two variables are highly positively correlated.



- 1. If the points fall in a narrow band from the upper left hand corner to the lower right hand corner, there will be a high degree of negative correlation.
- If the plotted points in the plane are spread all over the diagram there is no correlation between the two variables.



Ο

CLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

## Merits:

- 1. It is a simplest and attractive method of finding the nature of correlation between the two variables.
- 2. It is a non-mathematical method of studying correlation. It is easy to understand.
- 3. It is not affected by extreme items.
- 4. It is the first step in finding out the relation between the two variables.
- 5. We can have a rough idea at a glance whether it is a positive correlation or negative correlation.

## Demerits:

By this method we cannot get the exact degree or correlation between the two variables.

## Types of Correlation:

Correlation is classified into various types. The most important ones are

- i) Positive and negative.
- ii) Linear and non-linear.
- iii) Partial and total.
- iv) Simple and Multiple.

## Positive and Negative Correlation:

It depends upon the direction of change of the variables. If the two variables tend to move together in the same direction (ie) an increase in the value of one variable is accompanied by an increase in the value of the other, (or) a decrease in the value of one variable is accompanied by a decrease in the value of other, then the correlation is called positive or direct correlation. Price and supply, height and weight, yield and rainfall, are some examples of positive correlation.

If the two variables tend to move together in opposite directions so that increase (or) decrease in the value of one variable is accompanied by a decrease or increase in the value of the other variable, then the correlation is called negative (or) inverse correlation. Price and demand, yield of crop and price, are examples of negative correlation.

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

#### Linear and Non-linear correlation:

If the ratio of change between the two variables is a constant then there will be linear correlation between them.

Consider the following.

X	2	4	6	8	10	12
Y	3	6	9	12	15	18

Here the ratio of change between the two variables is the same. If we plot these points on a graph we get a straight line.

If the amount of change in one variable does not bear a constant ratio of the amount of change in the other. Then the relation is called Curvi-linear (or) non-linear correlation. The graph will be a curve.

## Simple and Multiple correlation:

When we study only two variables, the relationship is simple correlation. For example, quantity of money and price level, demand and price. But in a multiple correlation we study more than two variables simultaneously. The relationship of price, demand and supply of a commodity are an example for multiple correlation.

## Partial and total correlation:

The study of two variables excluding some other variable is called **Partial correlation**. For example, we study price and demand eliminating supply side. In total correlation all facts are taken into account.

## Computation of correlation:

When there exists some relationship between two variables, we have to measure the degree of relationship. This measure is called the measure of correlation (or) correlation coefficient and it is denoted by 'r'.

## Co-variation:

The covariation between the variables x and y is defined as  $Cov(x,y) = \frac{\Sigma(x-\overline{x})(y-\overline{y})}{n}$  where  $\overline{x}, \overline{y}$  are respectively means of x and y and 'n' is the number of pairs of observations.

Page 5/37

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

## Karl pearson's coefficient of correlation:

Karl pearson, a great biometrician and statistician, suggested a mathematical method for measuring the magnitude of linear relationship between the two variables. It is most widely used method in practice and it is known as pearsonian coefficient of correlation. It is denoted by 'r'. The formula for calculating 'r' is

(i)  $r = \frac{C \operatorname{ov}(x, y)}{\sigma_x \cdot \sigma_y}$  where  $\sigma_x$ ,  $\sigma_y$  are S.D of x and y respectively. (ii)  $r = \frac{\sum xy}{n \ \sigma_x \ \sigma_y}$ (iii)  $r = \frac{\sum XY}{\sqrt{\sum X^2 \cdot \sum Y^2}}$ ,  $X = x - \overline{x}$ ,  $Y = y - \overline{y}$ 

when the deviations are taken from the actual mean we can apply any one of these methods. Simple formula is the third one.

The third formula is easy to calculate, and it is not necessary to calculate the standard deviations of x and y series respectively.

#### Steps:

- 1. Find the mean of the two series x and y.
- 2. Take deviations of the two series from x and y.

$$X = x - x$$
,  $Y = y - y$ 

- 3. Square the deviations and get the total, of the respective squares of deviations of x and y and denote by  $\Sigma X^2$ ,  $\Sigma Y^2$  respectively.
- 4. Multiply the deviations of x and y and get the total and Divide by n. This is covariance.
- 5. Substitute the values in the formula.

$$r = \frac{\operatorname{cov}(x, y)}{\sigma x. \sigma y} = \frac{\sum (x - \overline{x}) (y - \overline{y})/n}{\sqrt{\frac{\sum (x - \overline{x})^2}{n}} \sqrt{\frac{\sum (y - \overline{y})^2}{n}}}$$

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

The above formula is simplified as follows

$$r = \frac{\Sigma XY}{\sqrt{\Sigma X^2 \cdot \Sigma Y^2}}, \quad X = x - \overline{x}, Y = y - \overline{y}$$

#### Example 1:

Find Karl Pearson's coefficient of correlation from the following data between height of father (x) and son (y).

X	64	65	66	67	68	69	70
Y	66	67	65	68	70	68	72

Comment on the result.

Solution:

X	Y	$X = x - \overline{x}$	$X^2$	$Y = y - \overline{y}$	$Y^2$	XY
	s	X = x - 67		$\mathbf{Y} = \mathbf{y} - 68$		
64	66	-3	9	-2	4	6
65	67	-2	4	-1	1	2
66	65	-1	1	-3	9	3
67	68	0	0	0	0	0
68	70	1	1	2	4	2
69	68	2	4	0	0	0
70	72	3	9	4	16	12
469	476	0	28	0	34	25
10000000000	$\frac{169}{7} = 6$		= 68	20.42		
123	$\Sigma$	XY :	25	25	25	0.01
<i>r</i> =	$\sqrt{\sum \mathbf{v}^2}$	$\sum Y^2 = \sqrt{28}$	× 34	$= \sqrt{952}$	30.85	= 0.81

Since r = +0.81, the variables are highly positively correlated. (ie) Tall fathers have tall sons.

#### Working rule (i)

We can also find r with the following formula

We have 
$$r = \frac{C \operatorname{ov}(x, y)}{\sigma_x \cdot \sigma_y}$$
  
 $\operatorname{Cov}(x, y) = \frac{\Sigma(x - \overline{x})(y - \overline{y})}{n} = \frac{\Sigma(xy + \overline{x}y - \overline{y}x - \overline{x}\overline{y})}{n}$ 

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

$$= \frac{\Sigma xy}{n} - \frac{\overline{y}\Sigma x}{n} - \frac{\overline{x}\Sigma y}{n} + \frac{\Sigma \overline{x}\overline{y}}{n}$$

$$Cov(x,y) = \frac{\Sigma xy}{n} - \overline{y}\overline{x} - \overline{x}\overline{y} + \overline{y}\overline{y} = \frac{\Sigma xy}{n} - \overline{x}\overline{y}$$

$$\varpi^{2}x^{2} = \frac{\Sigma x^{2}}{n} - \overline{x}^{2}, \quad \overline{\varpi}^{2}y^{2} = \frac{\Sigma y^{2}}{n} - \overline{y}^{2}$$

$$Now \ r = -\frac{Cov(x,y)}{\sigma_{x}.\sigma_{y}}$$

$$r = \frac{\frac{\Sigma xy}{n} - \overline{x}\overline{y}}{\sqrt{\left(\frac{\Sigma x^{2}}{n} - \overline{x}^{2}\right)} \cdot \sqrt{\left(\frac{\Sigma y^{2}}{n} - \overline{y}^{2}\right)}}$$

$$r = \frac{n\Sigma xy - (\Sigma x)(\Sigma y)}{\sqrt{[n\Sigma x^{2} - (\Sigma x)^{2}][n\Sigma y^{2} - (\Sigma y)^{2}]}}$$

**Note:** In the above method we need not find mean or standard deviation of variables separately.

#### Example 2:

Calculate coefficient of correlation from the following data.

Х	1	2	3	4	5	6	7	8	9	
Y	9	8	10	12	11	13	14	16	15	
X		N	1	X	2	v	2	Xy	7	
1		5	)	1		8	1	9		
2		8	3	4	21	64		16	5	
		1	0	9		100		30		
4		1.	2	16		144		48		
5		1	1	25		121		55		
6		1	3	36		169		78		
7		1	4	49		196		98		
8		1	6	64		25	256 128			
9		1.	5	81		225		135		
45		108	3	285		135	6	597		

■ ]

	EMY OF HIGHER EDUCATION	
CLASS: I B.Com(CA) COURSE NAME: COURSE CODE: 17CCU202 UN		ND STATISTICS TCH-2017-2020

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

$$r = \frac{n\Sigma xy - (\Sigma x) (\Sigma y)}{\sqrt{[n\Sigma x^2 - (\Sigma x)^2][n\Sigma y^2 - (\Sigma y)^2]}}$$

$$r = \frac{9 \times 597 - 45 \times 108}{\sqrt{(9 \times 285 - (45)^2).(9 \times 1356 - (108)^2)}}$$

$$r = \frac{5373 - 4860}{\sqrt{(2565 - 2025).(12204 - 11664)}}$$

$$= \frac{513}{\sqrt{540 \times 540}} = \frac{513}{540} = 0.95$$

## Working rule (ii) (shortcut method)

We have 
$$r = \frac{C \operatorname{ov}(x, y)}{\sigma_x \cdot \sigma_y}$$
  
where  $\operatorname{Cov}(x, y) = \frac{\Sigma(x - \overline{x})(y - \overline{y})}{n}$ 

Take the deviation from x as x - A and the deviation from y as y - B

$$Cov(x,y) = \frac{\sum [(x-A) \cdot (\overline{x} - A)] [(y-B) \cdot (\overline{y} - B)]}{n}$$
  
=  $\frac{1}{n} \sum [(x-A) (y-B) \cdot (x-A) (\overline{y} - B)]$   
 $\cdot (\overline{x} - A)(y-B) + (\overline{x} - A)(\overline{y} - B)]$   
=  $\frac{1}{n} \sum [(x-A) (y-B) \cdot (\overline{y} - B) \frac{\sum (x-A)}{n}$   
 $- (\overline{x} - A) \frac{\sum (y-B)}{n} + \frac{\sum (\overline{x} - A)(\overline{y} - B)}{n}$   
=  $\frac{\sum (x-A)(y-B)}{n} - (\overline{y} - B) (\overline{x} - \frac{nA}{n})$   
 $- (\overline{x} - A) (\overline{y} - \frac{nB}{n}) + (\overline{x} - A) (\overline{y} - B)$ 

Page 10/37

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

$$= \frac{\sum(x - A)(y - B)}{n} - (\overline{y} - B)(\overline{x} - A)$$
$$- (\overline{x} - A)(\overline{y} - B) + (\overline{x} - A)(\overline{y} - B)$$
$$= \frac{\sum(x - A)(y - B)}{n} - (\overline{x} - A)(\overline{y} - B)$$
Let  $x - A = u$ ;  $y - B = v$ ;  $\overline{x} - A = \overline{u}$ ;  $\overline{y} - B = \overline{v}$ 
$$\therefore \operatorname{Cov}(x, y) = \frac{\sum uv}{n} - \overline{uv}$$
$$\operatorname{cox}_{x}^{2} = \frac{\sum u^{2}}{n} - \overline{u^{2}} = \sigma u^{2}$$
$$\sum v^{2} = c$$

$$\sigma \mathfrak{P}_{y}^{2} = \frac{\Sigma v^{2}}{n} - \overline{v}^{2} = \sigma v^{2}$$
  
$$\therefore r = \frac{n\Sigma uv - (\Sigma u)(\Sigma v)}{\sqrt{\left[n\Sigma u^{2} - (\Sigma u)^{2}\right] \cdot \left[(n\Sigma v^{2}) - (\Sigma v)^{2}\right]}}$$

## Example 3:

]

## Calculate Pearson's Coefficient of correlation.

Х	45	55	56	58	60	65	68	70	75	80	85
Y	56	50	48	60	62	64	65	70	74	82	90

Х	Y	u = x-A	v = y-B	$u^2$	$v^2$	uv
45	56	-20	-14	400	196	280
55	50	-10	-20	100	400	200
56	48	-9	-22	81	484	198
58	60	-7	-10	49	100	70
60	62	-5	-8	25	64	40
65	64	0	-6	0	36	0
68	65	3	-5	9	25	-15
70	70	5	0	25	0	0
75	74	10	4	100	16	40
80	82	15	12	225	144	180
85	90	20	20	400	400	400
		2	-49	1414	1865	1393

CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

$$r = \frac{n\Sigma uv - (\Sigma u) (\Sigma v)}{\sqrt{[n\Sigma u^2 - (\Sigma u^2)] [n\Sigma v^2 - (\Sigma v)^2]}}$$

$$r = \frac{11 \times 1393 - 2 \times (-49)}{\sqrt{(1414 \times 11 - (2)^2) \times (1865 \times 11 - (-49)^2)}}$$

$$= \frac{15421}{\sqrt{15550 \times 18114}} = \frac{15421}{16783.11} = +0.92$$

# Limitations:

- Correlation coefficient assumes linear relationship regardless of the assumption is correct or not.
- Extreme items of variables are being unduly operated on correlation coefficient.
- Existence of correlation does not necessarily indicate causeeffect relation.

# Interpretation:

The following rules helps in interpreting the value of 'r'.

- 1. When r = 1, there is perfect +ve relationship between the variables.
- When r = -1, there is perfect –ve relationship between the variables.
- 3. When r = 0, there is no relationship between the variables.
- If the correlation is +1 or −1, it signifies that there is a high degree of correlation. (+ve or -ve) between the two variables.
- If r is near to zero (ie) 0.1,-0.1, (or) 0.2 there is less correlation.

KARPAGAM	ACADEMY OF HIGHEI	REDUCATION
CLASS: I B.Com(CA) COURSE I COURSE CODE: 17CCU202	NAME:BUSINESS MATI UNIT: IV	HEMATICS AND STATISTICS BATCH-2017-2020

# Rank Correlation:

It is studied when no assumption about the parameters of the population is made. This method is based on ranks. It is useful to study the qualitative measure of attributes like honesty, colour, beauty, intelligence, character, morality etc. The individuals in the group can be arranged in order and there on, obtaining for each individual a number showing his/her rank in the group. This method was developed by Edward Spearman in 1904. It is defined

as  $r = 1 - \frac{6\Sigma D^2}{n^3 - n}$  r = rank correlation coefficient.

Note: Some authors use the symbol  $\rho$  for rank correlation.  $\Sigma D^2 = \text{sum of squares of differences between the pairs of ranks.}$ n = number of pairs of observations.

The value of r lies between -1 and +1. If r = +1, there is complete agreement in order of ranks and the direction of ranks is also same. If r = -1, then there is complete disagreement in order of ranks and they are in opposite directions.

Computation for tied observations: There may be two or more items having equal values. In such case the same rank is to be given. The ranking is said to be tied. In such circumstances an average rank is to be given to each individual item. For example if the value so is repeated twice at the 5<sup>th</sup> rank, the common rank to be assigned to each item is  $\frac{5+6}{2} = 5.5$  which is the average of 5 and 6 given as 5.5, appeared twice.

If the ranks are tied, it is required to apply a correction factor which is  $\frac{1}{12}$  (m<sup>3</sup>-m). A slightly different formula is used when there is more than one item having the same value.

The formula is

r = 
$$1 - \frac{6[\Sigma D^2 + \frac{1}{12}(m^3 - m) + \frac{1}{12}(m^3 - m) + \dots]}{n^3 - n}$$

KARPAGAM	ACADEMY OF HIGHEI	REDUCATION
CLASS: I B.Com(CA) COURSE I COURSE CODE: 17CCU202	NAME:BUSINESS MATI UNIT: IV	HEMATICS AND STATISTICS BATCH-2017-2020

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

Where m is the number of items whose ranks are common and should be repeated as many times as there are tied observations.

# Example 6:

In a marketing survey the price of tea and coffee in a town based on quality was found as shown below. Could you find any relation between and tea and coffee price.

Price of tea	88	90	95	70	60	75	50
Price of coffee	120	134	150	115	110	140	100

Price of	Rank	Price of	Rank	D	$D^2$
tea		coffee			
88	3	120	4	1	1
90	2	134	3	1	1
95	1	150	1	0	0
70	5	115	5	0	0
60	6	110	6	0	0
75	4	140	2	2	4
50	7	100	7	0	0
					$\Sigma D^2 = 6$

$$r = 1 - \frac{6\Sigma D^2}{n^3 - n} = 1 - \frac{6 \times 6}{7^3 - 7}$$
$$= 1 - \frac{36}{7^3 - 7} = 1 - 0.1071$$

$$=1-\frac{1}{336}$$
 = 1-0

= 0.8929

The relation between price of tea and coffee is positive at 0.89. Based on quality the association between price of tea and price of coffee is highly positive.

# Example 7:

In an evaluation of answer script the following marks are awarded by the examiners.

$1^{st}$	88	95	70	960	50	80	75	85
$2^{nd}$	84	90	88	55	48	85	82	72

Page 16/37

CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV

BATCH-2017-2020

Do you agree the evaluation by the two examiners is fair?

Do you agree the evaluation by the two examiners is fair :								
Х	R1	У	R2	D	$D^2$			
88	2	84	4	2	4			
95	1	90	1	0	0			
70	6	88	2	4	16			
60	7	55	7	0	0			
50	8	48	8	0	0			
80	4	85	3	1	1			
85	3	75	6	3	9			
					30			

$$r = 1 - \frac{6\Sigma D^2}{n^3 - n} = 1 - \frac{6 \times 30}{8^3 - 8}$$
$$= 1 - \frac{180}{8^3 - 8} = 1 - \frac{0.357}{8^3 - 8} = 0$$

$$= 1 - \frac{180}{504} = 1 - 0.357 = 0.643$$

r = 0.643 shows fair in awarding marks in the sense that uniformity has arisen in evaluating the answer scripts between the two examiners.

# Example 8:

Rank Correlation for tied observations. Following are the marks obtained by 10 students in a class in two tests.

Students	Α	В	С	D	E	F	G	Η	Ι	J
Test 1	70	68	67	55	60	60	75	63	60	72
Test 2	65	65	80	60	68	58	75	63	60	70


Calculate the rank correlation coefficient between the marks of two tests.

Student	Test 1	R1	Test 2	R2	D	$D^2$
А	70	3	65	5.5	-2.5	6.25
В	68	4	65	5.5	-1.5	2.25
С	67	5	80	1.0	4.0	16.00
D	55	10	60	8.5	1.5	2.25
Е	60	8	68	4.0	4.0	16.00
F	60	8	58	10.0	-2.0	4.00
G	75	1	75	2.0	-1.0	1.00
Н	63	6	62	7.0	-1.0	1.00
Ι	60	8	60	8.5	0.5	0.25
J	72	2	70	3.0	-1.0	1.00
						50.00

Page 17/37

# CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

60 is repeated 3 times in test 1. 60,65 is repeated twice in test 2. m = 3; m = 2; m = 2 $\mathbf{r} = 1 - \frac{6[\Sigma D^2 + \frac{1}{12}(m^3 - m) + \frac{1}{12}(m^3 - m) + \frac{1}{12}(m^3 - m)}{n^3 - n}$  $= 1 - \frac{6[50 + \frac{1}{12}(3^3 - 3) + \frac{1}{12}(2^3 - 2) + \frac{1}{12}(2^3 - 2)]}{10^3 - 10}$  $= 1 - \frac{6[50 + 2 + 0.5 + 0.5]}{990}$  $= 1 - \frac{6 \times 53}{990} = \frac{672}{990} = 0.68$ 



CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

# 9.1 Introduction:

After knowing the relationship between two variables we may be interested in estimating (predicting) the value of one variable given the value of another. The variable predicted on the basis of other variables is called the "dependent" or the 'explained' variable and the other the 'independent' or the 'predicting' variable. The prediction is based on average relationship derived statistically by regression analysis. The equation, linear or otherwise, is called the regression equation or the explaining equation.

For example, if we know that advertising and sales are correlated we may find out expected amount of sales for a given advertising expenditure or the required amount of expenditure for attaining a given amount of sales.

The relationship between two variables can be considered between, say, rainfall and agricultural production, price of an input and the overall cost of product, consumer expenditure and disposable income. Thus, regression analysis reveals average relationship between two variables and this makes possible estimation or prediction.

# 9.1.1 Definition:

Regression is the measure of the average relationship between two or more variables in terms of the original units of the data.

# 9.2 Types Of Regression:

The regression analysis can be classified into:

- a) Simple and Multiple
- b) Linear and Non-Linear
- c) Total and Partial

# a) Simple and Multiple:

In case of simple relationship only two variables are considered, for example, the influence of advertising expenditure on sales turnover. In the case of multiple relationship, more than

# CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

two variables are involved. On this while one variable is a dependent variable the remaining variables are independent ones.

For example, the turnover (y) may depend on advertising expenditure (x) and the income of the people (z). Then the functional relationship can be expressed as y = f(x,z).

# b) Linear and Non-linear:

The linear relationships are based on straight-line trend, the equation of which has no-power higher than one. But, remember a linear relationship can be both simple and multiple. Normally a linear relationship is taken into account because besides its simplicity, it has a better predective value, a linear trend can be easily projected into the future. In the case of non-linear relationship curved trend lines are derived. The equations of these are parabolic.

# c) Total and Partial:

In the case of total relationships all the important variables are considered. Normally, they take the form of a multiple relationships because most economic and business phenomena are affected by multiplicity of cases. In the case of partial relationship one or more variables are considered, but not all, thus excluding the influence of those not found relevant for a given purpose.

# 9.3 Linear Regression Equation:

If two variables have linear relationship then as the independent variable (X) changes, the dependent variable (Y) also changes. If the different values of X and Y are plotted, then the two straight lines of best fit can be made to pass through the plotted points. These two lines are known as regression lines. Again, these regression lines are based on two equations known as regression equations. These equations show best estimate of one variable for the known value of the other. The equations are linear.

Linear regression equation of Y on X is

Y = a + bX .....(1)

 $\mathbf{X} = a + b\mathbf{Y}.\dots(2)$ 

*a*, *b* are constants.


# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

From (1) We can estimate Y for known value of X. (2) We can estimate X for known value of Y.


# 9.3.1 Regression Lines:

For regression analysis of two variables there are two regression lines, namely Y on X and X on Y. The two regression lines show the average relationship between the two variables.

For perfect correlation, positive or negative i.e.,  $r = \pm 1$ , the two lines coincide i.e., we will find only one straight line. If r = 0, i.e., both the variables are independent then the two lines will cut each other at right angle. In this case the two lines will be parallel to X and Y-axes.



Lastly the two lines intersect at the point of means of X and Y. From this point of intersection, if a straight line is drawn on X-axis, it will touch at the mean value of x. Similarly, a perpendicular drawn from the point of intersection of two regression lines on Y-axis will touch the mean value of Y.



KARPAGAM	ACADEMY OF HIGHEI	REDUCATION
CLASS: I B.Com(CA) COURSE I COURSE CODE: 17CCU202	NAME:BUSINESS MATI UNIT: IV	HEMATICS AND STATISTICS BATCH-2017-2020

CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

# 9.3.2 Principle of 'Least Squares':

Regression shows an average relationship between two variables, which is expressed by a line of regression drawn by the method of "least squares". This line of regression can be derived graphically or algebraically. Before we discuss the various methods let us understand the meaning of least squares.

A line fitted by the method of least squares is known as the line of best fit. The line adapts to the following rules:

 The algebraic sum of deviation in the individual observations with reference to the regression line may be equal to zero. i.e.,

 $\Sigma(X - Xc) = 0$  or  $\Sigma(Y - Yc) = 0$ 

Where Xc and Yc are the values obtained by regression analysis.


(ii) The sum of the squares of these deviations is less than the sum of squares of deviations from any other line. i.e.,  $\Sigma(Y - Yc)^2 \le \Sigma (Y - Ai)^2$ 

Where Ai = corresponding values of any other straight line.

(iii) The lines of regression (best fit) intersect at the mean values of the variables X and Y, i.e., intersecting point is  $\overline{x, y}$ .

# 9.4 Methods of Regression Analysis:

The various methods can be represented in the form of chart given below:



# KARPAGAM ACADEMY OF HIGHER EDUCATIONCLASS: I B.Com(CA)COURSE NAME:BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

# 9.4.1 Graphic Method: Scatter Diagram:

Under this method the points are plotted on a graph paper representing various parts of values of the concerned variables. These points give a picture of a scatter diagram with several points spread over. A regression line may be drawn in between these points either by free hand or by a scale rule in such a way that the squares of the vertical or the horizontal distances (as the case may be) between the points and the line of regression so drawn is the least. In other words, it should be drawn faithfully as the line of best fit leaving equal number of points on both sides in such a manner that the sum of the squares of the distances is the best.

# 9.4.2 Algebraic Methods:

(i) Regression Equation. The two regression equations for X on Y; X = a + bY And for Y on X; Y = a + bX Where X, Y are variables, and a,b are constants whose values are to be determined

For the equation, X = a + bYThe normal equations are  $\sum X = na + b \sum Y$  and  $\sum XY = a\sum Y + b\sum Y^2$ For the equation, Y = a + bX, the normal equations are  $\sum Y = na + b\sum X$  and  $\sum XY = a\sum X + b\sum X^2$ 

From these normal equations the values of a and b can be determined.

# Example 1:

Find the two regression equations from the following data:

X:	6	2	10	4	8
Y:	9	11	5	8	7

Prepared by K. Aarthiya & R. Gayathri, Asst Prof, Department of Mathematics, KAHE

Page 25/37

#### CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

# Solution:

X	Y	$X^2$	$Y^2$	XY
6	9	36	81	54
2	11	4	121	22
10	5	100	25	50
4	8	16	64	32
8	7	64	49	56
30	40	220	340	214

Regression equation of Y on X is Y = a + bX and the normal equations are

 $\sum Y = na + b\sum X$   $\sum XY = a\sum X + b\sum X^{2}$ Substituting the values, we get  $40 = 5a + 30b \dots (1)$   $214 = 30a + 220b \dots (2)$ Multiplying (1) by 6  $240 = 30a + 180b \dots (3)$  (2) - (3) - 26 = 40bor  $b = -\frac{26}{40} = -0.65$ Now, substituting the value of 'b' in equation (1)

$$40 = 5a - 19.5$$
  

$$5a = 59.5$$
  

$$a = \frac{59.5}{5} = 11.9$$

Hence, required regression line Y on X is Y = 11.9 - 0.65 X. Again, regression equation of X on Y is

X = a + bY and

The normal equations are

 $\sum X = na + b \sum Y$  and  $\sum XY = a \sum Y + b \sum Y^2$ 

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

Now, substituting the corresponding values from the above table, we get

 $30 = 5a + 40b \dots (3)$   $214 = 40a + 340b \dots (4)$  **Multiplying (3) by 8, we get**   $240 = 40a + 320 b \dots (5)$  (4) - (5) gives -26 = 20b  $b = -\frac{26}{20} = -1.3$ Substituting b = -1.3 in equation (3) gives 30 = 5a - 52 5a = 82 $a = \frac{82}{5} = 16.4$ 

Hence, Required regression line of X on Y is X = 16.4 - 1.3Y(ii) Regression Co-efficients:

The regression equation of Y on X is  $y_e = \overline{y} + r \frac{\sigma_y}{\sigma_x} (x - \overline{x})$ 

Here, the regression Co.efficient of Y on X is

$$b_{1} = b_{yx} = r \frac{\sigma_{y}}{\sigma_{x}}$$
$$y_{e} = \overline{y} + b_{1}(x - \overline{x})$$

The regression equation of X on Y is

$$X_e = \overline{x} + r \frac{\sigma_x}{\sigma_y} (y - \overline{y})$$

Here, the regression Co-efficient of X on Y

$$b_2 = b_{xy} = r \frac{\sigma_x}{\sigma_y}$$
$$X_e = \overline{X} + b_2(y - \overline{y})$$

Prepared by K. Aarthiya & R. Gayathri, Asst Prof, Department of Mathematics, KAHE

Page 27/37

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

#### If the deviation are taken from respective means of x and y

$$b_{1} = b_{yx} = \frac{\sum (X - \overline{X})(Y - \overline{Y})}{\sum (X - \overline{X})^{2}} = \frac{\sum xy}{\sum x^{2}} \text{ and}$$
$$b_{2} = b_{xy} = \frac{\sum (X - \overline{X})(Y - \overline{Y})}{\sum (Y - \overline{Y})^{2}} = \frac{\sum xy}{\sum y^{2}}$$

where  $x = X - \overline{X}, y = Y - \overline{Y}$ 

If the deviations are taken from any arbitrary values of x and y (short – cut method)

$$\mathbf{b}_{1} = \mathbf{b}_{yx} = \frac{n \sum uv - \sum u \sum v}{n \sum u^{2} - \left(\sum u\right)^{2}}$$

$$\mathbf{b}_2 = \mathbf{b}_{xy} = \frac{n\sum uv - \sum u\sum v}{n\sum v^2 - (\sum v)^2}$$

where u = x - A: v = Y-BA = any value in X

B = any value in Y

# 9.5 Properties of Regression Co-efficient:

- Both regression coefficients must have the same sign, ie either they will be positive or negative.
- 2. correlation coefficient is the geometric mean of the regression coefficients ie,  $r = \pm \sqrt{b_1 b_2}$
- The correlation coefficient will have the same sign as that of the regression coefficients.
- If one regression coefficient is greater than unity, then other regression coefficient must be less than unity.
- Regression coefficients are independent of origin but not of scale.
- 6. Arithmetic mean of  $b_1$  and  $b_2$  is equal to or greater than the

coefficient of correlation. Symbolically

$$\frac{b_1 + b_2}{2} \ge r$$

Prepared by K. Aarthiya & R. Gayathri, Asst Prof, Department of Mathematics, KAHE

Page 28/37

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

- 7. If r=0, the variables are uncorrelated, the lines of regression become perpendicular to each other.
- 8. If  $r=\pm 1$ , the two lines of regression either coincide or parallel to each other
- 9. Angle between the two regression lines is  $\theta = \tan^{-1} \left| \frac{m_1 m_2}{1 + m_1 m_2} \right|$

where  $m_1$  and, $m_2$  are the slopes of the regression lines X on Y and Y on X respectively.

10. The angle between the regression lines indicates the degree of dependence between the variables.

# Example 2:

If 2 regression coefficients are 
$$b_1 = \frac{4}{5}$$
 and  $b_2 = \frac{9}{20}$ . What would be

the value of r?

The correlation coefficient , 
$$r = \pm \sqrt{b_1 b_2}$$
  
$$= \sqrt{\frac{4}{5} x \frac{9}{20}}$$
$$= \sqrt{\frac{36}{100}} = \frac{6}{10} = 0.6$$

# Example 3:

Given 
$$b_1 = \frac{15}{8}$$
 and  $b_2 = \frac{3}{5}$ , Find r  
Solution:  
 $r = \pm \sqrt{b_1 b_2}$   
 $= \sqrt{\frac{15}{8} \times \frac{3}{5}}$   
 $= \sqrt{\frac{9}{8}} = 1.06$ 

It is not possible since r, cannot be greater than one. So the given values are wrong

Page 29/37

CLASS: I B.Com(CA) COURSE NAME:BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

# Example 4:

Compute the two regression equations from the following data.

Х	1	2	3	4	5
Y	2	3	5	4	6
11 1	1	1 (			

If x = 2.5, what will be the value of y?

# Solution:

Х	Y	$x = X - \overline{X}$	$y = Y - \overline{Y}$	x <sup>2</sup>	y <sup>2</sup>	ху	1
1	2	-2	-2	4	4	4	
2	3	-1	-1	1	1	-1	]
3	5	0	1	0	1	0	]
4	4	1	0	1	0	0	
5	6	2	2	4	4	4	
15	20	20		10	10	9	


$$\overline{X} = \frac{\sum X}{n} = \frac{15}{5} = 3$$
$$\overline{Y} = \frac{\sum Y}{n} = \frac{20}{5} = 4$$

Regression Co efficient of Y on X

$$b_{yx} = \frac{\sum xy}{\sum x^2} = \frac{9}{10} = 0.9$$

Hence regression equation of Y on X is

$$Y = \overline{Y} + b_{yx}(X - \overline{X})$$
  
= 4 + 0.9 (X-3)  
= 4 + 0.9X - 2.7  
= 1.3 + 0.9X  
when X = 2.5  
Y = 1.3 + 0.9 × 2.5  
= 3.55



#### CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 BATCH-2017-2020 UNIT: IV

#### Example 6:

In a correlation study, the following values are obtained

	X	Y
Mean	65	67
S.D	2.5	3.5

Co-efficient of correlation = 0.8

Find the two regression equations that are associated with the above values.

#### Solution:

Given.

$$\overline{X} = 65, \ \overline{Y} = 67, \ \sigma_x = 2.5, \ \sigma_y = 3.5, \ r = 0.8$$

The regression co-efficient of Y on X is

$$b_{yx} = b_1 = r \frac{\sigma_y}{\sigma_x}$$
$$= 0.8 \times \frac{3.5}{2.5} = 1.12$$

The regression coefficient of X on Y is

$$b_{xy} = b_2 = r \frac{\sigma_x}{\sigma_y}$$
$$= 0.8 \times \frac{2.5}{3.5} = 0.57$$
Ience, the regression equation of Y on

Η X is

$$\begin{split} Y_e &= \overline{Y} + b_1 (X - \overline{X}) \\ &= 67 + 1.12 \text{ (X-65)} \\ &= 67 + 1.12 \text{ X} - 72.8 \\ &= 1.12 \text{ X} - 5.8 \end{split}$$

CLASS: I B.Com(CA)COURSE NAME: BUSINESS MATHEMATICS AND STATISTICSCOURSE CODE: 17CCU202UNIT: IVBATCH-2017-2020

The regression equation of X on Y is  $X_e = \overline{X} + b_2(Y - \overline{Y})$  = 65 + 0.57 (Y-67) = 65 + 0.57Y - 38.19= 26.81 + 0.57Y

# 9.7 Uses of Regression Analysis:

- Regression analysis helps in establishing a functional relationship between two or more variables.
- Since most of the problems of economic analysis are based on cause and effect relationships, the regression analysis is a highly valuable tool in economic and business research.
- Regression analysis predicts the values of dependent variables from the values of independent variables.
- 4. We can calculate coefficient of correlation (r) and coefficient of determination ( $r^2$ ) with the help of regression coefficients.
- In statistical analysis of demand curves, supply curves, production function, cost function, consumption function etc., regression analysis is widely used.

CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

S.No	Correlation	Regression
1.	Correlation is the relationship between two or more variables, which vary in sympathy with the other in the same or the opposite direction.	Regression means going back and it is a mathematical measure showing the average relationship between two variables
2.	Both the variables X and Y are random variables	Here X is a random variable and Y is a fixed variable. Sometimes both the variables may be random variables.
3.	It finds out the degree of relationship between two variables and not the cause and effect of the variables.	It indicates the causes and effect relationship between the variables and establishes functional relationship

	B.Com(CA) COURSE NAME:BUSINESS MAT DDE: 17CCU202 UNIT: IV	BATCH-2017-2020	
4.	It is used for testing and verifying the relation between two variables and gives limited information.	Besides verification it is used for the prediction of one value, in relationship to the other given value.	1
5.	The coefficient of correlation is a relative measure. The range of relationship lies between -1 and +1	Regression coefficient is an absolute figure. If we know the value of the independent variable, we can find the value of the dependent variable.	
6.	There may be spurious correlation between two variables.	In regression there is no such spurious regression.	
7.	It has limited application, because it is confined only to linear relationship between the variables.	It has wider application, as it studies linear and non- linear relationship between the variables.	
8.	It is not very useful for further mathematical treatment.	It is widely used for further mathematical treatment.	
9.	If the coefficient of correlation is positive, then the two variables are positively correlated and vice-versa.	The regression coefficient explains that the decrease in one variable is associated with the increase in the other variable.	

#### CLASS: I B.Com(CA) COURSE NAME: BUSINESS MATHEMATICS AND STATISTICS COURSE CODE: 17CCU202 UNIT: IV BATCH-2017-2020

#### **2 MARK QUESTIONS**

- 1. Define Correlation.
- 2. Define Standard Error and Probable Error.
- 3. Give any two points on merits and Demerits of Spearman's Rank correlation and Regression
- 4. Distinguish between correlation and Regression
- 5: Mentione therdifferent correlation rate transmission equations.
- 7. If r = 0.78,  $\sigma_x = 2.5$ ,  $\sigma_y = 3.5$  then find  $b_{xy}$ 8. write the formula for Rank correlation

#### **8 MARK QUESTIONS**

- 1. Distinguish between correlation and Regression (minimum 8 Points)
- 2. Marks obtained by 8 students in Accountancy (X) and Statistics (Y) are given

below. Compute Rank Correlation Coefficient.

below.	Compute	Rank Col	relation C	oemcient	•					
	Х	25	20	28	22	40	60		20	
	Y	40	30	50	30	20	10		30	
3. You	are given	the follow	ving data:							
				X			Y			
	thmetic m			3	6		85			
	ndard dev			-	.1		8			
Cor			between 2		0.66					
		•	ression eq	uations.						
	ii) Find 1	r.								
4. Find			fficient of					0.0	0.6	07
	X : 10		-		100	99	97	98	96	95
5 5	Y: 98			97	95	92	95	94	90	91
		· ·	ow find th			es.				
X: Y:	10	12 1 38 4		16	15					
		38   4 4 $100   4$		37	43.					
· · · · · · · · · · · · · · · · · · ·		hen $Y = 20$								
,			coefficien	t of correl	ation from	n the ma	rke secure	d by 1	0	
			nd statistic		ation non	i the ma	iks secure	AU UY I	0	
student			ancy (X):		65 30	90 4	0 50 7	5 85	60	
		1 statistics			70 40 9		60 80			
7. In a			he followi				00 00	00 00	•	
		X		Ŷ						
	Mean	65		67						
	S.D		.5	3.5						
	Coeffic	cient of co	rrelation =	: 0.8						
	Find th	e two regi	ression equ	lations.						
8. Find	l the Spea	rman's ra	nk correla	tion coeff	icient for t	the follo	wing data	•		
X: 6		64 7		64	80	75	40	55	64	
Y: 6	2	58 6	8 45	81	60	68	48	50	70	

	SE CODE: 17CCU202	UNIT: IV		TICS AND STATISTI BATCH-2017-202	
Maths:24153971068. Develop the Regression Equation that best fit the data given below using annual come as an independent variable and amount of life insurance as dependent variable.8Annual Income (Rs. in 000's)627841538534	Following are the rank obtainent of the rank obtainent.	d by 10 students in two	subjects. Find t	he Spearman's	
. Develop the Regression Equation that best fit the data given below using annualcome as an independent variable and amount of life insurance as dependent variable.Annual Income (Rs. in 000's)627841538534	_				
come as an independent variable and amount of life insurance as dependent variable.Annual Income (Rs. in 000's)627841538534					
Annual Income (Rs. in 000's)         62         78         41         53         85         34	Develop the Regression Equation	ation that best fit the da	ta given below u	sing annual	
					1
	Amount of I				