



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

**Subject: Mathematical Modeling**  
**Class: II M.Sc**

**Subject Code: 16MMP303**  
**Semester: III**

**L T P C**  
**4 0 0 4**

**PO:** After the completion of this course, the learner gain clear knowledge about various aspects of Mathematical modeling which is the motivating tool in the areas such as applied mathematics, engineering etc.

**PLO:** To understand the mathematical model of ODE of first order & second order, Population dynamics, genetics and to be familiar with mathematical models of graphs.

**UNIT I**

Mathematical Modeling through Ordinary Differential Equations of First order: Linear Growth and Decay Models – Non-Linear Growth and Decay Models – Compartment Models – Dynamics problems – Geometrical problems.

**UNIT II**

Mathematical Modeling through Systems of Ordinary Differential Equations of First Order: Population Dynamics – Epidemics – Compartment Models – Economics – Medicine, Arms Race, Battles and International Trade – Dynamics.

**UNIT III**

Mathematical Modeling through Ordinary Differential Equations of Second Order: Planetary Motions – Circular Motion and Motion of Satellites – Mathematical Modeling through Linear Differential Equations of Second Order – Miscellaneous Mathematical Models.

**UNIT IV**

Mathematical Modeling through Difference Equations : Simple Models – Basic Theory of Linear Difference Equations with Constant Coefficients – Economics and Finance – Population Dynamics and Genetics – Probability Theory.

**UNIT V**

Mathematical Modeling through Graphs: Solutions that can be Modeled through Graphs – Mathematical Modeling in Terms of Directed Graphs, Signed Graphs, Weighted Digraphs and Un oriented Graphs.

**SUGGESTED READINGS**

**TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modelling, Wiley Eastern Limited, New Delhi.

**REFERENCES**

**R1:** Kapur, J. N., (1985). Mathematical Models in Biology and Medicine, Affiliated East –West Press Pvt Limited, New Delhi.

**R2:** Brain Albright, (2010). Mathematical Modeling with Excel, Jones and Bartlett Publishers, New Delhi.

**R3:** Frank. R. Giordano, Maurice. D. Weir, William P. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

**Subject: Mathematical Modeling**  
**Class:II M.Sc**

**Subject Code: 16MMP303**  
**Semester:III**

**L T P C**  
**4 0 0 4**

S.No	Lecture Duration Hour	Topics To Be Covered	Support Materials
<b>Unit-I MATHEMATICAL MODELING THROUGH ORDINARY DIFFERENTIAL EQUATIONS OF FIRST ORDER</b>			
1	1	Mathematical modeling: Introduction and Simple Illustrations	T1:Chapter 1, Sec 1.1 Pg.No : 1-15
2	1	Mathematical modeling Through Differential Equations, Linear Growth and Decay Models	T1 : Chapter 2,Sec 2.1 Pg.No : 30-32
3	1	Continuation of types of Linear Growth and Decay Models	T1 : Chapter 2,Sec 2.2 Pg.No : 32-35
4	1	Non-Linear Growth and Decay Models	T1 : Chapter 2,Sec 2.3 Pg.No :35-37
5	1	Continuation of Non-Linear Growth and Decay Models	T1 : Chapter 2,Sec 2.3 Pg.No :37-39
6	1	Compartment Models	T1 : Chapter 2,Sec 2.4 Pg.No :39-41
7	1	Continuation of types of Compartment Models	T1 : Chapter 2,Sec 2.4 Pg.No :41-43
8	1	Mathematical modeling in Dynamics Through Ordinary Differential Equations of First Order	T1 : Chapter 2,Sec 2.5 Pg.No :43-45
9	1	Continuation of Mathematical modeling in Dynamics Through Ordinary Differential Equations of First Order	T1 : Chapter 2,Sec 2.5 Pg.No :45-48
10	1	Mathematical modeling of Geometrical Problems Through Ordinary Differential Equations of First Order	R3 : Chapter 2, Pg.No :75 - 77
11	1	Continuation of Mathematical modeling of Geometrical Problems Through Ordinary Differential Equations of First Order	R3 : Chapter 2, Pg.No :77 - 79

12	1	Recapitulation and discussion of possible Question	
<b>Total</b>	<b>12 Hrs</b>		
<b>TEXT BOOK:</b> <b>T1:</b> J.N.Kapur,2015. Mathematical modeling,Wiley Eastern Limited,New Delhi. <b>REFERENCES:</b> <b>R3:</b> Frank.R.Giordano,Maurice.D.Weir, William P. Fox,2003,A first course in Mathematical modeling , Vikash Publishing House, UK.			
<b>Unit-II MATHEMATICAL MODELING THROUGH SYSTEMS OF ORDINARY DIFFERENTIAL EQUATIONS OF THE FIRST ORDER</b>			
1	1	Mathematical modeling in Population Dynamics	T1 : Chapter 3,Sec 3.1 Pg.No :53-60
2	1	Mathematical modeling of Epidemics Through Systems of Ordinary Differential Equations of First Order	R2 : Chapter 4, Pg.No :118-120
3	1	Continuation of Mathematical modeling of Epidemics Through Systems of Ordinary Differential Equations of First Order	R2 : Chapter 4, Pg.No :120-122
4	1	Continuation of Mathematical modeling of Epidemics Through Systems of Ordinary Differential Equations of First Order	R2 : Chapter 4, Pg.No :122-124
5	1	Compartment Models Through Systems of Ordinary Differential Equations	T1 : Chapter 3,Sec 3.3 Pg.No :63-64
6	1	Mathematical modeling in Economics Through Systems of Ordinary Differential Equations of First Order	T1 : Chapter 3,Sec 3.4 Pg.No :64 - 66
7	1	Continuation of types of Mathematical modeling in Economics Through Systems of Ordinary Differential Equations of First Order	T1 : Chapter 3,Sec 3.4 Pg.No :66 - 69
8	1	Mathematical Models in Medicine, Arms Race, Battles and International Trade in Terms of Systems of Ordinary Differential Equations	R3 : Chapter 9, Pg.No :350 - 355
9	1	Continuation of types of Mathematical Models in Medicine, Arms Race, Battles and International Trade in Terms of Systems of Ordinary Differential Equations	T1 : Chapter 3,Sec 3.5 Pg.No :69 - 72
10	1	Mathematical modeling in Dynamics through systems of Ordinary Differential Equations of First Order	T1 : Chapter 3,Sec 3.6 Pg.No :72-74
11	1	Continuation of Mathematical modeling in	T1 : Chapter 3,Sec 3.6

		Dynamics through systems of Ordinary Differential Equations of First Order	Pg.No :74-76
12	1	Recapitulation and discussion of possible Question	
<b>Total</b>	<b>12Hrs</b>		
<b>TEXT BOOK:</b> <b>T1:</b> J.N.Kapur,2015. Mathematical modeling,Wiley Eastern Limited,New Delhi. <b>REFERENCES:</b> <b>R2:</b> Brain Albright,2012. Mathematical modeling with Excel, Jones and Bartlett Publishers,New Delhi. <b>R3:</b> Frank.R.Giordano,Maurice.D.Weir, William P. Fox,2003,A first course in Mathematical modeling , Vikash Publishing House, UK.			
<b>Unit-III MATHEMATICAL MODELING THROUGH ORDINARY DIFFERENTIAL EQUATIONS OF SECOND ORDER</b>			
1	1	Mathematical modeling of Planetary Motions	T1 : Chapter 4,Sec 4.1 Pg.No :76-79
2	1	Continuation of types of Mathematical modeling of Planetary Motions	T1 : Chapter 4,Sec 4.1 Pg.No :79-82
3	1	Mathematical modeling of Circular Motion and Motion of Satellites	T1 : Chapter 4,Sec 4.2 Pg.No :82-85
4	1	Continuation of types of Mathematical modeling of Circular Motion and Motion of Satellites	T1 : Chapter 4,Sec 4.2 Pg.No :85-88
5	1	Mathematical modeling Through Linear Differential Equations of Second Order	T1 : Chapter 4,Sec 4.3 Pg.No :88-90
6	1	Continuation of types of Mathematical modeling Through Linear Differential Equations of Second Order	T1 : Chapter 4,Sec 4.3 Pg.No :90-93
7	1	Continuation of Problems on Mathematical modeling Through Linear Differential Equations of Second Order	R2 : Chapter 7, Pg.No :238-244
8	1	Miscellaneous Mathematical Model Through Ordinary Differential Equations of the Second Order	T1 : Chapter 4,Sec 4.4 Pg.No :93-96
9	1	Continuation of Problems on Miscellaneous Mathematical Model Through Ordinary Differential Equations of the Second Order	R3: Chapter 11, Pg.No : 437-440
10	1	Continuation of Problems on Miscellaneous Mathematical Model Through Ordinary Differential Equations of the Second Order	R3: Chapter 11, Pg.No : 440-445
11	1	Continuation of Problems on Miscellaneous	R3: Chapter 11, Pg.No :



		Mathematical Model Through Ordinary Differential Equations of the Second Order	445-452
12	1	Recapitulation and discussion of possible Question	
<b>Total</b>	<b>12 Hrs</b>		
<b>TEXT BOOK:</b> <b>T1:</b> J.N.Kapur,2015. Mathematical modeling,Wiley Eastern Limited,New Delhi. <b>REFERENCES:</b> <b>R2:</b> Brain Albright,2012. Mathematical modeling with Excel, Jones and Bartlett Publishers,New Delhi. <b>R3:</b> Frank.R.Giordano,Maurice.D.Weir, William P. Fox,2003,A first course in Mathematical modeling , Vikash Publishing House, UK.			
<b>Unit-IV MATHEMATICAL MODELING THROUGH DIFFERENCE EQUATIONS</b>			
1	1	The Need for Mathematical modeling Through Difference Equations : Some simple Models	T1 : Chapter 5,Sec 5.1 Pg.No :96-98
2	1	Basic Theory of Linear Difference Equations with Constant Coefficients	T1 : Chapter 5,Sec 5.2 Pg.No :98-101
3	1	Continuation of types of Basic Theory of Linear Difference Equations with Constant Coefficients	T1 : Chapter 5,Sec 5.2 Pg.No :101-105
4	1	Mathematical modeling Through Difference Equations in Economics and Finance	T1 : Chapter 5,Sec 5.3 Pg.No :105-107
5	1	Continuation of Mathematical modeling Through Difference Equations in Economics and Finance	T1 : Chapter 5,Sec 5.3 Pg.No :107-110
6	1	Mathematical modeling Through Difference Equations in Population Dynamics and Genetics	T1 : Chapter 5,Sec 5.4 Pg.No :110 - 113
7	1	Continuation of types of Mathematical modeling Through Difference Equations in Population Dynamics and Genetics	T1 : Chapter 5,Sec 5.4 Pg.No :113 - 117
8	1	Mathematical modeling Through Difference Equations in Probability Theory	R3: Chapter 6, Pg.No :217-220
9	1	Continuation of Mathematical modeling Through Difference Equations in Probability Theory	R3: Chapter 6, Pg.No :220-223
10	1	Miscellaneous Examples of Mathematical modeling Through Difference Equations	T1 : Chapter 5,Sec 5.6 Pg.No :121-122
11	1	Continuation of Miscellaneous Examples of Mathematical modeling Through Difference Equations	T1 : Chapter 5,Sec 5.6 Pg.No :122-124

12	1	Recapitulation and discussion of possible Question	
<b>Total</b>	<b>12 Hrs</b>		
<b>TEXT BOOK:</b> <b>T1:</b> J.N.Kapur,2015. Mathematical modeling,Wiley Eastern Limited,New Delhi. <b>REFERENCES:</b> <b>R3:</b> Frank.R.Giordano,Maurice.D.Weir, William P. Fox,2003,A first course in Mathematical modeling , Vikash Publishing House, UK.			
<b>Unit-V MATHEMATICAL MODELING THROUGH GRAPHS</b>			
1	1	Situations that can be Modelled through Graphs	T1 : Chapter 7,Sec 7.1 Pg.No :151-154
2	1	Mathematical Models in terms of Directed Graphs	T1 : Chapter 7,Sec 7.2 Pg.No :154-156
3	1	Continuation of types of Mathematical Models in terms of Directed Graphs	T1 : Chapter 7,Sec 7.2 Pg.No :156-161
4	1	Mathematical Models in terms of Signed Graphs	R3 : Chapter 3, Pg.No : 101-107
5	1	Mathematical modeling in terms of Weighted Digraphs	T1 : Chapter 7,Sec 7.4 Pg.No :164-170
6	1	Continuation of Mathematical modeling in terms of Weighted Digraphs	T1 : Chapter 7,Sec 7.4 Pg.No :164-170
7	1	Mathematical modeling in terms of Unoriented Graphs	T1 : Chapter 7,Sec 7.5 Pg.No :170-173
8	1	Continuation of Mathematical modeling in terms of Unoriented Graphs	T1 : Chapter 7,Sec 7.5 Pg.No :173-177
9	1	Recapitulation and discussion of possible Question	
10	1	Discussion of previous ESE question papers	
11	1	Discussion of previous ESE question papers	
12	1	Discussion of previous ESE question papers	
<b>Total</b>	<b>12Hrs</b>		
<b>TEXT BOOK:</b> <b>T1:</b> J.N.Kapur,2015. Mathematical modeling,Wiley Eastern Limited,New Delhi. <b>REFERENCES:</b> <b>R3:</b> Frank.R.Giordano,Maurice.D.Weir, William P. Fox,2003,A first course in Mathematical modeling , Vikash Publishing House, UK.			

**Total no.of hours for the course:60 Hours**

**TEXT BOOK:**

**T1:** J.N.Kapur,2015. Mathematical modeling,Wiley Eastern Limited,New Delhi.

**REFERENCES:**

**R1:** J.N.Kapur,1981. Mathematical Models in Biology and Medicine Affiliated East-West Press Pvt Limited,New Delhi.

**R2:**Brain Albright,2012. Mathematical modeling with Excel, Jones and Bartlett Publishers,New Delhi.

**R3:** Frank.R.Giordano,Maurice.D.Weir, William P. Fox,2003,A first course in Mathematical modeling , Vikash Publishing House, UK.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
(Deemed to be University Established Under Section 3 of UGC Act 1956)  
Pollachi Main Road, Eachanari (Po),  
Coimbatore –641 021  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

---

## **UNIT I**

Mathematical Modeling through Ordinary Differential Equations of First order: Linear Growth and Decay Models – Non-Linear Growth and Decay Models – Compartment Models – Dynamics problems – Geometrical problems.

### **SUGGESTED READINGS**

#### **TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

#### **REFERENCES**

**R1:**Kapur, J. N., (1985). Mathematical Models in Biology and Medicine, Affiliated East –West Press Pvt Limited, New Delhi.

**R3:**Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.

## MATHEMATICAL MODELLING THROUGH DIFFERENTIAL EQUATIONS

Mathematical Modelling in terms of differential equations arises when the situation modelled involves some *continuous* variable(s) varying with respect to some other continuous variable(s) and we have some reasonable hypotheses about the *rates of change* of dependent variable(s) with respect to independent variable(s).

When we have one dependent variable  $x$  (say population size) depending on one independent variable (say time  $t$ ), we get a mathematical model in terms of an *ordinary differential equation* of the *first order*, if the hypothesis is about the rate of change  $dx/dt$ . The model will be in terms of an *ordinary differential equation of the second order* if the hypothesis involves the rate of change of  $dx/dt$ .

If there are a number of dependent continuous variables and only one independent variable, the hypothesis may give a mathematical model in terms of a *system of first or higher order ordinary differential equations*.

If there is one dependent continuous variable (say velocity of fluid  $u$ ) and a number of independent continuous variables (say space coordinates  $x, y, z$  and time  $t$ ), we get a mathematical model in terms of a *partial differential equation*. If there are a number of dependent continuous variables and a number of independent continuous variables, we can get a mathematical model in terms of systems of *partial differential equations*.

## LINEAR GROWTH AND DECAY MODELS

### Populational Growth Models

Let  $x(t)$  be the population size at time  $t$  and let  $b$  and  $d$  be the birth and death rates, i.e. the number of individuals born or dying per individual

per unit time, then in time interval  $(t, t + \Delta t)$ , the numbers of births and deaths would be  $bx \Delta t + o(\Delta t)$  and  $dx \Delta t + o(\Delta t)$  where  $o(\Delta t)$  is an infinitesimal which approaches zero as  $\Delta t$  approaches zero, so that

$$\Delta x = x(t + \Delta t) - x(t) = (bx(t) - dx(t))\Delta t + o(\Delta t), \quad (1)$$

so that dividing by  $\Delta t$  and proceeding to the limit as  $\Delta t \rightarrow 0$ , we get



$$\frac{dx}{dt} = (b - d)x = ax \quad (\text{say}) \quad (2)$$

Integrating (2), we get

$$x(t) = x(0) \exp(at), \quad (3)$$

so that the population grows exponentially if  $a > 0$ , decays exponentially if  $a < 0$  and remains constant if  $a = 0$  (Figure 2.1)




Figure 2.1

(i) If  $a > 0$ , the population will become double its present size at time  $T$ , where

$$2x(0) = x(0) \exp(aT) \quad \text{or} \quad \exp(aT) = 2$$

$$\text{or} \quad T = \frac{1}{a} \ln 2 = (0.69314118)a^{-1} \quad (4)$$

$T$  is called the doubling period of the population and it may be noted that this doubling period is independent of  $x(0)$ . It depends only on  $a$  and is such that greater the value of  $a$  (i.e. greater the difference between birth and death rates), the smaller is the doubling period.

(ii) If  $a < 0$ , the population will become half its present size in time  $T'$  when

$$\frac{1}{2}x(0) = x(0) \exp(aT') \quad \text{or} \quad \exp(aT') = \frac{1}{2}$$

$$\text{or} \quad T' = \frac{1}{a} \ln \frac{1}{2} = -(0.69314118)a^{-1} \quad (5)$$

It may be noted that  $T'$  is also independent of  $x(0)$  and since  $a < 0$ ,  $T' > 0$ .  $T'$  may be called the half-life (period) of the population and it decreases as the excess of death rate over birth rate increases.

## Growth of Science and Scientists

Let  $S(t)$  denote the number of scientists at time  $t$ ,  $bS(t)\Delta t + u(\Delta t)$  be the number of new scientists trained in time interval  $(t, t + \Delta t)$  and let



$dS(t)\Delta t + 0(\Delta t)$  be the number of scientists who retire from science in the same period, then the above model applies and the number of scientists should grow exponentially.

The same model applies to the growth of Science, Mathematics and Technology. Thus if  $M(t)$  is the amount of Mathematics at time  $t$ , then the rate of growth of Mathematics is proportional to the amount of Mathematics, so that

$$dM/dt = aM \quad \text{or} \quad M(t) = M(0) \exp(at) \quad (6)$$

Thus according to this model, Mathematics, Science and Technology grow at an exponential rate and double themselves in a certain period of time. During the last two centuries this doubling period has been about ten years. This implies that if in 1900, we had one unit of Mathematics, then in 1910, 1920, 1930, 1940, ... 1980 we have 2, 4, 8, 16, 32, 64, 128, 256 unit of Mathematics and in 2000 AD we shall have about 1000 units of Mathematics. This implies that 99.9% of Mathematics that would exist at the end of the present century would have been created in this century and 99.9% of all mathematicians who ever lived, would have lived in this century.

The doubling period of mathematics is 10 years and the doubling period of the human population is 30-35 years. These doubling periods cannot obviously be maintained indefinitely because then at some point of time, we shall have more mathematicians than human beings. Ultimately the doubling period of both will be the same, but hopefully this is a long way away.

This model also shows that the doubling period can be shortened by having more intensive training programmes for mathematicians and scientists and by creating conditions in which they continue to do creative work for longer durations in life.

## Effects of Immigration and Emigration on Population Size

If there is immigration into the population from outside at a rate proportional to the population size, the effect is equivalent to increasing the birth rate. Similarly if there is emigration from the population at a rate proportional to the population size, the effect is the same as that of increase in the death rate.

If however immigration and emigration take place at constant rate  $i$  and  $e$  respectively, equation (3) is modified to



$$\frac{dx}{dt} = bx - dx + i - e = ax + k \quad (7)$$

Integrating (7) we get

$$x(t) + \frac{k}{a} = \left(x(0) + \frac{k}{a}\right)e^{at} \quad (8)$$

The model also applies to growth of populations of bacteria and micro-organisms, to the increase of volume of timber in forest, to the growth of malignant cells etc. In the case of forests, planting of new plants will correspond to immigration and cutting of trees will correspond to emigration.

## Radio-Active Decay

Many substances undergo radio-active decay at a rate proportional to the amount of the radioactive substance present at any time and each of them has a half-life period. For uranium 238 it is 4.55 billion years. For potassium it is 1.3 billion years. For thorium it is 13.9 billion years. For rubidium it is 50 billion years while for carbon 14, it is only 5568 years and for white lead it is only 22 years.

In radiogeology, these results are used for radioactive dating. Thus the ratio of radio-carbon to ordinary carbon (carbon 12) in dead plants and animals enables us to estimate their time of death. Radioactive dating has also been used to estimate the age of the solar system and of earth as 45 billion years.

### 2.2.7 Diffusion

According to Fick's law of diffusion, the time rate of movement of a solute across a thin membrane is proportional of the area of the membrane and to the difference in concentrations of the solute on the two sides of the membrane.

If the area of the membrane is constant and the concentration of solute on one side is kept fixed at  $a$  and the concentration of the solution on the other side initially is  $c_0 < a$ , then Fick's law gives

$$\frac{dc}{dt} = k(a - c), \quad c(0) = c_0, \quad (15)$$

so that

$$a - c(t) = (a - c(0))e^{-kt} \quad (16)$$

and  $c(t) \rightarrow a$  as  $t \rightarrow \infty$ , whatever be the value of  $c_0$ .

## Change of Price of a Commodity

Let  $p(t)$  be the price of a commodity at time  $t$ , then its rate of change is proportional to the difference between the demand  $d(t)$  and the supply  $s(t)$  of the commodity in the market so that

$$\frac{dp}{dt} = k(d(t) - s(t)), \quad (17)$$

where  $k > 0$ , since if demand is more than the supply, the price increases. If  $d(t)$  and  $s(t)$  are assumed linear functions of  $p(t)$ , i.e. if

$$d(t) = d_1 + d_2 p(t), \quad s(t) = s_1 + s_2 p(t), \quad d_2 < 0, s_2 > 0 \quad (18)$$

we get

$$\frac{dp}{dt} = k(d_1 - s_1 + (d_2 - s_2)p(t)) = k(a - \beta p(t)), \quad \beta > 0 \quad (19)$$

or

$$\frac{dp}{dt} = K(p_e - p(t)), \quad (20)$$

where  $p_e$  is the equilibrium price, so that

$$p_e - p(t) = (p_e - p(0))e^{-kt} \quad (21)$$

and

$$p(t) \rightarrow p_e \quad \text{as} \quad t \rightarrow \infty$$



**Logistic Law of Population Growth**

(i)  $x(0) < a/b \Rightarrow x(t) < a/b \Rightarrow dx/dt > 0 \Rightarrow x(t)$  is a monotonic increasing function of  $t$  which approaches  $a/b$  as  $t \rightarrow \infty$ .

(ii)  $x(0) > a/b \Rightarrow x(t) > a/b \Rightarrow dx/dt < 0 \Rightarrow x(t)$  is a monotonic decreasing function of  $t$  which approaches  $a/b$  as  $t \rightarrow \infty$ .

Now from (23)

$$\frac{d^2x}{dt^2} = a - 2bx, \quad (25)$$

so that  $d^2x/dt^2 \gtrless 0$  according as  $x \gtrless a/2b$ . Thus in case (i) the growth curve is convex if  $x < a/2b$  and is concave if  $x > a/2b$  and it has a point of inflexion at  $x = a/2b$ . Thus the graph of  $x(t)$  against  $t$  is as given in Figure 2.2.




Figure 2.2

- If  $x(0) < a/2b$ ,  $x(t)$  increases at an increasing rate till  $x(t)$  reaches  $a/2b$  and then it increases at a decreasing rate and approaches  $a/b$  as  $t \rightarrow \infty$
- If  $a/2b < x(0) < a/b$ ,  $x(t)$  increases at a decreasing rate and approaches  $a/b$  as  $t \rightarrow \infty$
- If  $x(0) = a/b$ ,  $x(t)$  is always equal to  $a/b$
- If  $x(0) > a/b$ ,  $x(t)$  decreases at a decreasing absolute rate and approaches  $a/b$  as  $t \rightarrow \infty$

**Spread of Technological Innovations and Infectious Diseases**

Let  $N(t)$  be the number of companies which have adopted a technological innovation till time  $t$ , then the rate of change of the number of these companies depends both on the number of companies which have adopted this innovation and on the number of those which have not yet adopted it, so that if  $R$  is the total number of companies in the region

$$\frac{dN}{dt} = kN(R - N), \quad (26)$$

which is the logistic law and shows that ultimately all companies will adopt this innovation.

Similarly if  $N(t)$  is the number of infected persons, the rate at which the number of infected persons increases depends on the product of the numbers of infected and susceptible persons. As such we again get (26), where  $R$  is the total number of persons in the system.

It may be noted that in both the examples, while  $N(t)$  is essentially an integer-valued variable, we have treated it as a continuous variable. This can be regarded as an idealisation of the situation or as an approximation to reality.

### Rate of Dissolution

Let  $x(t)$  be the amount of undissolved solute in a solvent at time  $t$  and let  $c_0$  be the maximum concentration or saturation concentration, i.e. the maximum amount of the solute that can be dissolved in a unit volume of the solvent. Let  $V$  be the volume of the solvent. It is found that the rate at which the solute is dissolved is proportional to the amount of undissolved solute and to the difference between the concentration of the solute at time  $t$  and the maximum possible concentration, so that we get

$$\frac{dx}{dt} = kx(t) \left( \frac{x(0) - x(t)}{V} - c_0 \right) = \frac{kx(t)}{V} ((x_0 - c_0V) - x(t)) \quad (27)$$



## Law of Mass Action: Chemical Reactions

Two chemical substances combine in the ratio  $a : b$  to form a third substance  $Z$ . If  $z(t)$  is the amount of the third substance at time  $t$ , then a proportion  $az(t)/(a + b)$  of it consists of the first substance and a proportion  $bz(t)/(a + b)$  of it consists of the second substance. The rate of formation of the third substance is proportional to the product of the amount of the two component substances which have not yet combined together. If  $A$  and  $B$  are the initial amounts of the two substances, then we get

$$\frac{dz}{dt} = k \left( A - \frac{az}{a+b} \right) \left( B - \frac{bz}{a+b} \right) \quad (28)$$

This is the non-linear differential equation for a second order reaction. Similarly for an  $n$ th order reaction, we get the non-linear equation

$$\frac{dz}{dt} = k(A_1 - a_1z)(A_2 - a_2z) \dots (A_n - a_nz), \quad (29)$$

where  $a_1 + a_2 + \dots + a_n = 1$ .

## EXERCISE

1. If in (24),  $a = 0.03134$ ,  $b = (1.5887)(10)^{-10}$ ,  $x(0) = 39 \times 10^6$ , show that

$$x(t) = \frac{313,400,000}{1.5887 + 78,7703^{-0.03134t}}$$

This is Verhulst model for the population of USA when time zero corresponds to 1790. Estimate the population of USA in 1800, 1850, 1900 and 1950. Show that the point of inflexion should have occurred in about 1914. Find also the limiting population of USA on the basis of this model.

2. In (26)  $k = 0.007$ ,  $R = 1000$ ,  $N(0) = 50$ , find  $N(10)$  and find when  $N(t) = 500$ .

## COMPARTMENT MODELS

In the last two sections, we got mathematical models in terms of ordinary differential equations of the first order, in all of which variables were separable. In the present section, we get models in terms of linear differential equations of first order.

We also use here the principle of continuity i.e. that the gain in amount of a substance in a medium in any time is equal to the excess of the amount that has entered the medium in the time over the amount that has left the medium in this time.

### A Simple Compartment Model

Let a vessel contain a volume  $V$  of a solution with concentration  $c(t)$  of a substance at time  $t$  (Figure 2.3). Let a solution with constant concentration  $C$  in an overhead tank enter the vessel at a constant rate  $R$  and after mixing thoroughly with the solution in the vessel, let the mixture with concentration  $c(t)$  leave the vessel at the same rate  $R$  so that the volume of the solution in the vessel remains  $V$ .




Figure 2.3

Using the principle of continuity, we get

$$V(c(t + \Delta t) - c(t)) = RC \Delta t - Rc(t)\Delta t + o(\Delta t)$$

giving

$$V \frac{dc}{dt} + Rc = RC \quad (30)$$

Integrating

$$c(t) = c(0) \exp\left(-\frac{R}{V}t\right) + C\left(1 - \exp\left(-\frac{R}{V}t\right)\right) \quad (31)$$

As  $t \rightarrow \infty$ ,  $c(t) \rightarrow C$ , so that ultimately the vessel has the same concentration as the overhead tank. Since

$$c(t) = C - (C - c_0) \exp\left(-\frac{R}{V}t\right), \quad (32)$$



if  $C > c_0$ , the concentration in the vessel increases to  $C$ ; on the other hand if  $C < c_0$ , the concentration in the vessel decreases to  $C$  (Figure 2.4).

If the rate  $R'$  at which the solution leaves the vessel is less than  $R$ , the equations of continuity gives

$$\frac{d}{dt}[(V_0 + (R - R')t)c(t)] = RC - R'(ct) \quad (33)$$

where  $V$  is the initial volume of the solution in the vessel. This is also a linear differential equation of the first order.




Figure 2.4

### Diffusion of Glucose or a Medicine in the Blood Stream

Let the volume of blood in the human body be  $V$  and let the initial concentration of glucose in the blood stream be  $c(0)$ . Let glucose be introduced in the blood stream at a constant rate  $I$ . Glucose is also removed from the blood stream due to the physiological needs of the human body at a rate proportional to  $c(t)$ , so that the continuity principle gives

$$V \frac{dc}{dt} = I - kc \quad (34)$$

which is similar to (30).

Now let a dose  $D$  of a medicine be given to a patient at regular intervals of duration  $T$  each. The medicine also disappears from the system at a rate proportional to  $c(t)$ , the concentration of the medicine in the blood stream, then the differential equation given by the continuity principle is

$$V \frac{dc}{dt} = -kc \quad (35)$$

Integrating

$$c(t) = D \exp\left(-\frac{k}{V}t\right), \quad 0 \leq t < T \quad (36)$$

At time  $T$ , the residue of the first dose is  $D \exp\left(-\frac{k}{V}T\right)$  and now another dose  $D$  is given so that we get



$$c(t) = \left( D \exp\left(-\frac{k}{V} T\right) + D \right) \exp\left(-\frac{k}{V}(t - T)\right), \quad (37)$$

$$= D \exp\left(-\frac{k}{V} t\right) + D \exp\left(-\frac{k}{V}(t - T)\right), \quad (38)$$

$$T \leq t < 2T$$

The first term gives the residual of the first dose and the second term gives the residual of the second dose. Proceeding in the same way, we get after  $n$  doses have been given

$$c(t) = D \exp\left(-\frac{k}{V} t\right) + D \exp\left(-\frac{k}{V}(t - T)\right) \\ + D \exp\left(-\frac{k}{V}(t - 2T)\right) + \dots + D \exp\left(-\frac{k}{V}(t - (n-1)T)\right) \quad (39)$$

$$= D \exp\left(-\frac{k}{V} t\right) \left( 1 + \exp\left(\frac{k}{V} T\right) + \exp\left(\frac{2k}{V} T\right) \right. \\ \left. + \dots + \exp\left((n-1) \frac{k}{V} T\right) \right) \\ = D \exp\left(-\frac{k}{V} t\right) \frac{\exp\left(n \frac{k}{V} T\right) - 1}{\exp\left(\frac{k}{V} T\right) - 1}, \quad (n-1)T \leq t < nT \quad (40)$$

$$c(nT - 0) = D \frac{1 - \exp\left(-\frac{k}{V} nT\right)}{\exp\left(\frac{kT}{V}\right) - 1} \quad (41)$$

$$c(nT + 0) = D \frac{\exp\left(\frac{kT}{V}\right) - \exp\left(-\frac{k}{V} nT\right)}{\exp\left(\frac{kT}{V}\right) - 1} \quad (42)$$

Thus the concentration never exceeds  $D/(1 - \exp(-\frac{kT}{V}))$ . The graph of  $c(t)$  is shown in Figure 2.5.




Figure 2.5

Thus in each interval, concentration decreases. In any interval, the concentration is maximum at the beginning of this interval and thus maximum concentration at the beginning of an interval goes on increasing as the number of intervals increases, but the maximum value is always below  $D/(1 - e^{-kT/V})$ . The minimum value in an interval occurs at the end of each interval. This also increases, but it lies below  $D/(\exp(kT/V) - 1)$ .

The concentration curve is piecewise continuous and has points of discontinuity at  $T, 2T, 3T, \dots$

By injecting glucose or penicillin in blood and fitting curve (36) to the data, we can estimate the value of  $k$  and  $V$ . In particular this gives a method for finding the volume of blood in the human body.

### 2.4.3 The Case of a Succession of Compartments

Let a solution with concentration  $c(t)$  of a solute pass successively into  $n$  tanks in which the initial concentrations of the solution are  $c_1(0), c_2(0), \dots, c_n(0)$ . The rates of inflow in each tank is the same as the rate of outflow from the tank. We have to find the concentrations  $c_1(t), c_2(t) \dots c_n(t)$  at time  $t$ . We get the equations

$$\begin{aligned} V \frac{dc_1}{dt} &= Rc - Rc_1 \\ V \frac{dc_2}{dt} &= Rc_1 - Rc_2 \\ &\dots \dots \dots \\ V \frac{dc_n}{dt} &= Rc_{n-1} - Rc_n \end{aligned} \quad (43)$$

By solving the first of these equations, we get  $c_1(t)$ . Substituting the value of  $c_1(t)$  and proceeding in the same way, we can find  $c_2(t), \dots, c_n(t)$ .

## MATHEMATICAL MODELLING IN DYNAMICS THROUGH ORDINARY DIFFERENTIAL EQUATIONS OF FIRST ORDER

Let a particle travel a distance  $x$  in time  $t$  in a straight line, then its velocity  $v$  is given by  $dx/dt$  and its acceleration is given by

$$dv/dt = (dv/dx)(dx/dt) = v dv/dx = d^2x/dt^2$$



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

---

**UNIT I(continuation)**

Mathematical Modeling through Ordinary Differential Equations of First order: Linear Growth and Decay Models – Non-Linear Growth and Decay Models – Compartment Models – Dynamics problems – Geometrical problems.

**SUGGESTED READINGS**

**TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

**REFERENCES**

**R1:**Kapur, J. N., (1985). Mathematical Models in Biology and Medicine, Affiliated East –West Press Pvt Limited, New Delhi.

**R3:**Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.



## Simple Harmonic Motion

Here a particle moves in a straight line in such a manner that its acceleration is always proportional to its distance from the origin and is always directed towards the origin, so that

$$v \frac{dv}{dx} = -\mu x \quad (44)$$

Integrating

$$v^2 = \mu(a^2 - x^2), \quad (45)$$

where the particle is initially at rest at  $x = a$ . Equation (44) gives

$$\frac{dx}{dt} = -\sqrt{\mu} \sqrt{a^2 - x^2} \quad (46)$$

We take the negative sign since velocity increases as  $x$  decreases (Figure 2.6).




Figure 2.6

Integrating again and using the condition that at  $t = 0$ ,  $x = a$

$$x(t) = a \cos \sqrt{\mu} t \quad (47)$$

so that

$$v(t) = -a\sqrt{\mu} \sin \sqrt{\mu} t, \quad (48)$$

Thus in simple harmonic motion, both displacement and velocity are periodic functions with period  $2\pi/\sqrt{\mu}$ .

The particle starts from  $A$  with zero velocity and moves towards  $0$  with increasing velocity and reaches  $0$  at time  $\pi/2\sqrt{\mu}$  with velocity  $\sqrt{\mu}a$ . It continues to move in the same direction, but now with decreasing velocity till it reaches  $A'$  ( $OA' = a$ ) where its velocity is again zero. It then begins moving towards  $0$  with increasing velocity and reaches  $0$  with velocity  $\sqrt{\mu}a$  and again comes to rest at  $A$  after a total time period  $2\pi/\sqrt{\mu}$ . The periodic motion then repeats itself.

As one example of SHM, consider a particle of mass  $m$  attached to one end of a perfectly elastic string, the other end of which is attached to a fixed point  $O$  (Figure 2.7). The particle moves under gravity in vacuum.

Let  $l_0$  be the natural length of the string and let  $a$  be its extension when the particle is in equilibrium so that by Hooke's law

$$mg = T_0 = \lambda \frac{a}{l_0} \quad (49)$$

where  $\lambda$  is the coefficient of elasticity. Now let the string be further stretched a distance  $x$  and then the mass be left free. The equation of motion which states that mass  $\times$  acceleration in any direction = force




Figure 2.7

on the particle in that direction, gives

$$mv \frac{dv}{dx} = mg - T = mg - \lambda \frac{a+x}{l_0} = -\frac{\lambda x}{l_0} \quad (50)$$

or

$$v \frac{dv}{dx} = \frac{\lambda}{m} \frac{x}{l_0} = -\frac{gx}{a}, \quad (51)$$

which gives a simple harmonic motion with time period  $2\pi \sqrt{\frac{a}{g}}$ .

### Motion Under Gravity in a Resisting Medium

A particle falls under gravity in a medium in which the resistance is proportional to the velocity. The equation of motion is

$$m \frac{dv}{dt} = mg - mkv$$

or

$$\frac{dv}{V-v} = k dt; \quad V = \frac{g}{k} \quad (52)$$



Integrating

$$V - v = Ve^{-kt} \quad (53)$$

If the particle starts from rest with zero velocity. Equation (50) gives

$$v = V(1 - e^{-kt}), \quad (54)$$

so that the velocity goes on increasing and approaches the limiting velocity  $g/k$  as  $t \rightarrow \infty$ . Replacing  $v$  by  $dx/dt$ , we get

$$\frac{dx}{dt} = V(1 - e^{-kt}) \quad (55)$$

Integrating and using  $x = 0$  when  $t = 0$ , we get

$$x = Vt + \frac{Ve^{-kt}}{k} - \frac{V}{k}. \quad (56)$$

## Motion of a Rocket

As a first idealisation, we neglect both gravity and air resistance. A rocket moves forward because of the large supersonic velocity with which gases produced by the burning of the fuel inside the rocket come out of the converging-diverging nozzle of the rocket (Figure 2.8).

Let  $m(t)$  be the mass of the rocket at time  $t$  and let it move forward with velocity  $v(t)$  so that the momentum at time  $t$  is  $m(t)v(t)$ .

In the interval of time  $(t, t + \Delta t)$ , the mass of the rocket becomes

$$m(t + \Delta t) = m(t) + \frac{dm}{dt} \Delta t + o(\Delta t)$$




Figure 2.8

Since the rocket is losing mass,  $dm/dt$  is negative and the mass of gases  $-dm/dt \Delta t$  moves with velocity  $u$  relative to the rocket, i.e. with a velocity  $v(t + \Delta t) - u$  relative to the earth so that the total momentum of the rocket and the gases at time  $t + \Delta t$  is

$$m(t + \Delta t)v(t + \Delta t) - \frac{dm}{dt}\Delta t(v(t + \Delta t) - u) \quad (57)$$

Since we are neglecting air resistance and gravity, there is no external force on the rocket and as such the momentum is conserved, giving the equation

$$\begin{aligned} m(t)v(t) &= \left(m(t) + \frac{dm}{dt}\Delta t\right) \left(v(t) + \frac{dv}{dt}\Delta t\right) \\ &\quad - \frac{dm}{dt}\Delta t(v - u) + 0(\Delta t)^2 \end{aligned} \quad (58)$$

Dividing by  $\Delta t$  and proceeding to the limit as  $\Delta t \rightarrow 0$ , we get

$$m(t) \frac{dv}{dt} = -u \frac{dm}{dt} \quad (59)$$

or

$$\frac{dm}{m} = -\frac{1}{u} dv \quad (60)$$

or

$$\ln \frac{m(t)}{m(0)} = -\frac{v(t)}{u} \quad (61)$$

assuming that the rocket starts with zero velocity.

As the fuel burns, the mass of the rocket decreases. Initially the mass of the rocket  $= m_P + m_F + m_S$  when  $m_P$  is the mass of the pay-load,  $m_F$  is the mass of the fuel and  $m_S$  is the mass of the structure. When the fuel is

completely burnt out,  $m_F$  becomes zero and if  $v_B$  is the velocity of the rocket at this stage, when the fuel is all burnt, then (60) gives

$$v_B = u \ln \frac{m_P + m_F + m_S}{m_P + m_S} = u \ln \left(1 + \frac{m_F}{m_P + m_S}\right) \quad (62)$$

This is the maximum velocity that the rocket can attain and it depends on the velocity  $u$  of efflux of gases and the ratio  $m_F/(m_P + m_S)$ . The larger the values of  $u$  and  $m_F/(m_P + m_S)$ , the larger will be the maximum velocity attained.



For the best modern fuels and structural materials, the maximum velocity this gives is about 7 km/sec. In practice it would be much less since we have neglected air resistance and gravity, both of which tend to reduce the velocity. However if a rocket is to place a satellite in orbit, we require a velocity of more than 7 km/sec.

The problem can be overcome by using the concept of multi-stage rockets.

The fuel may be carried in a number of containers and when the fuel of a container is burnt up, the container is thrown away, so that the rocket has not to carry any dead weight.

Thus in a three-stage rocket, let  $m_{F_1}$ ,  $m_{F_2}$ ,  $m_{F_3}$  be the masses of the fuels and  $m_{S_1}$ ,  $m_{S_2}$ ,  $m_{S_3}$  be the three corresponding masses of containers, then velocity at the end of the first stage is

$$v_1 = u \ln \frac{m_P + m_{F_1} + m_{S_1} + m_{F_2} + m_{S_2} + m_{F_3} + m_{S_3}}{m_P + m_{F_2} + m_{S_2} + m_{F_3} + m_{S_3}} \quad (63)$$

At the end the second stage, the velocity is

$$v_2 = v_1 + u \ln \frac{m_P + m_{F_2} + m_{F_3} + m_{S_3}}{m_P + m_{F_3} + m_{S_3}} \quad (64)$$

and at the end of the third stage, the velocity

$$v_3 = v_2 + u \ln \frac{m_P + m_{F_3}}{m_P} \quad (65)$$

In this way, a much larger velocity is obtained than can be obtained by a single-stage rocket.

## MATHEMATICAL MODELLING OF GEOMETRICAL PROBLEMS THROUGH ORDINARY DIFFERENTIAL EQUATIONS OF FIRST ORDER

### Simple Geometrical Problems

Many geometrical entities can be expressed in terms of derivatives and as such relations between these entities can give rise to differential equations whose solution will give us a family of curves for which the given relation between geometrical entities is satisfied.

(i) Find curves for which tangent at a point is always perpendicular to the line joining the point to the origin.

The slope of the tangent is  $dy/dx$  and the slope of line joining the point  $(x, y)$  to the origin is  $y/x$  and since these lines are given to be orthogonal

$$\frac{dy}{dx} = -\frac{x}{y} \quad (66)$$

Integrating

$$x^2 + y^2 = a^2 \quad (67)$$

which represents a family of concentric circle.




Figure 2.9

(ii) Find curves for which the projection of the normal on the x-axis is of constant length.

This condition gives

$$y \frac{dy}{dx} = k \quad (68)$$

Integrating

$$y^2 = 2kx + A, \quad (69)$$

which represents a family of parabolas, all with the same axis and same length of latus rectum.

(iii) Find curves for which tangent makes a constant angle with the radius vector.

Here it is convenient to use polar coordinates and the conditions of the problem gives

$$r \frac{d\theta}{dr} = \tan \alpha \quad (70)$$

Integrating

$$r = Ae^{\theta \cot \alpha}, \quad (71)$$

which represents a family of equiangular spirals.



## Orthogonal Trajectories

Let

$$f(x, y, a) = 0 \quad (72)$$

represent a family of curves, one curve for each value of the parameter  $a$ .

Differentiating (72), we get

$$\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx} = 0 \quad (73)$$

Eliminating  $a$  between (72) and (73), we get a differential equation of the first order

$$\phi\left(x, y, \frac{dy}{dx}\right) = 0, \quad (74)$$

of which (72) is the general solution.

Now we want a family of curves cutting every member of (72) at right angle at all points of intersection.

At a point of intersection of the two curves,  $x, y$  are the same but the slope of the second curve is negative reciprocal of the slope of the first curve. As such differential equation of the family of orthogonal trajectories is




Figure 2.10

$$\phi\left(x, y, -\frac{1}{dy/dx}\right) = 0 \quad (75)$$

Integrating (75), we get

$$g(x, y, b) = 0, \quad (76)$$

which give the orthogonal trajectories of the family (72).

(i) Let the original family be  $y = mx$ , when  $m$  is a parameter then

$$\frac{dy}{dx} = m$$

and eliminating  $m$ , we get the differential equation of this concurrent family of straight lines as

$$\frac{y}{x} = \frac{dy}{dx} \quad (77)$$

To get the orthogonal trajectories, we replace  $dy/dx$  by  $-1/(dy/dx)$  to get

$$\frac{y}{x} = -\frac{1}{dy/dx}$$

Integrating

$$x^2 + y^2 = a^2 \quad (78)$$

which gives the orthogonal trajectories as concentric circles (Figure 2.9a).

(i) Find the orthogonal trajectories of the family of confocal conics

$$\frac{x^2}{a^2 + \lambda} + \frac{y^2}{b^2 + \lambda} = 1, \quad (79)$$

where  $\lambda$  is a parameter. Differentiating, we get

$$\frac{x}{a^2 + \lambda} + \frac{y}{b^2 + \lambda} \frac{dy}{dx} = 0 \quad (80)$$

Eliminating  $\lambda$  between (79) and (80), we get

$$(xp - y)(x + py) = p(a^2 - b^2); \quad p = \frac{dy}{dx} \quad (81)$$

To get the orthogonal trajectories, we replace  $p$  by  $-\frac{1}{p}$  to get

$$\left(-\frac{x}{p}, -y\right)\left(x - \frac{y}{p}\right) = -\frac{1}{p}(a^2 - b^2)$$

or

$$(xp - y)(x + py) = p(a^2 - b^2) \quad (82)$$

However (81) and (82) are identical. As such the family of confocal conics is self-orthogonal, i.e. for every conic of the family, there is another with same focii which cuts it at right angles.




Figure 2.11

One family consists of confocal ellipses and the other consists of confocal hyperbolas with the same focii (Figure 2.11).

(iii) In polar coordinates after getting the differential equation of the family of curves, we have to replace  $r \frac{d\theta}{dr}$  by  $-1 / \left(r \frac{d\theta}{dr}\right)$  and then integrate the resulting differential equation.

Then if the original family is

$$r = 2a \cos \theta, \quad (83)$$

with  $a > 0$  as a parameter, its differential equation is obtained by eliminating  $a$  between (83) and

$$\frac{dr}{d\theta} = -2a \sin \theta \quad (84)$$

to get

$$r \frac{d\theta}{dr} = -\cot \theta$$


Replacing  $r \frac{d\theta}{dr}$  by  $-\left(r \frac{d\theta}{dr}\right)^{-1}$ , we get

$$r \frac{d\theta}{dr} = \tan \theta$$

Integrating we get

$$r = 2b \sin \theta$$

The orthogonal trajectories are shown in Figure 2.12.



The circles of both families pass through the origin, but while the centre of one family lie on  $x$ -axis, the centres of the orthogonal family lie on  $y$ -axis.



**POSSIBLE QUESTIONS**

**Part B (6 Marks)**

1. Write a note on Radio – Active Decay.
2. Discuss a simple Compartment Model.
3. Give a brief note on diffusion of a medicine in the blood stream.
4. Suppose the population of the world now is 4 billion and its doubling period is 35 years,  
what will be the population of the world after 350 years?
5. Design a mathematical model for motion of a rocket.
6. Give an explanatory note on simple compartment models
7. Explain about simple harmonic motion.
8. Discuss in detail about motion under gravity in a resisting medium.
9. Find the relation between doubling, tripling and quadrupling times a population.

**Part C (10 Marks)**

1. Discuss about logistic law of population growth.
2. Discuss a simple Compartment Model.
3. Give an explanatory note on simple compartment models.
4. Explain about simple geometric problems.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
 (Deemed to be University Established Under Section 3 of UGC Act 1956)  
 Pollachi Main Road, Eachanari (Po),  
 Coimbatore –641 021

**UNIT-I**

**Subject: Mathematical Modeling**

**Subject Code: 16MMP303**

**Mathematical Modelling Through ODE of first order**  
**Part-A(20X1=20 Marks)**  
**(Question Nos. 1 to 20 Online Examinations)**  
**Multiple Choice Questions**

Question	Opt 1	Opt 2	Opt 3	Opt 4	Answer
In growth if sciences and scientists the no of scientist should grow	Logically	exponentially	inversely	proportionally	Exponentially
If there is immigration into the population from outside at a rate _____ to the population size	Logically	exponentially	inversely	proportionally	Proportionally
e is the amount of an initial capital of 1 unit invested for 1 unit of time then the interest at unit rate is _____ continuously	simple	compounded	cumulative	principle	compounded
e is the amount of an initial capital of 1 unit invested for _____ unit of time then the interest at unit rate is compounded continuously	1	2	3	4	1
In radio geology the of age solar system is used to estimate _____	Radio active	Diffusion	Decay	immigration	Radio active
The ratio of radio carbon to ordinary carbon in dead plants and animals enables to estimate their _____	Time of birth	time of death	Time of dating	None	Time of death
_____ law is used in the model 'decrease of temperature'	Fick's	Hooke's	Newton's	Gauss	Newton's
_____ law is used in the model 'Diffusion'	Fick's	Hooke's	Newton's	Gauss	Fick's
If P(t) price of commodity and its rate of change is proportional to the _____ between demand and supply	Addition	Difference	Division	Multiplication	Difference
If P(t) price of commodity and its rate of change is _____ to the difference between demand and supply	Logically	exponentially	inversely	Proportional	Proportional
In the model 'change of price of commodity S(t) denotes	System	Supply	Size	None	Supply
In the model 'change of price of commodity d(t) denotes	Demand	Death	Decrease	Diffusion	Demand
In the model 'change of price of commodity $p_e$ denotes	Equilibrium price	Eligible price	Essential price	Evaluation price	Equilibrium price
As population increases the birth rate be decrease and death rate be _____	Increases	stable	decreases	None	Increases
As population increases the birth rate be _____ and death rate be increases	Increases	stable	decreases	None	Decreases
In the model spread of technological innovation and infestious diseases $kN(R-N)$ , R denotes	Total no of companies	companies adopted technological innovation	region	rate	Total no of companies
In rate of dissonation CO be _____ concentration	Maximum	Minimum	Both	None	Maximum
Two chemical substances combined in the ratio _____ to form the third substances Z	a:b	a:2b	2a:b	a:3b	a:b
The gain in amount of a substance in a medium in any time is _____ to the excess of the amount that has entered the medium	Equal	Proportional	Linear	Exponential	Equal
The gain in amount of a substance in a medium in any time is equal to the excess of the amount that has _____ the medium	exit	entered	outer	None	entered
A particle moves in a straight line then its acceleration is _____ to its distance from the origin	Logically	exponentially	inversely	Proportional	proportional
A particle moves in a straight line then its acceleration is proportional to its distance from the origin states that	SHM	MOR	MUG	None	SHM
A particle falls under _____ in a medium in which resistance is proportional to the velocity	Gravity	Sense	Force	Mass	Gravity
A particle falls under gravity in a medium in which resistance is proportional to the _____	Velocity	Sense	Force	Mass	Velocity
The equation of motion which states that mass x acceleration in any direction is _____ on the particle	Velocity	Sense	Force	Mass	Force
A rocket moves forward because of the large _____ velocity	Ultra	Supersonic	Infrared	None	Supersonic
$m(t)$ be the mass of rocket at time t with velocity $v(t)$ then momentum is	$m(t)+v(t)$	$m(t)v(t)$	$m(t)-v(t)$	$m(t)/v(t)$	$m(t)v(t)$
mass of the rocket = $m_F+m_P+m_S$ then F is	Fuel	pay load	structure	ferrocity	Fuel
mass of the rocket = $m_F+m_P+m_S$ then P is	Pressure	pay load	structure	ferrocity	pay load
mass of the rocket = $m_F+m_P+m_S$ then S is	System	pay load	structure	ferrocity	structure
Curves for which tangent at a point is _____ to the line joining the point to the origin.	Equal	Proportional	Perpendicular	Exponential	Perpendicular
Curves for which tangent at a point is perpendicular to the line joining the point to the _____.	centre	point	origin	parallel	origin
Curves for which the projection of the normal on the x axis is of _____ length.	Variable	Constant	unit	y axis	Constant



Curves for which the _____ of the normal on the x axis is of constant length.	Projection	Property	Process	parameter	Projection
Curves for which the projection of the _____ on the x axis is of constant length.	Normal	Proportional	Perpendicular	Exponential	Normal
Curves for which _____ makes a constant angle with the radius vector.	tangent	Secant	Cosecant	Cot	tangent
Curves for which tangent makes a constant angle with the _____ vector.	diameter	radius	unit	scalar	radius
Curves for which tangent makes a _____ angle with the radius vector.	Variable	Constant	unit	y axis	Constant
The point of intersection of two curves the slope of second curve is _____ reciprocal of the first curve.	Positive	negative	unity	trajective	negative
The point of intersection of two curves the slope of second curve is negative _____ of the first curve.	Proportional	reciprocal	Exponential	Logically	reciprocal
The circles of both families pass through _____	Point	centre	Origin	None	Origin
The centres of one family lie on x axis the centres of orthogonal family lie on _____	X axis	Y axis	Both axes	None	Y axis
The centres of one family lie on x axis the centres of _____ family lie on Y axis.	Proportional	linear	unit	orthogonal	orthogonal
The centres of one family lie on _____ the centres of orthogonal family lie on Y axis.	X axis	Y axis	Both axes	None	X axis
The family of confocal conics are _____	self orthogonal	proportional	orthogonal	linear	self orthogonal



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

---

## UNIT II

Mathematical Modeling through Systems of Ordinary Differential Equations of First Order: Population Dynamics – Epidemics – Compartment Models – Economics – Medicine, Arms Race, Battles and International Trade – Dynamics.

### SUGGESTED READINGS

#### TEXT BOOK

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

#### REFERENCES

**R2:** Brain Albright, 2012. Mathematical modeling with Excel, Jones and Bartlett Publishers, New Delhi.

**R3:** Frank. R. Giordano, Maurice. D. Weir, William P. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.

## MATHEMATICAL MODELLING IN POPULATION DYNAMICS

### Prey-Predator Models

Let  $x(t)$ ,  $y(t)$  be the populations of the prey and predator species at time  $t$ . We assume that

(i) if there are no predators, the prey species will grow at a rate proportional to the population of the prey species,

(ii) if there are no prey, the predator species will decline at a rate proportional to the population of the predator species,

(iii) the presence of both predators and preys is beneficial to growth of predator species and is harmful to growth of prey species. More specifically the predator species increases and the prey species decreases at rates proportional to the product of the two populations.

These assumptions give the systems of non-linear first order ordinary differential equations

$$\frac{dx}{dt} = ax - bxy = x(a - by), \quad a, b > 0 \quad (1)$$

$$\frac{dy}{dt} = -py + qxy = -y(p - qx), \quad p, q > 0 \quad (2)$$

Now  $dx/dt$ ,  $dy/dt$  both vanish if

$$x = x_e = \frac{p}{q}, \quad y = y_e = \frac{a}{b} \quad (3)$$

If the initial populations of prey and predator species are  $p/q$  and  $a/b$  respectively, the populations will not change with time. These are the equilibrium sizes of the populations of the two species. Of course  $x = 0$ ,  $y = 0$  also gives another equilibrium position.

From (1) and (2)

$$\frac{dy}{dx} = - \frac{y(p - qx)}{x(a - by)} \quad \frac{dy/dt}{dx/dt} \quad (4)$$

or

$$\frac{a - by}{y} dy = - \frac{p - qx}{x} dx; \quad x_0 = x(0), \quad y_0 = y(0) \quad (5)$$



Integrating

$$a \ln \frac{y}{y_0} + p \ln \frac{x}{x_0} = b(y - y_0) + q(x - x_0) \quad (6)$$

Thus through every point of the first quadrant of the  $x$ - $y$  plane, there is a unique trajectory. No two trajectories can intersect, since intersection will imply two different slopes at the same point.

If we start with  $(0, 0)$  or  $(p/q, a/b)$ , we get point trajectories. If we start with  $x = x_0, y = 0$ , from (1) and (2), we find that  $x$  increases while  $y$  remains zero. Similarly if we start with  $x = 0, y = y_0$ , we find that  $x$  remains zero while  $y$  decreases. Thus positive axes of  $x$  and  $y$  give two line trajectories (Figure 3.1).




Figure 3.1

Since no two trajectories intersect, no trajectory starting from a point situated within the first quadrant will intersect the  $x$ -axis and  $y$ -axis trajectories. Thus all trajectories corresponding to positive initial populations will lie strictly within the first quadrant. Thus if the initial populations are positive, the populations will be always positive. If the population of one (or both) species is initially zero, it will always remain zero.

The lines through  $(p/q, a/b)$  parallel to the axes of coordinates divide the first quadrant into four parts I, II, III and IV. Using (1), (2), we find that

in I,	$dx/dt < 0,$	$dy/dt > 0,$	$dy/dx < 0$
in II,	$dx/dt < 0,$	$dy/dt < 0,$	$dy/dx > 0$
in III,	$dx/dt > 0,$	$dy/dt < 0,$	$dy/dx < 0$
in IV,	$dx/dt > 0,$	$dy/dt > 0,$	$dy/dx > 0$

This give the direction field at all points as shown in Figure 3.1. Each trajectory is a closed convex curve. These trajectories appear relatively cramped near the axes.

In I and II, prey species decreases and in III and IV, it increases. Similarly in IV and I, predator species increases and in II and III, it decreases. After a certain period, both species return to their original sizes and thus both species sizes vary periodically with time.

## Competition Models

Let  $x(t)$  and  $y(t)$  be the populations of two species competing for the same resources, then each species grows in the absence of the other species, and the rate of growth of each species decreases due to the presence of the other species. This gives the system of differential equations

$$\frac{dx}{dt} = ax - bxy = bx\left(\frac{a}{b} - y\right); \quad a > 0, \quad b > 0. \quad (7)$$

$$\frac{dy}{dt} = py - qxy = y(p - qx) = qy\left(\frac{p}{q} - x\right); \quad p > 0, \quad q > 0 \quad (8)$$

There are two equilibrium positions viz.  $(0, 0)$  and  $(p/q, a/b)$ . There are two point trajectories viz.  $(0, 0)$  and  $(p/q, a/b)$  and there are two line trajectories viz.  $x = 0$  and  $y = 0$ .

$$\text{In I} \quad \frac{dx}{dt} < 0, \quad \frac{dy}{dt} < 0, \quad \frac{dy}{dx} > 0 \quad (9)$$

$$\text{In II} \quad \frac{dx}{dt} < 0, \quad \frac{dy}{dt} > 0, \quad \frac{dy}{dx} < 0$$

$$\text{In III} \quad \frac{dx}{dt} > 0, \quad \frac{dy}{dt} > 0, \quad \frac{dy}{dx} > 0$$

$$\text{In IV} \quad \frac{dx}{dt} > 0, \quad \frac{dy}{dt} < 0, \quad \frac{dy}{dx} < 0 \quad (10)$$


This gives the direction field as shown in Figure 3.2. From (7) and (8)

$$\frac{dy}{dx} = \frac{y(p - qx)}{x(a - by)} \quad \text{or} \quad \frac{a - by}{y} dy = \frac{p - qx}{x} dx \quad (11)$$

Integrating

$$a \ln \frac{y}{y_0} - b(y - y_0) = p \ln \frac{x}{x_0} - q(x - x_0) \quad (12)$$





The trajectory which passes through  $(p/q, a/b)$  is

$$a \ln \frac{by}{a} - by + a = p \ln \frac{qx}{p} - qx + p \quad (13)$$

If the initial populations correspond to the point  $A$ , ultimately the first species dies but and the second species increases in size to infinity. If the initial populations correspond to the point  $B$ , then ultimately the second species dies out and the first species tends to infinity. Similarly if the initial populations correspond to point  $C$ , the first species dies out and the second species goes to infinity and if the initial populations correspond to point  $D$ , the second species dies out and the first species goes to infinity.

If the initial populations correspond to point  $E$  or  $F$ , the species populations converge to equilibrium populations  $p/q, a/b$  and if the initial population correspond to point  $G, H$ , the first and second species die out respectively.

Thus except when the initial populations correspond to points on curves  $O'E$  and  $O'F$ , only one species will survive in the competition process and the species can coexist only when the initial population sizes correspond to points on the curve  $EF$ .





If  $x_{10}, x_{20}, \dots, x_{n0}$  is an equilibrium position, we can discuss its local stability by substituting

$$x_1 = x_{10} + u_1, \quad x_2 = x_{20} + u_2, \dots, \quad x_n = x_{n0} + u_n \quad (15)$$

14) and getting a system of linear differential equations

$$\begin{aligned} \frac{du_1}{dt} &= c_{11}u_1 + c_{12}u_2 + \dots + c_{1n}u_n \\ \frac{du_2}{dt} &= c_{21}u_1 + c_{22}u_2 + \dots + c_{2n}u_n \\ &\vdots \\ \frac{du_n}{dt} &= c_{n1}u_1 + c_{n2}u_2 + \dots + c_{nn}u_n, \end{aligned} \quad (16)$$

by neglecting squares, products and higher powers of  $u_i$ 's. We can try the solutions  $u_1 = A_1 e^{\lambda t}$ ,  $u_2 = A_2 e^{\lambda t}$ ,  $\dots$ ,  $u_n = A_n e^{\lambda t}$  to get

$$\begin{vmatrix} c_{11} - \lambda & c_{12} & c_{13} & \dots & c_{1n} \\ c_{21} & c_{22} - \lambda & c_{23} & \dots & c_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ c_{n1} & c_{n2} & c_{n3} & \dots & c_{nn} - \lambda \end{vmatrix} = 0 \quad (17)$$

Thus the equilibrium position would be stable if the real parts of all the eigenvalues of the matrix  $[c_{ij}]$  are negative. The conditions for this are given by Routh-Hurwitz criterion which states that all the roots of

$$a_0 x^n + a_1 x^{n-1} + \dots + a_n = 0, \quad a_0 > 0 \quad (18)$$

will have negative real parts if and only if  $T_0, T_1, T_2, \dots$  are positive where

$$T_0 = a_0, \quad T_1 = a_1, \quad T_2 = \begin{vmatrix} a_1 & a_0 \\ a_3 & a_2 \end{vmatrix}, \quad T_3 = \begin{vmatrix} a_1 & a_0 & 0 \\ a_3 & a_2 & a_1 \\ a_5 & a_4 & a_3 \end{vmatrix}$$

$$T_4 = \begin{vmatrix} a_1 & a_0 & 0 & 0 \\ a_3 & a_2 & a_1 & 0 \\ a_5 & a_4 & a_3 & a_2 \\ a_7 & a_6 & a_5 & a_4 \end{vmatrix} \quad (19)$$



This is true if and only if  $a_i > 0$  and either all even-numbered  $T_k$  or all odd-numbered  $T_k$  are positive. Alternatively (18) will have all roots with negative real parts iff this is true for the  $(n - 1)$ th degree equation

$$a_1 x^{n-1} + a_2 x^{n-2} + a_3 x^{n-3} + \dots - \frac{a_0}{a_1} a_3 x^{n-2} - \frac{a_0}{a_1} a_5 x^{n-4} - \dots = 0 \quad (20)$$

The above method will enable us to discuss only local stability of a position of equilibrium, i.e. this will decide that if the populations of different species are changed slightly from these equilibrium values, whether the population sizes will return to their original equilibrium values or not. The problem of discussing the global stability i.e. of discussing whether the populations will return to these equilibrium values, whatever be the magnitudes of the disturbances, is a more difficult problem and it is possible to solve this problem in special cases only.

### Age-Structured Population Models

Let  $x_1(t), x_2(t), \dots, x_p(t)$  be the populations of the  $p$  pre-reproductive age-groups; let  $x_{p+1}(t), \dots, x_{p+q}(t)$  be the populations of  $q$  reproductive age-groups and let  $x_{p+q+1}(t), \dots, x_{p+q+r}(t)$  be the populations of the  $r$  post-reproductive age-groups. Let  $b_{p+1}, b_{p+2}, \dots, b_{p+q}$  be the birth rates in the  $q$  reproductive age-groups, let  $d_i$  be the death rates in the  $i$ th age-group ( $i = 1, 2, \dots, p + q + r$ ) and let  $m_j$  be the rate of migration from the  $j$ th age-group to the  $(j + 1)$ th age-group ( $j = 1, 2, \dots, p + q + r - 1$ ), then we get the system of differential equations

$$\frac{dx_1}{dt} = b_{p+1}x_{p+1} + \dots + b_{p+q}x_{p+q} - (d_1 + m_1)x_1 \quad (21)$$

$$\frac{dx_2}{dt} = m_1x_1 - (d_2 + m_2)x_2$$

$$\dots \dots \dots$$

$$\frac{dx_n}{dt} = m_{n-1}x_{n-1} - d_nx_n; \quad n = p + q + r$$

or

$$\frac{d}{dt} \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix}$$



$$= \begin{bmatrix} -(d_1 + m_1) & 0 & \dots & b_{p+1} & \dots & b_{p+q} & \dots & 0 & 0 \\ m_1 & -(d_2 + m_2) & \dots & 0 & \dots & 0 & \dots & 0 & 0 \\ 0 & m_2 & \dots & 0 & \dots & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & 0 & \dots & m_{n-1} & -d_n \end{bmatrix}$$

$$\times \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix} \quad (22)$$

or

$$\frac{dX}{dt} = AX(t), \quad (23)$$

where  $A$  is a matrix, all of whose diagonal elements are negative, all of whose main subdiagonal elements are positive,  $q$  other elements of the first row are positive and all other elements are zero. Equation (22) has the solution

$$X(t) = \exp(At)X(0) \quad (24)$$

# MATHEMATICAL MODELLING OF EPIDEMICS THROUGH SYSTEMS OF ORDINARY DIFFERENTIAL EQUATIONS OF FIRST ORDER

## A Simple Epidemic Model

Let  $S(t)$  and  $I(t)$  be the number of susceptibles (i.e. those who can get a disease) and infected persons (i.e. those who have already got the disease).

Initially let there be  $n$  susceptible and one infected person in the system so that

$$S(t) + I(t) = n + 1, \quad S(0) = n, \quad I(0) = 1 \quad (25)$$

The number of infected persons grows at a rate proportional to the product of susceptible and infected persons and the number of susceptible persons decreases at the same rate so that we get the system of differential equations

$$\frac{dS}{dt} = -\beta SI, \quad \frac{dI}{dt} = \beta SI, \quad (26)$$

so that

$$\frac{dS}{dt} + \frac{dI}{dt} = 0, \quad S(t) + I(t) = \text{constant} = n + 1 \quad (27)$$

and

$$\frac{dS}{dt} = -\beta S(n + 1 - S), \quad (28)$$

$$\frac{dI}{dt} = \beta I(n + 1 - I).$$

Integrating

$$S(t) = \frac{n(n + 1)}{n + e^{(n+1)\beta t}}, \quad I(t) = \frac{(n + 1)e^{(n+1)\beta t}}{n + e^{(n+1)\beta t}}, \quad (29)$$

so that

$$\lim_{t \rightarrow \infty} S(t) = 0, \quad \lim_{t \rightarrow \infty} I(t) = n + 1 \quad (30)$$







**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

## **UNIT II(Continuation)**

Mathematical Modeling through Systems of Ordinary Differential Equations of First Order: Population Dynamics – Epidemics – Compartment Models – Economics – Medicine, Arms Race, Battles and International Trade – Dynamics.

### **SUGGESTED READINGS**

#### **TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

#### **REFERENCES**

**R2:** Brain Albright, 2012. Mathematical modeling with Excel, Jones and Bartlett Publishers, New Delhi.

**R3:** Frank. R. Giordano, Maurice. D. Weir, William P. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.

### A Susceptible-Infected-Susceptible (SIS) Model

Here, a susceptible person can become infected at a rate proportional to  $SI$  and an infected person can recover and become susceptible again at a rate  $\gamma I$ , so that

$$\frac{dS}{dt} = -\beta SI + \gamma I, \quad \frac{dI}{dt} = \beta SI - \gamma I, \quad (31)$$

which gives

$$\frac{dI}{dt} = (\beta(n+1) - \gamma)I - \beta I^2 \quad (32)$$

### SIS Model with Constant Number of Carriers

Here infection is spread both by infectives and a constant number  $C$  of carriers, so that (30) becomes

$$\begin{aligned} \frac{dI}{dt} &= \beta(I+C)S - \gamma I \\ &= \beta C(n+1) + \beta(n+1-C-\gamma/\beta)I - \beta I^2. \end{aligned} \quad (33)$$

### Simple Epidence Model with Carriers

In this model, only carriers spread the disease and their number decreases exponentially with time as these are identified and eliminated, so that we get

$$\begin{aligned} \frac{dS}{dt} &= -\beta S(t)C(t) + \gamma I(t), & \frac{dI}{dt} &= \beta C(t)S(t) - \gamma I(t), \\ \frac{dC}{dt} &= -\alpha C \end{aligned} \quad (34)$$

so that

$$S(t) + I(t) = S_0 + I_0 = N \text{ (say)}, \quad C(t) = C_0 \exp(-\alpha t) \quad (35)$$

and

$$\frac{dI}{dt} = \beta C_0 N \exp(-\alpha t) - [\beta C_0 \exp(-\alpha t) + \gamma]I \quad (36)$$

### Model with Removal

Here infected persons are removed by death or hospitalisation at a rate proportional to the number of infectives, so that the model is

$$\frac{dS}{dt} = -\beta SI, \quad \frac{dI}{dt} = \beta SI - \gamma I = \beta I \left( S - \frac{\gamma}{\beta} \right)$$



$$= \beta I(S - \rho); \quad \rho = \frac{\gamma}{\beta} \quad (37)$$

with initial conditions

$$\begin{aligned} S(0) = S_0 > 0, \quad I(0) = I_0 > 0, \quad R(0) = R_0 = 0, \\ S_0 + I_0 = N. \end{aligned} \quad (38)$$

## Model with Removal and Immigration

We modify the above model to allow for the increase of susceptibles at a constant rate  $\mu$  so that the model is

$$\frac{dS}{dt} = -\beta SI + \mu, \quad \frac{dI}{dt} = \beta SI - \gamma I, \quad \frac{dR}{dt} = \gamma I. \quad (39)$$

## COMPARTMENT MODELS THROUGH SYSTEMS OF ORDINARY DIFFERENTIAL EQUATIONS

Pharmokinetics (also called drug kinetics or tracer kinetics or multi-compartment analysis) deals with the distribution of drugs, chemicals, tracers or radio-active substances among various compartments of the body where compartments are real or fictitious spaces for drugs.

Let  $x_i(t)$  be the amount of the drug in the  $i$ th compartment at time  $t$ . We shall assume that the amount that can be transferred from the  $i$ th to the  $j$ th compartment ( $j \neq i$ ) in the time interval  $(t, t + \Delta t)$  is  $k_{ij}x_i(t)\Delta t + 0(\Delta t)$  where  $k_{ij}$  is called the transfer coefficient from the  $i$ th to the  $j$ th compartment. The total change  $\Delta x_i$  in time  $\Delta t$  is given by the amount entering the  $i$ th compartment from other compartments which is reduced by the amount leaving the  $i$ th compartment for other compartments including the zeroeth compartment that denotes the outside system.

Thus we get

$$\Delta x_i = - \sum_{\substack{j=0 \\ j \neq i}}^n k_{ij}x_i\Delta t + \sum_{\substack{j=1 \\ j \neq i}}^n k_{ji}x_j\Delta t + 0(\Delta t) \quad (40)$$

Dividing by  $\Delta t$  and proceeding to the limit as  $\Delta t \rightarrow 0$ , we get

$$\frac{dx_i}{dt} = -x_i \sum_{\substack{j=1 \\ j \neq i}}^n k_{ij} + \sum_{\substack{j=1 \\ j \neq i}}^n k_{ji}x_j \quad (41)$$

$$= \sum_{j=1}^n k_{ji}x_j, \quad (i = 1, 2, \dots, n, \quad (42)$$



where we define

$$k_{ii} = - \sum_{j=1, j \neq i}^n k_{ij}, \quad (i = 1, 2, \dots, n) \quad (43)$$

In matrix notation, we have

$$dX/dt = KX, \quad (44)$$

where

$$X(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix}, \quad K = \begin{bmatrix} k_{11} & k_{21} & \cdots & k_{n1} \\ k_{12} & k_{22} & \cdots & k_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ k_{1n} & k_{2n} & \cdots & k_{nn} \end{bmatrix} \quad (45)$$

If  $X = Be^{\lambda t}$ , when  $B$  is a column matrix, (44) gives

$$\lambda Be^{\lambda t} = KBe^{\lambda t} \quad (46)$$

This gives a constant system of equations to determine  $B$  if

$$|K - \lambda I| = 0 \quad (47)$$

where  $I$  is  $n \times n$  unit matrix. Thus  $\lambda$  has to be an eigenvalue of the matrix  $K$ . We note that all the diagonal elements of  $K$  are negative, all the non-diagonal elements are non-negative and the sum of element of every column is greater than or equal to zero. For such a matrix, it can be shown that the real parts of the eigenvalues are always less than or equal to zero, and the imaginary part is non-zero only when the real part is strictly less than zero. Thus if  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the eigenvalues then

$$\begin{aligned} \operatorname{Re}(\lambda_i) &\leq 0 \\ \operatorname{Im}(\lambda_i) &\neq 0 \text{ only if } \operatorname{Re}(\lambda_i) < 0 \end{aligned} \quad (48)$$

If the drug is injected at a constant rate given by the column vector  $D$  with components  $D_1, D_2, \dots, D_n$ , (44) becomes

$$dX/dt = KX + D \quad (49)$$

Equations (44) and (49) constitute the basic equations for the analysis of drug distribution in the  $n$ -compartment system.

## MATHEMATICAL MODELLING IN ECONOMICS BASED ON SYSTEMS OF ORDINARY DIFFERENTIAL EQUATIONS OF FIRST ORDER

### Domar Macro Model

Let  $S(t)$ ,  $I(t)$ ,  $Y(t)$  be the Savings, Investment and National Income at time  $t$ , then it is assumed that

(i) Savings are proportional to national income, so that

$$S(t) = \alpha Y(t), \quad \alpha > 0$$

(ii) Investment is proportional to the rate of increase of national income so that

$$I(t) = \beta Y'(t), \quad \beta > 0 \quad (51)$$

(iii) All savings are invested, so that

$$S(t) = I(t) \quad (52)$$

We get a system of three ordinary differential equations of first order for determining  $S(t)$ ,  $Y(t)$ ,  $I(t)$ . Solving we get

$$Y(t) = Y(0) e^{\alpha t/\beta}, \quad I(t) = \alpha Y(0) e^{\alpha t/\beta} = S(t), \quad (53)$$

so that the national income, investment and savings all increase exponentially.

### Domar First Debt Model

Let  $D(t)$ ,  $Y(t)$  denote the total national debt and total national income respectively, then we assume that

(i) Rate at which national debt changes is proportional to national income so that

$$D'(t) = \alpha Y(t) \quad (54)$$

(ii) National income increases at a constant rate, so that

$$Y'(t) = \beta \quad (55)$$

$$\text{Solving} \quad D(t) = D(0) + \alpha Y(0)t + \frac{1}{2}\alpha\beta t^2 \quad (56)$$

$$Y(t) = Y(0) + \beta t \quad (57)$$

$$\text{so that} \quad \frac{D(t)}{Y(t)} = \frac{D(0) + \alpha Y(0)t + 1/2\alpha\beta t^2}{Y(0) + \beta t} \quad (58)$$

In this model, the ratio of national debt to national income tends to increase without limit.

### Domar's Second Debt Model

In this model, the first assumption remains the same, but the second assumption is replaced by the assumption that the rate of increase of national income is proportional to the national income so that

$$Y'(t) = \beta Y(t) \quad (59)$$



Solving (54) and (59)

$$Y(t) = Y(0)e^{\beta t} \quad (60)$$

$$D(t) = D(0) + \frac{\alpha}{\beta} Y(0)(e^{\beta t} - 1) \quad (61)$$

$$\frac{D(t)}{Y(t)} = \frac{D(0)}{Y(0)e^{\beta t}} + \frac{\alpha}{\beta}(1 - e^{-\beta t}) \quad (62)$$

In this case  $D(t)/Y(t) \rightarrow \alpha/\beta$  as  $t \rightarrow \infty$ . Thus when debt increases at a rate proportional to income, then if the ratio of debt to income is not to increase indefinitely, income must increase exponentially.

## Samuelson's Investment Model

Let  $K(t)$  represent the capital and  $I(t)$  the investment at time  $t$ , then we assume that

(i) the investment gives the rate of increase of capital so that

$$\frac{dK}{dt} = I(t) \quad (69)$$

(ii) the deficiency of capital below a certain equilibrium level leads to an acceleration of the rate of investment proportional to this deficiency and a surplus of capital above this equilibrium level leads to a declaration of the rate of investment, again proportional to the surplus, so that

$$\frac{dI}{dt} = -m(K(t) - K_e), \quad (70)$$

where  $K_e$  is the capital equilibrium level. If  $k(t) = K(t) - K_e$ , we get

$$\frac{dk}{dt} = I(t), \quad \frac{dI}{dt} = -mk(t), \quad (71)$$

so that

$$-mk(t) = \frac{dI}{dt} = \frac{dI}{dk} \frac{dk}{dt} = I \frac{dI}{dk} \quad (72)$$

Integrating

$$I^2 = m(k_0^2 - k^2); \quad k_0 = k(0); \quad I(0) = 0, \quad (73)$$

so that

$$\frac{dk}{dt} = -\sqrt{m} \sqrt{k_0^2 - k^2} \quad (74)$$



and 
$$k(t) = k(0) \cos \sqrt{m} t \quad (75)$$

$$I(t) = -k(0) \sqrt{m} \sin \sqrt{m} t \quad (76)$$

so that both  $k(t)$  and  $I(t)$  oscillate with a time period  $2\pi/\sqrt{m}$ .

It will be noted that if we put  $k(t) = x(t)$ ,  $I(t) = v(t)$ , equations (71) are the equations for simple harmonic motion. Thus the mathematical models for the oscillation of a particle in a simple harmonic motion and for the oscillation of capital about its equilibrium value are the same.

### Samuelson's Modified Investment Model

In this case, the rate of investment is slowed not only by excess capital as before, but it is also slowed by a high investment level so that (71) become

$$\frac{dk}{dt} = I(t), \quad \frac{dI}{dt} = -mk(t) - nI(t), \quad (77)$$

so that

$$I \frac{dI}{dk} + mk(t) + nI(t) = 0, \quad (78)$$

or

$$\frac{d^2k}{dt^2} + n \frac{dk}{dt} + mk = 0, \quad (79)$$

which are the equations for damped harmonic motion corresponding to the case when a particle performing SHM is acted as by a resistance force proportional to the velocity.

### MATHEMATICAL MODELS IN MEDICINE, ARMS RACE BATTLES AND INTERNATIONAL TRADE IN TERMS OF SYSTEMS OF ORDINARY DIFFERENTIAL EQUATIONS

### 3.5.1 A Model for Diabetes Mellitus

Let  $x(t)$ ,  $y(t)$  be the blood sugar and insulin levels in the blood stream at time  $t$ . The rate of change  $dy/dt$  of insulin level is proportional to (i) the

excess  $x(t) - x_0$  of sugar in blood over its fasting level, since this excess makes the pancreas secrete insulin into the blood stream (ii) the amount  $y(t)$  of insulin since insulin left to itself tends to decay at a rate proportional to its amount and (iii) the insulin dose  $d(t)$  injected per unit time. This gives

$$\frac{dy}{dt} = a_1 (x - x_0) H(x - x_0) - a_2 y + a_3 d(t), \quad (95)$$

where  $a_1, a_2, a_3$  are positive constants and  $H(x)$  is a step function which takes the value unity when  $x > 0$  and taken the value zero otherwise. This occurs in (95) because if blood sugar level is less than  $x_0$ , there is no secretion of insulin from the pancreas.

Again the rate of change  $dx/dt$  of sugar level is proportional to (i) the product  $xy$  since the higher the levels of sugar and insulin, the higher is the metabolism of sugar (ii)  $x_0 - x$  since if sugar level falls below fasting level, sugar is released from the level stores to raise the sugar level to normal (iii)  $x - x_0$  since if  $x > x_0$ , there is a natural decay in sugar level proportional to its excess over fasting level (iv) function of  $t - t_0$  where  $t_0$  is the time at which food is taken

$$\begin{aligned} \frac{dx}{dt} = & -b_1 xy + b_2(x_0 - x) H(x_0 - x) - b_3(x - x_0) H(x - x_0) \\ & + b_4 z(t - t_0), \end{aligned} \quad (96)$$

where a suitable form for  $z(t - t_0)$  can be

$$\begin{aligned} z(t - t_0) &= 0, & t < t_0 \\ &= Qe^{-\alpha(t-t_0)}, & t > t_0 \end{aligned} \quad (97)$$

Equations (95) and (96) give two simultaneous differential equations to determine  $x(t)$  and  $y(t)$ . These equations can be numerically integrated.



### Richardson's Model for Arms Race

Let  $x(t)$ ,  $y(t)$  be the expenditures on arms by two countries  $A$  and  $B$ , then the rate of change  $dx/dt$  of the expenditure by the country  $A$  has a term proportional to  $y$ , since the larger the expenditure in arms by  $B$ , the larger will be the rate of expenditure on arms by  $A$ . Similarly it has a term proportional to  $(-x)$  since its own arms expenditure has an inhibiting effect on the rate of expenditure on arms by  $A$ . It may also contain a term independent of the expenditures depending on mutual suspicions or mutual goodwill. With these considerations, Richardson gave the model

$$\frac{dx}{dt} = ay - mx + r, \quad \frac{dy}{dt} = bx - ny + s \quad (98)$$

Here  $a, b, m, n$  are all  $> 0$ .  $r$  and  $s$  will be positive in the case of mutual suspicions and negative in the case of mutual goodwill.

A position of equilibrium  $x_0, y_0$ , if it exists, will be given by

$$\begin{aligned} amx_0 - say_0 - r &= 0 \\ bx_0 - ny_0 + s &= 0 \end{aligned} \quad \text{or} \quad \frac{x_0}{as - nr} = \frac{y_0}{-br - ms} = \frac{1}{-mn + ab}$$

$$\text{or} \quad x_0 = \frac{as + nr}{mn - ab}, \quad y_0 = \frac{ms + br}{mn - ab} \quad (99)$$

If  $r, s$  are positive, a position of equilibrium exists if  $ab < mn$ . If  $X = x - x_0$ ,  $Y = y - y_0$ , we get

$$\frac{dX}{dt} = aY - mX, \quad \frac{dY}{dt} = bX - nY \quad (100)$$

$X = Ae^{\lambda t}$ ,  $Y = Be^{\lambda t}$  will satisfy these equations if

$$\begin{vmatrix} \lambda + m & -a \\ -b & \lambda + n \end{vmatrix} = 0, \quad \lambda^2 + \lambda(m + n) + mn - ab = 0 \quad (101)$$

Now the following cases arise:

(i)  $mn - ab > 0$ ,  $r > 0$ ,  $s > 0$ . In this case  $x_0 > 0$ ,  $y_0 > 0$  and from (101)  $\lambda_1 < 0$ ,  $\lambda_2 < 0$ . As such there is a position of equilibrium and it is stable.

(ii)  $mn - ab > 0$ ,  $r < 0$ ,  $s < 0$ , there is no position of equilibrium since  $x_0 < 0$ ,  $y_0 < 0$ . However since  $\lambda_1 < 0$ ,  $\lambda_2 < 0$ ,  $X(t) \rightarrow 0$ ,  $Y(t) \rightarrow 0$  as  $t \rightarrow \infty$ , so that  $x(t) \rightarrow x_0$ ,  $y(t) \rightarrow y_0$ . However  $x_0$  and  $y_0$  are negative and populations cannot become negative. In any case to become negative, they have to pass through zero values. As such, as  $x(t)$  becomes zero, (98) is modified to

$$\frac{dy}{dt} = -ny + s \quad (102)$$



and since  $s < 0$ ,  $y(t)$  decreases till it reaches zero. Similarly if  $y(t)$  becomes zero first, (98) is modified to

$$\frac{dx}{dt} = -mx + r, \quad (103)$$

and since  $r < 0$ ,  $x(t)$  decreases till it reaches zero. Thus if  $mn - ab > 0$ ,  $r < 0$ ,  $s < 0$ , there will ultimately be complete disarmament.

(iii)  $ma - ab < 0$ ,  $r > 0$ ,  $s > 0$ . These give  $x_0 < 0$ ,  $y_0 < 0$ , one of  $\lambda_1, \lambda_2$  is positive and the other is negative. In this case there will be a run-away arms race.

(iv)  $ma - ab < 0$ ,  $r < 0$ ,  $s < 0$ . These give  $x_0 > 0$ ,  $y_0 > 0$  one of  $\lambda_1, \lambda_2$  is positive and the other is negative. In this case there will be a run-away arms race or disarmament depending on the initial expenditure on arms.

## Lanchester's Combat Model

Let  $x(t)$  and  $y(t)$  be the strengths of the two forces engaged in combat and let  $M$  and  $N$  be the fighting powers of individuals depending on physical fitness, types of arms and training, then Lanchester postulated that the reduction in strength of each force is proportional to the effective fighting strength of the opposite force, so that

$$\frac{dx}{dt} = -ayN, \quad \frac{dy}{dt} = -axM \quad (104)$$

giving  $\frac{dx}{yN} = \frac{dy}{xM}$  or  $Mx^2 - Ny^2 = \text{constant}$  (105)

If the proportional reduction of strengths in the two forces are the same

$$\frac{1}{x} \frac{dx}{dt} = \frac{1}{y} \frac{dy}{dt} \quad \text{or} \quad \frac{Ny}{x} = \frac{Mx}{y} \quad \text{or} \quad Mx^2 = Ny^2 \quad (106)$$

This is the square law. The fighting strength of an army depends on the square of its numerical strength and directly on the fighting quality of individuals.

## International Trade Model

Since international trade is beneficial to all parties, we can consider the model





## Motion of a Projectile

A particle of mass  $m$  is projected from the origin in vacuum with velocity  $V$  inclined at an angle  $\alpha$  to the horizontal. Suppose at time  $t$ , it is at position  $x(t), y(t)$  and its horizontal and vertical velocity components are  $u(t), v(t)$  respectively, then the equations of motion are:

$$m \frac{du}{dt} = 0 \quad m \frac{dv}{dt} = -mg \quad (108)$$




Figure 3.3

Integrating  $u = V \cos \alpha, \quad v = V \sin \alpha - gt, \quad (109)$

so that  $\frac{dx}{dt} = V \cos \alpha, \quad \frac{dy}{dt} = V \sin \alpha - gt \quad (110)$

Integrating again  $x = V \cos \alpha t, \quad y = V \sin \alpha t - \frac{1}{2}gt^2 \quad (111)$

Eliminating  $t$  between these two equations, we get

$$y = x \tan \alpha - \frac{1}{2} \frac{gx^2}{V^2 \cos^2 \alpha} \quad (112)$$

which is a parabola, since the terms of the second degree form a perfect square. The parabola cuts  $y = 0$ , when

$$x = 0 \quad \text{or} \quad x = \frac{V^2 \sin 2\alpha}{g} \quad (113)$$

corresponding to position 0 and A in Figure 3.3 so that the range of the particle is given by



$$R = \frac{V^2 \sin 2\alpha}{g} \quad (114)$$

Putting  $y = 0$  in (111) we get

$$t = 0 \quad \text{or} \quad t = \frac{2V \sin \alpha}{g} \quad (115)$$

This gives the time  $T$  of flight. Since the horizontal velocity is constant and equal to  $V \cos \alpha$ , the total horizontal distance travelled is

$$V \cos \alpha (2V \sin \alpha / g) = V^2 \sin 2\alpha / g$$

which gives us the same range.

### POSSIBLE QUESTIONS

#### Part B (6 Marks)

1. Discuss in detail prey-predator models.
2. Discuss in detail on Samuelson's investment model.
3. Derive a Simple Epidemic Model.
4. Show that national income, investment and savings increase exponentially.
5. Design any two mathematical models in economics based on ordinary differential equations of first order give by Domar.
6. Give a detailed note on multi-species models.
7. Explain about motion of a projectile.

#### Part C (10 Marks)

1. Discuss in detail on Samuelson's investment model.
2. Explain a simple epidemic model.
3. Discuss in detail Domar Macro model.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
(Deemed to be University Established Under Section 3 of UGC Act 1956)  
Pollachi Main Road, Eachanari (Po),  
Coimbatore -641 021

**UNIT-II**

**Subject: Mathematical Modeling**

**Subject Code: 16MMP303**

**Mathematical Modeling through Systems of ODE First Order**

**Part-A(20X1=20 Marks)**

**(Question Nos. 1 to 20 Online Examinations)**

**Multiple Choice Questions**

Question	Opt 1	Opt 2	Opt 3	Opt 4	Answer
If there are no predators the _____ species will grow at a rate proportional to the population.	Prey	trajectory	permanent	persuieng	Prey
If there are no predators the prey species will grow at a rate _____-1 to the population.	Proportional	reciprocal	Exponential	Logically	Proportional
The predator species _____ and the prey species decreases at a rate proportional to the product of two populations.	increases	decreases	uniformly	stable	increases
The predator species increases and the prey species _____ at a rate proportional to the product of two populations.	increases	decreases	uniformly	stable	decreases
The predator species increases and the prey species decreases at a rate _____ to the product of two populations.	Proportional	linear	unit	orthogonal	Proportional
The predator species increases and the prey species decreases at a rate proportional to the _____ of two populations.	Addition	subtraction	product	division	product
If there are no prey the _____ species will decline at a rate proportional to the population.	Prey	Predator	permanent	persuieng	Predator
If there are no prey the predator species will _____ at a rate proportional to the population.	decline	denied	different	decrease	decline
If there are no prey the predator species will decline at a rate _____ to the population.	Proportional	reciprocal	Exponential	Logically	Proportional
The initial populations of prey and preador species are _____	p/q and a/b	a/b and p/q	a/b	p/q	p/q and a/b
The population of x=0 and y=0 is called _____ position.	zero	equilibrium	unit	none	equilibrium
x(t) and y(t) are the populations of two species competing for the same resources stands _____ model	epidemic	population	dynamic	competition	competition
The rate of growth of each species _____ due to the presence of the other.	increases	decreases	uniformly	stable	decreases
The rate of growth of each species decreases due to the _____ of the other.	presence	absence	both	none	presence
$x_1(t), x_2(t), \dots, x_n(t)$ represent the populations of n species states _____ model.	multi-species	single-species	prey	predator	multi-species
The real parts of all the eigenvalues of the matrix $[c_{ij}]$ is negative are called	Rout-Herwitz	fick's	newtyons	gauss	Rout-Herwitz
Age structured population model deals _____ age groups	productive	reproductive	decline	increase	reproductive
In simple epidemic mode $S(t)$ denotes	susceptible	system	synopsis	success	susceptible
In simple epidemic mode $I(t)$ denotes	Infected	increase	innovation	intensity	Infected
In simple epidemic mode $S(t)+I(t) =$	n	n+1	n-1	2n	n+1
In simple epidemic mode limit t tends to infinity of $S(t)$ denotes	0	1	2	3	0
In simple epidemic mode limit t tends to infinity of $I(t)$ denotes	n	n+1	n-1	2n	n+1
A susceptible person can infected at a rate proportional to	SI	SIS	SHM	MOC	SI
In SIS Infected person can recover and become susceptible at a rate	Gamma I	SI	SHM	SIS	Gamma I
A susceptible person can infected at a rate _____ to SI	Proportional	linear	unit	orthogonal	Proportional
Only carriers spread the disease deals _____ model	Simple epidence	epidemic	SIS	SI	Simple epidence
The model with removal deals the infected persons are removed by _____	Death	moving	migration	none	Death
The model with removal deals the infected persons are removed by _____	Hospitalisation	moving	migration	none	Hospitalisation
Model with removal and immigration allows the _____ of susceptible.	Increases	decreases	decline	equate	Increases
Model with removal and immigration allows the increase of susceptible.	infected	susceptible	preys	predators	susceptible
_____ deals the distribution of drugs , chemicals tracers or radio active.	Pharmokinetics	kinetics	medicine	diffusion	Pharmokinetics
Parmokinetics deals the distribution of _____	Drugs	blood	Glucose	Rice	Drugs
Parmokinetics deals the distribution of _____	Chemicals	blood	Glucose	Rice	Chemicals
Parmokinetics deals the distribution of _____	Tracers	blood	Glucose	Rice	Tracers
Parmokinetics deals the distribution of _____	Radio active	blood	Glucose	Rice	Radio active
In Domar Macro Model $S(t)$ denotes	Savings	Success	Susceptible	System	Savings
In Domar Macro Model $I(t)$ denotes	Increases	Investment	innovation	Instalment	Investment
In Domar Macro Model $Y(t)$ denotes	Income	National Income	Debt	National debt	National Income
In Domar Macro Model savings are proportional to	Income	National Income	Debt	National debt	National Income
In Domar Macro Model Investment is proportional to the rate of increase of _____	Income	National Income	Debt	National debt	National Income
In Domar Macro Model all savings are Investment so that	$S(t) = I(t)$	$S(t) = 1/2 I(t)$	$2S(t) = I(t)$	None	$S(t) = I(t)$



In Domar first Debt model $D(t)$ denotes	debt	total national debt	income	national income	total national debt
In Domar first Debt model $Y(t)$ denotes	income	total income	national income	total national income	total national income
In Domar first Debt model rate at which national debt changes is _____ to the national income.	Proportional	linear	unit	orthogonal	Proportional
In Domar first Debt model rate at which national debt changes is proportional to the _____.	income	total income	national income	total national income	national income
In Domar first Debt model national income increases at a _____ rate.	Variable	Constant	unit	orthogonal	Constant
In Domar first Debt model national income _____ at constant rate.	Increases	decreases	decline	equate	Increases
In Domar's Second Debt model the ratio of debt to on come is not to increase indefinitely income must increase _____	Proportional	reciprocal	Exponential	Logically	Exponential
In Allen's Speculative Model $d(t)$ denotes	demand	supply	price of a commodity	debt	demand
In Allen's Speculative Model $s(t)$ denotes	demand	supply	price of a commodity	System	supply
In Allen's Speculative Model $p(t)$ denotes	demand	supply	price of a commodity	Prey	price of a commodity
In Samuelson's Investment model $K(t)$ denotes	Capital	investment	savings	debt	Capital
In Samuelson's Investment model the investment gives rate of increase of _____	Capital	investment	savings	debt	Capital
In Samuelson's Investment model the investment gives rate of _____ of capital.	Increases	decreases	investment	decline	Increases
In Samuelson's Modified Investment model a particle performing _____ is acted by a resistance force proportional to velocity.	SHM	MOC	SIS	none	SHM
In Samuelson's Modified Investment model a particle performing SHM is acted by a resistance force _____ to velocity.	Proportional	linear	unit	orthogonal	Proportional
In a model for Diabetes Mellitus $x(t)$ denotes	blood sugar	salt	urea	fat	blood sugar
In a model for Diabetes Mellitus $y(t)$ denotes	insulin	thyroid	salt	urea	insulin
In Leontief's Inter - Industries relation model, the notation of contribution from the $r$ th industry to $s$ th industry per unit time is	$x_{rs}$	$x_r$	$X_r$	$\xi_r$	$x_{rs}$
In Leontief's Inter - Industries relation model, the notation of contribution from the $r$ th industry to consumers per unit time is	$x_{rs}$	$x_r$	$X_r$	$\xi_r$	$x_r$
In Leontief's Inter - Industries relation model, the notation of total output of the $r$ th industry per unit time is	$x_{rs}$	$x_r$	$X_r$	$\xi_r$	$X_r$
In Leontief's Inter - Industries relation model, the notation of input of the labour in the $r$ th industry is	$x_{rs}$	$x_r$	$X_r$	$\xi_r$	$\xi_r$
In Leontief's Inter - Industries relation model, the notation of price per unit of the product of the $r$ th industry is	$p_r$	$x_r$	$X_r$	$\xi_r$	$p_r$
In Leontief's Inter - Industries relation model, the notation of wage per unit of labour per unit time is	$w$	$x_r$	$X_r$	$\xi_r$	$w$
In Leontief's Inter - Industries relation model, the notation of total labour input into the system is	$Y$	$x_r$	$X_r$	$\xi_r$	$Y$
In Leontief's Inter - Industries relation model, the notation of stock of the product of the $r$ th industry held by the $s$ th industry is	$S_{rs}$	$x_r$	$X_r$	$\xi_r$	$S_{rs}$
In Leontief's Inter - Industries relation model, the notation of stock of the $r$ th industry is	$S_r$	$S_{rs}$	$X_r$	$\xi_r$	$S_r$
The excess of sugar in blood over its fasting level makes _____ secrete insulin into the blood stream.	thyroid	hormone	pancreas	none	pancreas
The fighting strength of an army depends on the _____ of its numerical strength and directly on the fighting quality of individuals.	square	circle	rectangle	ellipse	square
A particle of mass $m$ is projected from the origin in vacuum with velocity inclined at an angle proportional to the _____	vertical	slope	horizontal	equal	horizontal
A particle of mass $m$ is projected from the origin in vacuum with velocity inclined at an angle _____ to the horizontal.	Proportional	reciprocal	Exponential	Logically	proportional
In the case of intercontinental ballistic missiles eating and _____ have to be considered.	aerodynamics	dynamics	mechanics	aeromechanics	aerodynamics
Both range and maximum height of projectile are reduced by _____ resistance.	air	water	liquid	solid	air



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

**Subject: Mathematical Modeling****Subject Code: 16MMP303****L T P C****Class:II M.Sc****Semester:III****4 1 0 4****UNIT III**

Mathematical Modeling through Ordinary Differential Equations of Second Order:  
 Planetary Motions – Circular Motion and Motion of Satellites – Mathematical  
 Modeling through Linear Differential Equations of Second Order – Miscellaneous  
 Mathematical Models.

**SUGGESTED READINGS****TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New  
 Delhi.

**REFERENCES**

**R2:** Brain Albright, (2010). Mathematical Modeling with Excel, Jones and Bartlett  
 Publishers, New Delhi.

**R3:** Frank. R. Giordano, Maurice. D. Weir, William P. Fox, (2003). A first course in  
 Mathematical Modelling, Vikash Publishing House, UK.

## MATHEMATICAL MODELLING OF PLANETARY MOTIONS

Every planet moves mainly under the gravitational attractive force exerted by the Sun. If  $S$  and  $P$  are masses of the Sun and the planet and  $G$  is the universal constant of gravitation, then the forces of gravitational attraction on the Sun and planet are both  $GSP/r^2$ , where  $r$  is the distance between the Sun and the planet. Accordingly the acceleration (Fig. 4.1) of the Sun towards the planet is  $GP/r^2$  and the acceleration of the planet towards the Sun is  $GS/r^2$ . The acceleration of the planet relative to the Sun is

$$G(S + P)/r^2 = \mu/r^2.$$

Now we take the Sun as fixed, then the planet can be said to move under a central force  $\mu/r^2$  per unit mass i.e. under a force which is always directed towards a fixed centre  $S$ .




Figure 4.1

We shall for the present also regard  $P$  as a particle so that to study the motion of the planet, we have to study the motion of a particle moving under a central force. We can take  $S$  as origin so that the central force is always along the radius vector. To study this motion, it is convenient to use polar coordinates and to find the components of the velocity and acceleration along and perpendicular to the radius vector.

### Components of Velocity and Acceleration Vectors along Radial and Transverse Directions

As the particle moves from  $P$  to  $Q$ , the displacement along the radius vector

$$= ON - OP = (r + \Delta r) \cos \Delta\theta - r \quad (1)$$

and the radial component  $u$  of velocity is

$$\begin{aligned} u &= \lim_{\Delta t \rightarrow 0} \frac{(r + \Delta r) \cos \Delta\theta - r}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{\Delta r}{\Delta t} = \frac{dr}{dt} \end{aligned} \quad (2)$$






Figure 4.2

Similarly the displacement perpendicular to the radius vector

$$= (r + \Delta r) \sin \Delta \theta \quad (3)$$

and the transverse component  $v$  of the velocity is given by

$$v = \lim_{\Delta t \rightarrow 0} \frac{(r + \Delta r) \sin \Delta \theta}{\Delta t} = \lim_{\Delta t \rightarrow 0} r \frac{\sin \Delta \theta}{\Delta \theta} \frac{\Delta \theta}{\Delta t} = r \frac{d\theta}{dt} \quad (4)$$

As such the velocity components in polar coordinates are

$$u = \frac{dr}{dt} = r' \quad \text{and} \quad v = r \frac{d\theta}{dt} = r\theta' \quad (5)$$

Now the change in the velocity along the radius vector

$$= (u + \Delta u) \cos \Delta \theta - (v + \Delta v) \sin \Delta \theta - u \quad (6)$$




Figure 4.3

and the radial component of acceleration

$$\begin{aligned} &= \lim_{\Delta t \rightarrow 0} \frac{(u + \Delta u) \cos \Delta \theta - (v + \Delta v) \sin \Delta \theta - u}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{\Delta u - v \Delta \theta}{\Delta t} = \frac{du}{dt} - v \frac{d\theta}{dt} = \frac{d}{dt}(r') - r\theta' \theta' \\ &= r'' - r\theta'^2 \end{aligned} \quad (7)$$

Similarly the transverse component of acceleration

$$\begin{aligned}
 &= \lim_{\Delta t \rightarrow 0} \frac{(u + \Delta u) \sin \Delta\theta + (v + \Delta v) \cos \Delta\theta - v}{\Delta t} \\
 &= \lim_{\Delta t \rightarrow 0} \frac{u \Delta\theta + \Delta v}{\Delta t} \\
 &= u \frac{d\theta}{dt} + \frac{dv}{dt} = r'\theta' + \frac{d}{dt}(r\theta') = \frac{1}{r} \frac{d}{dt}(r^2\theta')
 \end{aligned} \quad (8)$$

Thus the radial and transverse components of acceleration are

$$r'' - r\theta'^2 \quad \text{and} \quad \frac{1}{r} \frac{d}{dt}(r^2\theta') \quad (9)$$

## Motion Under a Central Force

Let the force acting on a particle of mass  $m$  be  $mF(r)$  and let it be directed towards the origin, then the equations of motion are

$$m(r'' - r\theta'^2) = -mF(r) \quad (10)$$

$$\frac{m}{r} \frac{d}{dt}(r^2\theta') = 0 \quad (11)$$

From (11)

$$r^2\theta' = \text{constant} = h \text{ (say)}, \quad (12)$$

then (10) gives

$$r'' - r\theta'^2 = -F(r) \quad (13)$$

We can eliminate  $t$  between (12) and (13) to get a differential equation between  $r$  and  $\theta$ . We find it convenient to use  $u = 1/r$  instead of  $r$ , so that making use of (12), we get

$$r' = \frac{dr}{dt} = \frac{dr}{du} \frac{du}{d\theta} \frac{d\theta}{dt} = -\frac{1}{u^2} \frac{du}{d\theta} \frac{h}{r^2} = -h \frac{du}{d\theta} \quad (14)$$

and

$$\begin{aligned}
 r'' &= \frac{d}{dt} \left( -h \frac{du}{d\theta} \right) = \frac{d}{d\theta} \left( -h \frac{du}{d\theta} \right) \frac{d\theta}{dt} \\
 &= -h \frac{d^2u}{d\theta^2} h u^2 = -h^2 u^2 \frac{d^2u}{d\theta^2}
 \end{aligned} \quad (15)$$

From (12), (13) and (15)

$$-F(r) = -h^2 u^2 \frac{d^2u}{d\theta^2} - \frac{1}{u} h^2 u^4 = -h^2 u^2 \left( \frac{d^2u}{d\theta^2} + u \right)$$

or

$$\frac{d^2u}{d\theta^2} + u = \frac{F}{h^2 u^2}, \quad (16)$$

where  $F$  can be easily expressed as a function of  $u$ . This is the differential equation of the second order whose integration will give the relation between  $u$  and  $\theta$  or between  $r$  and  $\theta$  i.e. the equation of the path described by a particle moving under a central force  $F$  per unit mass.

### Motion Under the Inverse Square Law

If the central force per unit mass is  $\mu/r^2$  or  $\mu u^2$ , Equation (16) gives

$$\frac{d^2u}{d\theta^2} + u = \frac{\mu}{h^2} \quad (17)$$

Integrating this linear equation with constant coefficients, we get

$$u = A \cos(\theta - \alpha) + \frac{\mu}{h^2}$$

$$\text{or } \frac{h^2/u}{r} = \frac{L}{r} = 1 + e \cos(\theta - \alpha); h^2 = \mu L, \quad (18)$$

which represents a conic with a focus at the centre of force. Thus if a particle moves under a central force  $\mu/r^2$  per unit mass, the path is a conic section with a focus at the centre. The conic can be an ellipse, parabola, or hyperbola according as  $e \leq 1$ .

Now the velocity  $V$  of the particle is given by

$$V^2 = \dot{r}^2 + r^2\dot{\theta}^2 = \left(\frac{dr}{du} \frac{du}{d\theta} \frac{d\theta}{dt}\right)^2 + \frac{1}{u^2}(hu^2)^2$$

$$= h^2 \left(\frac{du}{d\theta}\right)^2 + h^2 u^2 \quad (19)$$

Using (18)

$$L \frac{du}{d\theta} = -e \sin(\theta - \alpha) \quad (20)$$

From (19) and (20)

$$V^2 = \mu L \left( \frac{e^2 \sin^2(\theta - \alpha)}{L^2} + \frac{(1 + e \cos(\theta - \alpha))^2}{L^2} \right)$$

$$= \frac{\mu}{L} (1 + e^2 + 2e \cos(\theta - \alpha))$$

$$= \frac{\mu}{L} (e^2 - 1 + 2(1 + e \cos(\theta - \alpha)))$$

$$= \frac{\mu}{L} (e^2 - 1) + \frac{2\mu}{r} \quad (21)$$

$$\begin{aligned} \text{If the path is an ellipse } L &= a(1 - e^2) \\ \text{If the path is a parabola } e &= 1 \end{aligned} \quad (22)$$



If the path is a hyperbola  $L = a(e^2 - 1)$ ,

so that  $V^2 = \mu \left( \frac{2}{r} + \frac{1}{a} \right)$  in the case of a hyperbola

$$= \mu \left( \frac{2}{r} \right) \text{ in the case of a parabola} \quad (23)$$

$$= \mu \left( \frac{2}{r} - \frac{1}{a} \right) \text{ in the case of an ellipse.}$$

Thus if the particle is projected with velocity  $V$  from a point at a distance  $r$  from the centre of force, the path will be a hyperbola, parabola or ellipse according as

$$V^2 - \frac{2\mu}{r} \begin{matrix} \geq 0 \\ < 0 \end{matrix} \quad (24)$$

We have proved that if the central force is  $\mu/r^2$  per unit mass, the path is a conic section with the centre of forces at one focus. Conversely if we know that the path is a conic section

$$\frac{L}{r} = Lu = 1 + e \cos(\theta - \alpha), \quad (25)$$

with a focus at the centre of force, then the force per unit mass is given by

$$\begin{aligned} F &= h^2 u^2 \left( \frac{d^2 u}{d\theta^2} + u \right) \\ &= h^2 u^2 \left( \frac{-e \cos(\theta - \alpha)}{L} + \frac{1 + \cos(\theta - \alpha)}{L} \right) \\ &= \frac{h^2}{L} u^2 = \frac{\mu}{r^2}, \end{aligned} \quad (26)$$

so that the central force follows the inverse square law.

Since all planets are observed to move in elliptic orbits with the Sun at one focus, it follows that the law of attraction between different planets and Sun must be the inverse square law.

### Kepler's Laws of Planetary Motions

On the basis of the long period of observations of planetary motions by his predecessors and by Kepler himself, Kepler deduced the following three laws of motion empirically

- (i) Every planet describes an ellipse with the Sun at one focus
- (ii) The radius vector from the Sun to a planet describes equal areas in equal intervals of time.
- (iii) The squares of periodic time of planets are proportional to the cubes of the semimajor axes of the orbits of the planets

We can deduce all these three laws from the mathematical modelling of planetary motion discussed above, when the law of attraction is the inverse square law.

(i) We have already seen that under the inverse square law, the path has to be a conic section and this includes elliptic orbits.

(ii) Since  $r^2\theta' = h$ , we get

$$\lim_{\Delta t \rightarrow 0} \frac{1}{2} \frac{r^2 \Delta \theta}{\Delta t} = \frac{1}{2} h \quad (27)$$

From Figure 4.2, the area  $\Delta A$  bounded by radius vectors  $OP$  and  $OQ$  and the arc  $PQ$  is  $1/2 r^2 \sin \Delta \theta$  so that (27) gives

$$\frac{dA}{dt} = \frac{1}{2} h, \quad (28)$$

and the rate of description of sectorial area is constant and equal areas are described in equal intervals of time. This is Kepler's second law.

(iii) The total area of the ellipse is  $\pi ab$  and since the areal velocity is  $\frac{1}{2}h$ , the periodic time  $T$  is given by

$$T = \frac{\pi ab}{\frac{1}{2}h} = \frac{2\pi ab}{\sqrt{\mu L}} = \frac{2\pi ab}{\sqrt{\mu} \sqrt{b^2/a}} = \frac{2\pi}{\sqrt{\mu}} a^{3/2} \quad (29)$$

For two different planets of masses  $P_1, P_2$ , and semiaxes of orbits  $a_1, a_2$ , this gives

$$\frac{T_1}{T_2} = \frac{\sqrt{\mu_2}}{\sqrt{\mu_1}} \frac{a_1^{3/2}}{a_2^{3/2}} = \frac{\sqrt{G(S+P_2)}}{\sqrt{G(S+P_1)}} \frac{a_1^{3/2}}{a_2^{3/2}} \quad (30)$$

or

$$\frac{T_1^2}{T_2^2} = \frac{S+P_2}{S+P_1} \frac{a_1^3}{a_2^3} = \frac{1 + \frac{P_2}{S}}{1 + \frac{P_1}{S}} \frac{a_1^3}{a_2^3} \quad (31)$$

Since  $P_1, P_2$  are very small compared with  $S$ , this gives, as a very good approximation

$$\frac{T_1^2}{T_2^2} = \frac{a_1^3}{a_2^3} \quad (32)$$

which is Kepler's third law of planetary motion.



Deduction of Kepler's three laws of planetary motion from the universal law of gravitation was an important success of mathematical modelling. Results which took hundreds of years to obtain by observation could be obtained in a very short time by using mathematical modelling.

Here we have neglected the forces of attraction of other planets on the given planet. These are very small as compared with the attractive force of the Sun. However these can be taken into account. In fact possibly the most sensational achievement of mathematical modelling was achieved when the discrepancies from the above theory observed in the motion of planets were explained as possibly due to the existence of another small planet. The position of this planet, not observed till that time, was calculated, and when the telescope was pointed out to that position in the sky, the planet was there!

Again the occurrence of many of the fundamental particles in physics has been theoretically predicted on the basis of mathematical modelling.

The advantages of developing a successful theoretical model over relying on purely observational and empirical models are that (i) this development can suggest development of mathematical models for similar situations elsewhere and those new models can later be validated and (ii) the theoretical models, unlike empirical models, can be generalised. Thus the model developed by Newton for planetary motion could be easily extended to apply to motion of artificial satellites. Similarly in urban transportation, a gravity model was developed by trial and error and ad hoc empirical methods extending over a period of thirty to forty years. When the same model was obtained theoretically from the principle of maximum entropy, it could be easily generalised for many more complex situations than could ever be handled by the empirical methods.

## MATHEMATICAL MODELLING OF CIRCULAR MOTION AND MOTION OF SATELLITES

### Circular Motion

When a particle moves in a circle of radius  $a$  so that  $r = a$ , the radial component of velocity  $= r' = 0$ , the transverse component of velocity  $=$

$r\theta' = a\theta'$  the radial component of acceleration  $= r'' - r\theta'^2 = -a\theta'^2$ , the transverse component of acceleration  $= \frac{1}{r} \frac{d}{dt} (r^2\theta') = \frac{1}{a} \frac{d}{dt} (a^2\theta') = a\theta''$ .



Thus the velocity is  $a\theta'$  along the tangent and the acceleration has two components  $a\theta''$  along the tangent and  $a\theta'^2$  along the normal.

If a particle moves in a circle of radius  $a$ , its equations of motion are

$ma\theta''$  = external force in the direction of the tangent

$ma\theta'^2$  = external force in the direction of the inward normal.




Figure 4.4

Thus if a particle is attached to one end of a string, the other end of which is fixed and the particle moves in a vertical circle, the equations of motion are (Figure 4.4)

$$ma\theta'' = -mg \sin \theta \quad (33)$$

$$ma\theta'^2 = T - mg \cos \theta \quad (34)$$

If  $\theta$  is small, (33) gives

$$\theta'' = -\frac{g}{a} \theta, \quad (35)$$

which is the equation for a simple harmonic motion. Thus for small oscillations of a simple pendulum, the time period is

$$T = 2\pi\sqrt{a/g} \quad (36)$$

If  $\theta$  is not necessarily small, integration of (33) gives

$$a\theta'^2 = 2g \cos \theta + \text{constant} \quad (37)$$

If the particle is projected from the lowest point with velocity  $u$ , then  $a\theta' = u$  when  $\theta = 0$ , so that

$$a\theta'^2 = \frac{v^2}{a} = \frac{u^2}{a} - 2g(1 - \cos \theta), \quad (38)$$

where  $v$  is the velocity of the particle, so that

$$v^2 = u^2 - 2ga(1 - \cos \theta) \quad (39)$$

$$\text{or} \quad \frac{1}{2} mv^2 = \frac{1}{2} mu^2 - mga(1 - \cos \theta) = \frac{1}{2} mu^2 - mgh \quad (40)$$

where  $h$  is the vertical distance travelled by the particle. Equation (40) can be obtained directly from the principle of conservation of energy. Equation (34) then gives

$$T = m \frac{v^2}{a} + mg \cos \theta = m \frac{u^2}{a} - 2mg + 3mg \cos \theta \quad (41)$$

At the highest point  $\theta = \pi$  and  $T = m \frac{u^2}{a} - 5mg$ . If  $u^2 \geq 5ag$ , the particle will move in the complete vertical circle again and again. However if


$u^2 < 5ag$ , tension will vanish before the particle reaches the highest point. When the tension vanishes, the particle begins to move freely under gravity and describes a parabolic path till the string again becomes tight and the circular motion is started again.

### Motion of a Particle on a Smooth or Rough Vertical Wire


(a) If the particle moves on the inside of a smooth wire, the equations of motion (Fig. 4.5a) are:

$$ma\theta'' = -mg \sin \theta \quad (42)$$

$$ma\theta'^2 = R - mg \cos \theta \quad (43)$$



(a)



(b)

Figure 4.5

These are the same as (33) and (34) when  $T$  is replaced by the normal reaction  $R$ . As such if  $u^2 \geq 5ag$ , the particle makes an indefinite number of complete rounds of the circular wire. If  $u^2 < 5ag$ , the reaction vanishes before the particle reaches the highest point, the particle leaves the curve, describes a parabolic path till it meets the circular wire again and it again describes a circular path. This motion is repeated again and again.

(b) If the particle moves on the outside of the smooth vertical wire (Fig. 4.5b), the equations of motion are

$$ma\theta'' = mg \sin \theta \quad (44)$$

$$ma\theta'^2 = -R + mg \cos \theta \quad (45)$$



Integrating (44)  $\theta'^2 = u^2 + 2ga(1 - \cos \theta)$  (46)

Using (45)  $R = 3mg \cos \theta - \frac{mu^2}{a} - 2mg$  (47)

At the highest point  $\theta = 0, R = mg - \frac{mu^2}{a}$  (48)

At the point A,  $\theta = \pi/2, R = -\frac{mu^2}{a} - 2mg$  (49)

If  $u^2 > ag$ , the particle leaves contact with the wire immediately and describes a parabolic path.

If  $u^2 < ga$ , the particle remains in contact for some distance, but leaves contact when  $R$  vanishes i.e. before it reaches A and then it describes a parabolic path.

(c) If the particle moves on the inside of rough vertical circular wire, then there is an additional frictional force  $\mu R$  along the tangent opposing the motion. As such equations (42) and (43) are modified to

$$ma\theta'' = -mg \sin \theta - \mu R \quad (50)$$

$$ma\theta'^2 = -mg \cos \theta + R \quad (51)$$

Eliminating  $R$  between these equations, we get a non-linear differential equation

$$a\theta'' = -g \sin \theta - \mu(-g \cos \theta - a\theta'^2) \quad (52)$$

which can be integrated by substituting  $\theta' = w, \theta'' = w dw/d\theta$ .

Similarly (44) and (45) are modified to

$$ma\theta'' = mg \sin \theta - \mu R \quad (53)$$

$$ma\theta'^2 = -R + mg \cos \theta \quad (54)$$

We can again eliminate  $R$ , solve for  $\theta'$  and  $\theta$  and find the value of  $\theta$  when  $R$  vanishes.





**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

### UNIT III(Continuation)

Mathematical Modeling through Ordinary Differential Equations of Second Order:  
 Planetary Motions – Circular Motion and Motion of Satellites – Mathematical  
 Modeling through Linear Differential Equations of Second Order – Miscellaneous  
 Mathematical Models.

### SUGGESTED READINGS

#### TEXT BOOK

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New  
 Delhi.

#### REFERENCES

**R2:**Brain Albright, (2010). Mathematical Modeling with Excel, Jones and Bartlett  
 Publishers, New Delhi.

**R3:**Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in  
 Mathematical Modelling, Vikash Publishing House, UK.

## Circular Motion of Satellites

Just as planets move in elliptic orbits with the Sun in one focus, the man-

made artificial satellites move in elliptic (or circular) orbits with the Earth (or rather its centre) at one focus.

If the Earth is of mass  $M$  and radius  $a$  and a satellite of mass  $m$  ( $\ll M$ ) is projected from a point  $P$  at a height  $h$  above the Earth with velocity  $V$  at right angles to  $OP$  (Figure 4.6) it will move under a central force  $GmM/r^2$ . Since the central force of a circular orbits is  $mV^2/r$ , we get, if the path is to be circular,




Figure 4.6

$$\frac{mV^2}{a+h} = \frac{GmM}{(a+h)^2} \quad \text{or} \quad V^2 = \frac{GM}{a+h} \quad (55)$$

If  $g$  is the acceleration due to gravity, then the gravitational force on a particle of mass  $m$  on the surface of the Earth is  $mg$ . Alternatively from Newton's inverse square law, it is  $GMm/a^2$  so that

$$\frac{GMm}{a^2} = mg \quad \text{or} \quad Gm = ga^2 \quad (56)$$

From (55) and (56)

$$V^2 = \frac{ga^2}{a+h} \quad (57)$$

This gives the velocity of a satellite describing a circular orbit at a height  $h$  above the surface of the Earth. Its time period is given by

$$T = \frac{2\pi(a+h)}{V} = \frac{2\pi(a+h)}{\sqrt{ga}} (a+h)^{1/2} = \frac{2\pi}{\sqrt{ga}} (a+h)^{3/2} \quad (58)$$

The earth completes one revolution about its axis in twenty-four hours. As such if  $T$  is 24 hours, the satellite would have the same period as the Earth and would appear stationary, to an observer on the Earth. Now taking  $g = 32 \text{ ft/sec}^2$ ,  $a = 4000 \text{ miles}$ ,  $T = 24 \text{ hours}$ , we get if  $h$  is measured in miles

$$((4000 + h) \times 1760 \times 3)^{3/2} = \frac{24 \times 60 \times 60 \sqrt{32 \times 4000 \times 1760 \times 3 \times 7}}{2 \times 22}$$

$$= 1642607.416 \times 10^6$$

$$(4000 + h) \times 5280 = 13919.3408 \times 10^4$$

$$4000 + h = 26.36238788 \times 10^3 = 26362.38788$$

$$h = 22362.38788 \text{ miles}$$

This gives the height of the synchronous or synchron satellite, which is very useful for communication purposes.

**Elliptic Motion of Satellites**

If a satellite is projected at a height  $a + h$  above the centre of the Earth with a velocity different from  $\sqrt{ga}/\sqrt{a+h}$  or if it is not projected at right angles to the radius vector, the orbit will not be circular, but can be elliptic, parabolic or hyperbolic depending on  $V$  and the angle of projection.

If the angle of projection is  $90^\circ$  and the orbit is an elliptic with semi major axis  $a'$  and eccentricity  $e$ , then there are two possibilities depending on whether the point of projection is the apogee or the perigee

Using equation (23)




Figure 4.7

$$V^2 = \mu \left( \frac{2}{a'(1+e)} - \frac{1}{a'} \right), \quad a'(1+e) = a+h \quad (59)$$

or 
$$V^2 = \mu \left( \frac{2}{a'(1-e)} - \frac{1}{a'} \right), \quad a'(1-e) = a+h \quad (60)$$

i.e. 
$$V^2 = \frac{ga^2}{a+h} (1-e) \quad \text{or} \quad V^2 = \frac{ga^2}{a+h} (1+e)$$

i.e. 
$$V^2 = V_0^2(1-e) \quad \text{or} \quad V^2 = V_0^2(1+e), \quad (61)$$

where  $V_0$  is the velocity required for a circular orbit for which  $e = 0$ . Thus if  $V > V_0$ , the point of projection is nearest point of the orbit to the centre of the Earth and if  $V < V_0$ , this point is the furthest point.

For the elliptic orbit, the time period is

$$T = \frac{2\pi}{\sqrt{ga}} a'^{3/2} \quad (62)$$

where if  $V < V_0$ ,  $e = \sqrt{1 - \frac{V^2}{V_0^2}}$ ,  $a' = \frac{a+h}{1 + \sqrt{1 - V^2/V_0^2}}$  (63)

and if  $V > V_0$ ,  $e = \sqrt{\frac{V^2}{V_0^2} - 1}$ ,  $a' = \frac{a+h}{1 - \sqrt{V^2/V_0^2 - 1}}$  (64)



If  $h_{\max}$  and  $h_{\min}$  are the maximum and minimum heights of a satellite above the Earth's surface and  $a$  is the radius of the Earth, we get

$$\begin{aligned} \frac{a'(1+e)}{a'(1-e)} &= \frac{a+h_{\max}}{a+h_{\min}} \text{ or } \frac{1+e}{a+h_{\max}} = \frac{1-e}{a+h_{\min}} \\ &= \frac{2}{2a+h_{\max}+h_{\min}} \\ \text{or } \frac{1+e}{a+h_{\max}} &= \frac{1}{a+\frac{h_{\max}+h_{\min}}{2}} = \frac{e}{\frac{h_{\max}-h_{\min}}{2}} \\ \text{or } e &= \frac{h_{\max}-h_{\min}}{2a+h_{\max}-h_{\min}} \end{aligned} \quad (65)$$

## MATHEMATICAL MODELLING THROUGH LINEAR DIFFERENTIAL EQUATIONS OF SECOND ORDER

### Rectilinear Motion

Let one end  $O$  of an elastic string of natural length  $L(=OA)$  be fixed (Figure 4.8) and let the other end to which a particle of mass  $m$  is attached



Figure 4.8

be stretched a distance  $a$  and then released. At any time  $t$ , let  $x(t)$  be the extension, then the equation of motion of the particle is

$$m \frac{d^2x}{dt^2} = -\lambda \frac{x}{L} = -kx, \quad (66)$$

where  $k$  is the elastic constant. If the particle moves in a resisting medium with resistance proportional to the velocity  $x'$ , (66) becomes

$$mx'' + cx' + kx = 0, \quad (67)$$

which is a linear differential equation of the second order. Its solution is

$$x(t) = A_1 e^{\lambda_1 t} + A_2 e^{\lambda_2 t} \quad (68)$$

where  $\lambda_1, \lambda_2$  are the roots of

$$m\lambda^2 + c\lambda + k = 0 \quad (69)$$

Here  $\lambda_1 + \lambda_2 = -\frac{c}{m}$ ,  $\lambda_1 \lambda_2 = \frac{k}{m}$ . The sum of the roots is negative and the product of the roots is positive.

Case (i)  $c^2 > 4km$ , the roots are real and distinct and are negative. As such  $x(t) \rightarrow 0$  as  $t \rightarrow \infty$ . The motion is said to be *overdamped*.

Case (ii)  $c^2 = 4km$ , the roots are real and equal and

$$x(t) = (A_1 + A_2 t) \exp\left(-\frac{c}{2m} t\right) \quad (70)$$

and again  $x(t) \rightarrow 0$  as  $t \rightarrow \infty$ . In this case the motion is said to be *critically damped*.

Case (iii)  $c^2 < 4km$ , the roots are complex conjugate with the real parts of the roots negative.  $x(t)$  always oscillates but oscillations are damped out and tend to zero. In this case, the motion is said to be *under damped*.

Next we consider the case when there is an external force  $m \cdot F(t)$  acting on the particle. In this case (67) becomes

$$mx'' + cx' + kx = mF(t) \quad (71)$$

A particular case of interest is given by the model

$$x'' + w_0^2 x = F \cos wt \quad (72)$$

i.e., when in the absence of the external force, the motion is simple harmonic with period  $2\pi/w_0$  and the external force is periodic with period  $2\pi/w$ . The solution of (72) is given by

$$x(t) = A \cos(w_0 t - \alpha) + F \cos wt / (w_0^2 - w^2) \quad w \neq w_0 \quad (73)$$

$$= A \cos(w_0 t - \alpha) + \frac{F}{2w_0} t \sin w_0 t \quad w = w_0 \quad (74)$$

When  $w = w_0$ , the first term is periodic and its amplitude never exceeds  $|A|$ . However as  $t \rightarrow \infty$  along a sequence for which  $\sin w_0 t = \pm 1$ , the magnitude of the second term approaches infinity.

The phenomenon we have discussed here is known as of *pure* or *undamped resonance*. It occurs when  $c = 0$  and the input and natural frequencies are equal. We shall get a similar phenomenon when  $c$  is small. The forcing function  $F \cos wt$  is then said to be in resonance with the system.

Bridges, cars, planes, ships are vibrating systems and an external periodic force with the same frequency as their natural frequency can damage them. This is the reason why soldiers crossing a bridge are not allowed to march in step. However resonance phenomenon can also be used to advantage e.g. in uprooting trees or in getting a car out of a ditch.

When  $w$  and  $w_0$  differ only slightly, the solution represents superposition of two sinusoidal waves whose periods differ only slightly and this leads to the occurrence of beats.



### Electrical Circuits

Figure 4.9 shows an electrical circuit. The current  $i(t)$  amperes represents the time rate of change of charge  $q$  flowing in the circuits, so that

$$\frac{dq}{dt} = i(t) \quad (75)$$

(i) There is a resistance of  $R$  Ohms in the circuit. This may be provided by a light bulb, an electric heater or any other electrical device opposing the motion of the charge and causing a potential drop of magnitude  $E_R = Ri$  volts.




Figure 4.9

(ii) There is an induction of inductance  $L$  henrys which produces a potential drop  $E_L = L di/dt$ .

(iii) There is a capacitance  $C$  which produces a potential drop

$$E_c = \frac{1}{C} q.$$

All these potential drops are balanced by the battery which produces a voltage  $E$  volts. Now according to Kirchhoff's second law, the algebraic sum of the voltage drops round a closed circuit is zero so that

$$Ri + L \frac{di}{dt} + \frac{1}{C} q = E(t) \quad (76)$$

Differentiating and using (75), we get

$$L \frac{d^2i}{dt^2} + R \frac{di}{dt} + \frac{1}{C} i = \frac{dE}{dt} \quad (77)$$



Also substituting for (75) in (76) we get

$$L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{1}{C} q = E(t) \quad (78)$$

Both (77) and (78) represent linear differential equations with constant coefficients and their solutions will determine  $i(t)$  and  $q(t)$ .

Comparing (71) and (78), we get the correspondences

$$\begin{aligned} \text{mass } m &\leftrightarrow \text{inductance } L \\ \text{friction coefficient } c &\leftrightarrow \text{resistance } R \\ \text{spring constant } k &\leftrightarrow \text{inverse capacitance } 1/C \\ \text{impressed force } F &\leftrightarrow \text{impressed voltage } E \\ \text{displacement } x &\leftrightarrow \text{charge } q \\ \text{velocity } v = dx/dt &\leftrightarrow \text{current } i = \frac{dq}{dt} \end{aligned}$$

This shows the correspondence between mechanical and electrical systems. This forms the basis of analogue computers. A linear differential equation of the second order can be solved by forming an electrical circuit and measuring the electric current in it. Similar analogues exist between hydrodynamical and electrical systems. Mathematical modelling brings out the isomorphisms between mathematical structures of quite different systems and gives a method for solving all these models in terms of the simplest of these models.

We can have analogues of (71), (78) in economic system when  $k(t)$  represents the excess of the capital invested over the equilibrium capital and  $E(t)$  can represent external investments.

### Phillip's Stabilization Model for a Closed Economy

The assumptions of the model are:

- (i) The producers adjust the national production  $Y$  of a product according to the aggregate demand  $D$ . If  $D > Y$ , they increase production and if  $D < Y$ , they decrease production so that we get

$$dY/dt = \alpha(D - Y), \alpha > 0, \quad (79)$$

where  $\alpha$  is a reaction coefficient representing the velocity of adjustment.

(ii) Aggregate demand  $D$  is the sum of private demand, government demand  $G$  and an exogenous disturbance  $u$ . The private demand is proportional to the national income or output so that

$$D = (1 - L) Y + G - u \quad (80)$$

where  $1 - L$  is the marginal propensity to spend i.e. it is the marginal propensity to consume plus the marginal propensity to invest. We assume that  $0 < L < 1$ .

(iii) The government adjusts its demand to bring the national out-put to a desired level, which without loss of generality may be taken as zero.

The Government decides its demand according to one of the following policies:

(a) *proportionate stabilization policy* according to which

$$G^* = -f_p Y \quad (81)$$

where  $f_p > 0$  is the coefficient of proportionality and we use the negative sign on the right hand side since if the output is less than the described level, government will come out with a positive demand.

(b) *derivative stabilization policy* according to which

$$G^* = -f_d Y', \quad (82)$$

where  $f_d > 0$  and the government demand is proportional to  $Y'$ .

(c) *mixed proportionate derivative policy* according to which

$$G^* = -f_p Y - f_d Y' \quad (83)$$

(d) *integral stabilization policy* according to which

$$G^* = -f_I \int_0^t Y dt, \quad f_I > 0 \quad (84)$$

(iv)  $G^*$  is the potential demand which the Government may like to make, but the actual demand  $G$  will be gradually adjusted so that

$$G' = \beta(G^* - G), \quad (85)$$

where  $\beta$  is the reaction coefficient.  $\beta > 0$  since if  $G < G^*$ , the government tends to increase the demand to reach  $G^*$ .

Now from (79) and (80)

$$dY/dt = \alpha((1 - L) Y + G - u - Y), \quad (86)$$

so that

$$d^2 Y/dt^2 = -\alpha L dY/dt + \alpha dG/dt \quad (87)$$



Eliminating  $G$  between (85), (86) and (87)

$$\frac{d^2 Y/dt^2}{\alpha} + L \frac{dY}{dt} = \beta \left( G^* - \frac{dY/dt}{\alpha} - (Ly + u) \right) \quad (88)$$

or 
$$d^2 Y/dt^2 + dY/dt (\alpha L + \beta) + \alpha \beta L Y + \alpha \beta u = \alpha \beta G^* \quad (89)$$

If we substitute for  $G^*$  from (81), (82) or (83), we get a linear differential equation of the second order with constant coefficients. If however the government uses integral stabilization policy, we use (84) to get the third order differential equation

$$d^3 Y/dt^3 + (\alpha L + \beta) d^2 Y/dt^2 + \alpha \beta dY/dt + \alpha \beta f_I Y = 0 \quad (90)$$

The equations (89) and (90) can be easily solved. Even without solving these, the stability of the solutions and their behaviour as  $t \rightarrow \infty$  can be easily obtained.

## MISCELLANEOUS MATHEMATICAL MODELS THROUGH ORDINARY DIFFERENTIAL EQUATIONS OF THE SECOND ORDER

### The Catenary

A perfectly inflexible string is suspended under gravity from two fixed points  $A$  and  $B$  (Fig. 4.10).




Figure 4.10



Consider the equilibrium of the part  $CD$  of the string of length  $s$  where  $C$  is the lowest point of the string at which the tangent is horizontal.

The forces acting on this part of the string are (i) tension  $T_0$  at  $C$  (ii) tension  $T$  at point  $D$  along tangent at  $D$  (iii) weight  $ws$  of the string.

Equating the horizontal and vertical components of forces, we get

$$T \cos \psi = T_0, \quad T \sin \psi = ws \quad (91)$$

Let  $T_0$  be equal to weight of length  $c$  of the string, then (91) give

$$\tan \psi = \frac{ws}{T_0} = \frac{ws}{wc} = \frac{s}{c} \quad (92)$$

$$\frac{ds}{d\psi} = \rho = c \sec^2 \psi, \quad (93)$$

where  $\rho$  is radius of curvature of the string at  $D$ ; so that

$$\frac{\left(1 + \left(\frac{dy}{dx}\right)^2\right)^{3/2}}{\frac{d^2y}{dx^2}} = c \left(1 + \left(\frac{dy}{dx}\right)^2\right)$$

or 
$$c \left(\frac{d^2y}{dx^2}\right) = \left(1 + \left(\frac{dy}{dx}\right)^2\right)^{1/2}, \quad (94)$$

which is a non-linear differential equation of second order. If  $\frac{dy}{dx} = p$ , then (94) gives

$$c \frac{dp}{\sqrt{1+p^2}} = dx \quad (95)$$

Integrating 
$$\sinh^{-1} p = \frac{x}{c} + A \quad (96)$$

When  $x = 0$ ,  $p = 0$ , so that  $A = 0$  and

$$\frac{dy}{dx} = \sinh \frac{x}{c} \quad (97)$$

Integrating

$$y = c \cosh \frac{x}{c}, \quad (98)$$

where we choose  $x$ -axis in such a way that  $y = c$  when  $x = 0$ . This is the equation of the common catenary.

It may be noted that here we get a differential equation of the second order from a problem of statics rather than from a problem of dynamics.

**A Curve of Pursuit**

A ship at the point  $(a, 0)$  sights a ship at  $(0, 0)$  moving along  $y$ -axis with a uniform velocity  $ku$  ( $0 < k < 1$ ). It begins to pursue ship  $B$  with a velocity  $u$  always moving in the direction of the ship  $B$  so that at any time  $AB$  is along the tangent to the path of  $A$ .

From Figure 4.11


$$\tan(\pi - \psi) = \frac{kut - y}{x}$$

or  $-\frac{dy}{dx} = -\frac{y}{x} + \frac{kut}{x}$

or  $x \frac{dy}{dx} - y = -kut$  (99)

Differentiating with respect to  $x$ , we get

$$x \frac{d^2y}{dx^2} = -ku \frac{dt}{dx} \quad (100)$$



**Figure 4.11**

Now  $dx/dt$  = horizontal component of velocity of  $A = u \cos(\pi - \phi)$

$$= -u \cos \phi = -\frac{u}{\sqrt{1 + \left(\frac{dy}{dx}\right)^2}} \quad (101)$$

so that from (99) and (100)

$$x \frac{d^2y}{dx^2} = k \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \quad (102)$$

Putting,  $\frac{dy}{dx} = p$ , we get

$$\frac{dp}{\sqrt{1 + p^2}} = k \frac{dx}{x} \quad (103)$$

Integrating  $\frac{dy}{dx} = k \left( \sinh^{-1} \left( \ln \frac{x}{a} \right) \right) \quad (104)$

Integrating once again, we get  $y$  as a function of  $x$ .



**POSSIBLE QUESTIONS****Part B (6 Marks)**

1. Derive the components of velocity and acceleration vectors along radial and transverse directions.
2. Find the height of synchronous from the circular motion of satellites.
3. Explain about the catenary.
4. Design a mathematical model for motion of a projectile.
5. Explain on elliptic motion of satellites.
6. Discussion detail on a curve of pursuit.
7. Discuss motion of a particle on a rough vertical wire.

**Part C (10 Marks)**

1. Explain in detail Kepler's law of planetary motion.
2. Explain on circular motion of satellites.
3. Discuss motion of a particle on a rough vertical wire.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

---

## **UNIT IV**

Mathematical Modeling through Difference Equations : Simple Models – Basic Theory of Linear Difference Equations with Constant Coefficients – Economics and Finance – Population Dynamics and Genetics – Probability Theory.

## **SUGGESTED READINGS**

### **TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

### **REFERENCES**

**R3:** Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.

## THE NEED FOR MATHEMATICAL MODELLING THROUGH DIFFERENCE EQUATIONS: SOME SIMPLE MODELS

We need difference equation models when either the independent variable is discrete or it is mathematically convenient to treat it as a discrete variable.

Thus in Genetics, the genetic characteristics change from generation to generation and the variable representing a generation is a discrete variable.

In Economics, the price changes are considered from year to year or from month to month or from week to week or from day to day. In every case, the time variable is discretized.

In Population Dynamics, we consider the changes in population from one age-group to another and the variable representing the age-group is a discrete variable.

In finding the probability of  $n$  persons in a queue or the probability of  $n$  persons in a state or the probability of  $n$  successes in a certain number of trials, the independent variable is discrete.

For mathematical modelling through differential equations, we give an increment  $\Delta x$  to independent variable  $x$ , find the change  $\Delta y$  in  $y$  and let  $\Delta x \rightarrow 0$  to get differential equations. In most cases, we cannot justify the limiting process rigorously. Thus for modelling fluid motion, making  $\Delta x \rightarrow 0$  has no meaning since a fluid consists of a large number of particles and the distance between two neighbouring particles cannot be made arbitrary small. Continuum mechanics is only an approximation (though fortunately a very good one) to reality.

Even if the limiting process can be justified e.g. when the independent variable is time, the resulting differential equation may not be solvable analytically. We then solve it numerically and for this purpose, we again replace the differential equation by a system of difference equations. Numerical methods of solving differential equations essentially mean solving difference equations.



We now give simple difference equation models parallel to the differential equation models studied in earlier chapters.

(i) *Population Growth Model*: If the population at time  $t$  is  $x(t)$ , then assuming that the number of births and deaths in the next unit interval of time are proportional to the populations at time  $t$ , we get the model:

$$x(t+1) - x(t) = bx(t) - dx(t) \quad \text{or} \quad x(t+1) = ax(t), \quad (1)$$

so that

$$x(t) = ax(t-1) = a^2x(t-2) = a^3x(t-3) = \dots = a^tx(0) \quad (2)$$

This may be compared with the differential equation model:

$$\frac{dx}{dt} = ax \quad \text{with the solution} \quad x(t) = x(0)e^{at} \quad (3)$$

For solving the difference equation model, we require only simple algebra, but for solving the differential equation model, we require knowledge of calculus, differential equation and exponential functions.

(ii) *Logistic Growth Model*: This is given by

$$x(t+1) - x(t) = ax(t) - bx^2(t) \quad (4)$$

This is not easy to solve, but given  $x(0)$ , we can find  $x(1)$ ,  $x(2)$ ,  $x(3)$ , ... in succession and we can get a fairly good idea of the behaviour of the model with the help of a pocket calculator.

(iii) *Prey-Predator Model*: This is given by

$$\left. \begin{aligned} x(t+1) - x(t) &= -ax(t) + bx(t)y(t) \\ y(t+1) - y(t) &= py(t) - qx(t)y(t) \end{aligned} \right\} \begin{aligned} a, b &> 0 \\ p, q &> 0 \end{aligned} \quad (5)$$

and again given  $x(0)$ ,  $y(0)$ , we can find  $x(1)$ ,  $y(1)$ ;  $x(2)$ ,  $y(2)$ ;  $x(3)$ ,  $y(3)$ , ... in succession.

(iv) *Competition Model*: This is given by

$$\left. \begin{aligned} x(t+1) - x(t) &= ax(t) - bx(t)y(t) \\ y(t+1) - y(t) &= px(t) - qx(t)y(t) \end{aligned} \right\} \begin{aligned} a, b &> 0 \\ p, q &> 0 \end{aligned} \quad (6)$$

(v) *Simple Epidemics Model*: This is given by

$$\left. \begin{aligned} x(t+1) - x(t) &= -\beta x(t)y(t) \\ y(t+1) - y(t) &= \beta x(t)y(t) \end{aligned} \right\}, \quad \beta > 0 \quad (7)$$

## The Complementary Function

We try the solution  $x_t = a^t$ . If this satisfies (11), we get

$$g(\lambda) = a_0\lambda^n + a_1\lambda^{n-1} + a_2\lambda^{n-2} + \dots + a_n = 0 \quad (13)$$

This algebraic equation of  $n$ th degree has  $n$  roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , real or complex. The complementary function is then given by

$$G_1(t) = c_1 \lambda_1^t + c_2 \lambda_2^t + \dots + c_n \lambda_n^t \quad (14)$$

**Case (i):** If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are all real and distinct, (14) gives us the complementary function when  $c_1, c_2, \dots, c_n$  are any  $n$  arbitrary real constants.

**Case (ii):** If two of the roots  $\lambda_1, \lambda_2$  are equal, then (14) contains only  $n - 1$  arbitrary constants and as such it cannot be the most general solution. We try the solution  $c t \lambda_1^t$ . We get

$$a_0(t+n)\lambda_1^n + a_1(t+n-1)\lambda_1^{n-1} + \dots + a_n = 0$$

$$\text{or} \quad t g(\lambda_1) + g'(\lambda_1) = 0, \quad (15)$$

which is identically satisfied since both  $g(\lambda_1) = 0$  and  $g'(\lambda_1) = 0$  as  $\lambda_1$  is a repeated root. In this case

$$G_1(t) = (c_1 + c_2 t) \lambda_1^t + c_3 \lambda_3^t + c_4 \lambda_4^t + \dots + c_n \lambda_n^t \quad (16)$$

**Case (iii):** If a root  $\lambda_1$  is repeated  $k$  times, the complementary function is

$$G_1(t) = (c_1 + c_2 t + c_3 t^2 + \dots + c_k t^{k-1}) \lambda_1^t + c_{k+1} \lambda_{k+1}^t + \dots + c_n \lambda_n^t \quad (17)$$

**Case (iv):** Let  $g(\lambda) = 0$  have two complex roots  $\alpha \pm i\beta$ , then their contribution to complementary function is

$$c_1(\alpha + i\beta)^t + c_2(\alpha - i\beta)^t \quad (18)$$

Putting  $\alpha = r \cos \theta$ ,  $\beta = r \sin \theta$  and using De Moivre's theorem, this reduces to

$$\begin{aligned} & c_1 r^t (\cos \theta + i \sin \theta)^t + c_2 r^t (\cos \theta - i \sin \theta)^t \\ &= r^t \cos(\theta t) (c_1 + c_2) + r^t \sin(\theta t) (i c_1 - i c_2) \\ &= r^t (d_1 \cos(\theta t) + d_2 \sin(\theta t)) \\ &= (x^2 + \beta^2)^{t/2} (d_1 \cos(\theta t) + d_2 \sin(\theta t)), \end{aligned} \quad (19)$$

$$\text{where } \tan \theta = \frac{\beta}{\alpha} \quad (20)$$

and  $d_1, d_2$  are arbitrary constants.

**Case (v):** If the complex roots  $\alpha \pm i\beta$  are repeated  $k$  times, then contribution to the complementary function is

$$\begin{aligned} & (x^2 + \beta^2)^{t/2} ((d_0 + d_1 t + \dots + d_{k-1} t^{k-1}) \cos(\theta t) \\ &+ (f_0 + f_1 t + \dots + f_{k-1} t^{k-1}) \sin(\theta t)) \end{aligned} \quad (21)$$

where  $d_0, d_1, \dots, d_{k-1}, f_0, \dots, f_{k-1}$  are  $2k$  arbitrary constants.

## The Particular Solution



Here we want a solution of (10) not containing any arbitrary constant.  
**Case (i):** Let  $\varphi(t) = AB^t$ ,  $B$  is not a root of  $g(\lambda) = 0$  (22)

We try the solution  $CB^t$ . Substituting in (10), we get

$$CB^t(a_0B^n + a_1B^{n-1} + \dots + a_n) = AB^t \quad (23)$$

If  $B \neq \lambda_1, \lambda_2, \dots, \lambda_n$ , we get

$$C = \frac{A}{a_0B^n + a_1B^{n-1} + \dots + a_n} \quad (24)$$

and the particular solution is

$$\frac{AB^t}{a_0B^n + a_1B^{n-1} + \dots + a_n} \quad (25)$$

**Case (ii):** Let

$$\varphi(t) = AB^t, B \text{ is a non-repeated root of } g(\lambda) = 0 \quad (26)$$

We try the solution  $CtB^t$ . Substituting in (10), we get

$$B^t(Ct g(B) + Cg'(B)) = AB^t \quad (27)$$

Since  $g(B) = 0$ ,  $g'(B) \neq 0$

$$C = \frac{A}{g'(B)}, \quad (28)$$

so that the particular solution is

$$\frac{AtB^t}{a_0nB^{n-1} + a_1(n-1)B^{n-2} + \dots + a_{n-1}} \quad (29)$$

**Case (iii):** Let

$$\varphi(t) = AB^t, \quad g(B) = 0, \quad g'(B) = 0, \dots,$$

$$g^{(k-1)}(B) = 0, \quad g^{(k)}(B) \neq 0, \quad (30)$$

then the particular solution is

$$\frac{At^{k-1}B^t}{g^{(k)}(B)} \quad (31)$$

**Case (iv):** Let

$$\varphi(t) = At^k \quad (32)$$

We try the solution

$$d_0t^k + d_1t^{k-1} + d_2t^{k-2} + \dots + d_k \quad (33)$$

Substituting in (10) we get

$$\begin{aligned} & a_0(d_0(t+n)^k + d_1(t+n)^{k-1} + d_2(t+n)^{k-2} + \dots + d_k) \\ & + a_1(d_0(t+n-1)^k + d_1(t+n-1)^{k-1} + d_2(t+n-1)^{k-2} \\ & + \dots + d_k) + \dots + a_n(d_0t^k + d_1t^{k-1} + d_2t^{k-2} + \dots + d_k) \\ & = 0 \end{aligned} \quad (34)$$



Equating the coefficients of  $t^k, t^{k-1}, \dots, t^0$ , on both sides, we get  $(k+1)$  equations which in general will enable us to determine  $d_0, d_1, d_2, \dots, d_k$  and thus the particular solution will be determined.

$$\begin{aligned}x_1(t+1) &= x_2(t) \\x_2(t+1) &= x_3(t) \\\dots &\dots \\x_{n-1}(t+1) &= x_n(t) \\x_n(t+1) &= -\frac{a_1}{a_0}x_n(t) - \frac{a_2}{a_0}x_{n-1}(t) - \dots - \frac{a_n}{a_0}x_1(t),\end{aligned}\tag{37}$$

which can be written in the matrix form

$$\begin{bmatrix} x_1(t+1) \\ x_2(t+1) \\ \vdots \\ x_n(t+1) \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\frac{a_n}{a_0} & -\frac{a_{n-1}}{a_0} & -\frac{a_{n-2}}{a_0} & \dots & -\frac{a_1}{a_0} \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix}\tag{38}$$

$$\text{or } X(t+1) = AX(t),\tag{39}$$

where  $X(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix}$ ,

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -\frac{a_n}{a_0} & -\frac{a_{n-1}}{a_0} & -\frac{a_{n-2}}{a_0} & \cdots & -\frac{a_1}{a_0} \end{bmatrix} \quad (40)$$

Applying (39) repeatedly

$$X(k) = A^k X(0), \quad (41)$$

where

$$X(0) = \begin{bmatrix} x_1(0) \\ x_2(0) \\ x_3(0) \\ \vdots \\ x_n(0) \end{bmatrix} = \begin{bmatrix} x_1(0) \\ x_1(1) \\ x_1(2) \\ \vdots \\ x_1(n-1) \end{bmatrix} = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_{n-1} \end{bmatrix} \quad (42)$$

Thus knowing the values of  $x_1$  at times  $0, 1, 2, \dots, n-1$ , we can find its value at all subsequent times.

### Solution of Linear Difference Equations by Using z-Transform

Let  $\{u_n\}$  be an infinite sequence, then its z-transform is defined by

$$Z(u_n) = \sum_{n=0}^{\infty} u_n z^{-n}, \quad (51)$$

whenever this infinite series converges. If  $\{u_n\}$  is a probability distribution and  $z = 1/s$ , it will be the same as the probability generating function.

The following results can be easily established

(i) If  $k > 0$ ,  $Z(u_{n-k}) = z^{-k} Z(u_n)$  (52)

(ii) If  $k > 0$ ,  $Z(u_{n+k}) = z^k [Z(u_n) - \sum_{m=0}^{k-1} u_m z^{-m}]$  (53)

(iii)  $u_n$ :  $1, a^n, e^{an}$   
 $Z(u_n)$ :  $z/(z-1), z/(z-a), z/(z-e^a)$  (54)

Taking z-transform of both sides of a linear difference equation, we can find  $Z(u_n)$  and expanding it in powers of  $1/z$  and finding the coefficient of  $z^{-n}$ , we can get  $u_n$ .



## The Cobweb Model

Let  $p_t$  = price of a commodity in the year  $t$  and

$q_t$  = amount of the commodity available in the market in year  $t$ ,

then we make the following assumptions

(i) Amount of the commodity produced this year and available for sale is a linear function of the price of the commodity in the last year, i.e.

$$q_t = \alpha + \beta p_{t-1}, \quad (68)$$

where  $\beta > 0$  since if the last year's price was high, the amount available this year would also be high.

(ii) The price of the commodity this year is a linear function of the amount available this year i.e.

$$p_t = \gamma + \delta q_t, \quad (69)$$

where  $\delta < 0$ , since if  $q_t$  is large, the price would be low. From (68) and (69)

$$p_t - \beta\delta p_{t-1} = \gamma + \alpha\delta, \quad (70)$$

which has the solution

$$\left(p_t - \frac{\alpha\delta + \gamma}{1 - \beta\delta}\right) = \left(p_0 - \frac{\alpha\delta + \gamma}{1 - \beta\delta}\right)(\beta\delta)^t, \quad (71)$$

so that

$$\left(p_t - \frac{\alpha\delta + \gamma}{1 - \beta\delta}\right) = \left(p_{t-1} - \frac{\alpha\delta + \gamma}{1 - \beta\delta}\right)(\beta\delta) \quad (72)$$

Since  $\beta\delta$  is negative  $p_0, p_1, p_2, p_3, \dots$  are alternatively greater and less than  $(\alpha\delta + \gamma)/(1 - \beta\delta)$ .

If  $|\beta\delta| > 1$ , the deviation of  $p_t$  from  $(\alpha\delta + \gamma)/(1 - \beta\delta)$  goes on increasing. On the other hand if  $|\beta\delta| < 1$ , this deviation goes on decreasing and ultimately  $p_t \rightarrow (\alpha\delta + \gamma)/(1 - \beta\delta)$  as  $t \rightarrow \infty$ .

Figures 5.1a and 5.1b show how the price approaches the equilibrium price  $p_e = (\alpha\delta + \gamma)/(1 - \beta\delta)$  as  $t$  increases in the two cases when  $p_0 > p_e$  and  $p_0 < p_e$  respectively.






Figure 5.1

In the same way, eliminating  $p_t$  from (67), (68) we get

$$q_t = \alpha + \beta\gamma + \beta\delta q_{t-1}, \quad (73)$$

which has the solution

$$\left(q_t - \frac{\alpha + \beta\gamma}{1 - \beta\delta}\right) = \left(q_e - \frac{\alpha + \beta\gamma}{1 - \beta\delta}\right)(\beta\delta)^t, \quad (74)$$

so that  $q_t$  also oscillates about the equilibrium quantity level

$$q_t = (\alpha + \beta\gamma)/(1 - \beta\delta) \quad \text{if} \quad |\beta\delta| < 1$$

The variation of both prices and quantities is shown simultaneously in Figure 5.2.

Suppose we start in the year zero with price  $p_0$ , and quantity  $q_0$  represented by the point  $A$ . In year 1, the quantity  $q_1$  is given by  $\alpha + \beta p_0$  and the price is given by  $p_1 = \gamma + \delta q_1$ . This brings us to the point  $C$  in two steps via  $B$ . The path of prices and quantities is thus given by the Cobweb path  $ABCDEFGHI, \dots$  and the equilibrium price and quantity are given by the intersection of the two straight lines.

## Application to Actuarial Science

One important aspect of actuarial science is what is called mathematics of finance or mathematics of investment.

If a sum  $S_0$  is invested at compound interest of  $i$  per unit amount per unit time and  $S_t$  is the amount at the end of time  $t$ , then we get the difference equation

$$S_{t+1} = S_t + iS_t = (1 + i)S_t, \quad (78)$$

which has the solution

$$S_t = S_0(1 + i)^t, \quad (79)$$

which is the well-known formula for compound interest.

Suppose a person borrows a sum  $S_0$  at compound interest  $i$  and wants to amortize his debt, i.e. he wants to pay the amount and interest back by payment of  $n$  equal instalments, say  $R$ , the first payment to be made at the end of the first year.

Let  $S_t$  be the amount due at the end of  $t$  years, then we have the difference equation

$$S_{t+1} = S_t + iS_t - R = (1 + i)S_t - R \quad (80)$$

Its solution is

$$S_t = \left(S_0 - \frac{R}{i}\right)(1 + i)^t + \frac{R}{i} \quad (81)$$

$$= S_0(1 + i)^t - R \frac{(1 + i)^t - 1}{i} \quad (82)$$

If the amount is paid back in  $n$  years,  $S_n = 0$ , so that



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**(Deemed to be University Established Under Section 3 of UGC Act 1956)**  
**Pollachi Main Road, Eachanari (Po),**  
**Coimbatore –641 021**  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

---

**UNIT IV(cont)**

Mathematical Modeling through Difference Equations : Simple Models – Basic Theory of Linear Difference Equations with Constant Coefficients – Economics and Finance – Population Dynamics and Genetics – Probability Theory.

**SUGGESTED READINGS**

**TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

**REFERENCES**

**R3:** Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.



$$R = S_0 \frac{i}{1 - (1 + i)^{-n}} = S_0 \frac{1}{a_{\overline{n}|i}}, \quad (83)$$

where  $a_{\overline{n}|i}$  called the amortization factor is the present value of an annuity of 1 per unit time for  $n$  periods at an interest rate  $i$ .

The functions  $a_{\overline{n}|i}$  and  $(a_{\overline{n}|i})^{-1}$  are tabulated for common values of  $n$  and  $i$ .

Suppose an amount  $R$  is deposited at the end of every period in a bank and let  $S_t$  be the amount at the end of  $t$  periods, then

$$S_{t+1} = S_t(1 + i) + R, \quad (84)$$

so that (since  $S_0 = 0$ )

$$S_n = R \frac{(1 + i)^n - 1}{i} = RS_{\overline{n}|i} \quad (85)$$

From (83) and (85)

$$S_{\overline{n}|i} = (1 + i)^n a_{\overline{n}|i} \quad (86)$$

or

$$\frac{1}{S_{\overline{n}|i}} = \frac{(1 + i)^{-n}}{a_{\overline{n}|i}} \quad (87)$$

If a person has to pay an amount  $S$  at the end of  $n$  years, he can do it by paying into a sinking fund an amount  $R$  per period where

$$R = S \frac{1}{S_{\overline{n}|i}} \quad (88)$$

where  $\frac{1}{S_{\overline{n}|i}}$  is the sinking fund factor and can be tabulated by using (87).

## MATHEMATICAL MODELLING THROUGH DIFFERENCE EQUATIONS IN POPULATION DYNAMICS AND GENETICS

### Non-Linear Difference Equations Model for Population Growth: Non-Linear Difference Equations

Let  $x_t$  be the population at time  $t$  and let births and deaths in time-interval  $(t, t + 1)$  be proportional to  $x_t$ , then the population  $x_{t+1}$  at time  $t + 1$  is given by

$$x_{t+1} = x_t + bx_t - dx_t = x_t(1 + a) \quad (89)$$

This has the solution

$$x_t = x_0(1 + a)^t, \quad (90)$$



so that the population increases or decreases exponentially according as  $a > 0$  or  $a < 0$ . We now consider the generalisation when births and deaths  $b$  and  $d$  per unit population depend linearly on  $x_t$  so that

$$\begin{aligned} x_{t+1} &= x_t + (b_0 - b_1 x_t)x_t - (d_0 + d_1 x_t)x_t \\ &= mx_t - rx_t^2 = mx_t \left(1 - \frac{r}{m} x_t\right) \end{aligned} \quad (91)$$

This is the simplest non-linear generalisation of (90) and gives the discrete version of the logistic law of population growth. However this model shows many new features not present in the continuous version of the logistic model. Let  $rx_t/m = y_t$ , then (91) becomes

$$y_{t+1} = my_t(1 - y_t) \quad (92)$$

#### One-Period Fixed Points and Their Stability

A one-period fixed point of this equation is that value of  $y_t$  for which  $y_{t+1} = y_t$  i.e. for which

$$y_t = my_t(1 - y_t), \quad (93)$$

so that there are two one-period fixed points 0 and  $(m-1)/m$ . If  $y_0 = 0$ , then  $y_1, y_2, y_3, \dots$  are all zero and the population remains fixed at zero value:

If  $y_0 = (m-1)/m$ , then  $y_1, y_2, y_3, \dots$  are all equal to  $(m-1)/m$ . The second fixed point exists only if  $m > 1$ .

We now discuss the stability of equilibrium of each of these equilibrium positions.

Putting  $y_t = 0 + u_t$  in (92) and neglecting squares and higher powers of  $u_t$ , we get  $u_{t+1} = mu_t$  and since  $m > 0$ , the first equilibrium position is one of unstable equilibrium.

Again putting  $y_t = (m-1)/m + u_t$  in (92) and neglecting squares and higher powers of  $u_t$ , we get

$$u_{t+1} = (2-m)u_t, \quad (94)$$

so that the second position of equilibrium is stable only if  $-1 < 2-m < 1$  or if  $1 > m-2 > -1$  or if  $1 < m < 3$ .

Thus if  $0 < m < 1$ , there is only one one-period fixed point and it is unstable. If  $1 < m < 3$ , there are two one-period fixed points, the first is unstable and the second is stable. If  $m > 3$ , there are two one-period fixed points, both of which are unstable.

#### Two-Period Fixed Points and Their Stability

A point is called a two-period fixed point if it repeats itself after two periods i.e. if  $y_{t+2} = y_t$  i.e. if

$$y_{t+2} = my_{t+1}(1 - y_{t+1}) = m^2 y_t(1 - y_t)(1 - my_t + my_t^2) = y_t \quad (95)$$



or

$$y_t(m y_t - (m - 1))(m^2 y_t^2 - m(1 + m)y_t + (1 - m)) = 0 \quad (96)$$

This is a fourth degree equation and as such there can be four two-period fixed points. Two of these are the same as the one-period fixed points. This is obvious from the consideration that every one-period fixed point is also a two-period fixed point. The genuine two-period fixed points are obtained by solving the equation

$$m^2 y_t^2 - m(1 + m)y_t + (1 - m) = 0 \quad (97)$$

Its roots are real if  $m > 3$ . Thus if  $m > 3$ , the two one-period fixed points become unstable, but two new two-period fixed points exist and we can discuss their stability as before.

It can be shown that if  $m_2 < m < m_4$ , where  $m_2 = 3$  and  $m_4$  is a number slightly greater than 3, then the two two-period fixed points are stable but if  $m > m_4$ , all the four one- and two-periods become unstable, but four new four-period fixed points exist which are stable if  $m_4 < m < m_8$  and become unstable if  $m > m_8$ .

#### *2<sup>n</sup>-Period Fixed Points and Their Stability*

It can be shown that there exists an increasing infinite sequence of real numbers  $m_2, m_4, m_8, \dots, m_{2^n}, m_{2^{n+1}}, \dots$  such that when  $m_{2^n} < m < m_{2^{n+1}}$  there are  $2^{n+1}$ -period fixed points, out of which  $2^n$  fixed points are also fixed points of lower order time periods and all these are unstable and the remaining  $2^n$  points are genuine  $2^{n+1}$  period fixed points and are stable.

From 5.3 represents the stable fixed period points.




Figure 5.3

When  $m$  lies between  $m_1$  and  $m_2$ , there is one stable one-period fixed point.  
When  $m$  lies between  $m_2$  and  $m_4$  there are two stable two-period fixed points.



When  $m$  lies between  $m_4$  and  $m_8$ , there are four stable four-period fixed points, and so on.

#### *Fixed Points of other Periods*

The sequence  $m_2, m_4, m_8, \dots$  is bounded above by a fixed number  $m^*$ . If  $m > m^*$ , there can be a three-period fixed point and if there is a three-period fixed point, there will also be fixed points of periods,

$$\begin{aligned} &3, 5, 7, 9, \dots \\ &2 \cdot 3, 2 \cdot 5, 2 \cdot 7, 2 \cdot 9, \dots \\ &2^2 \cdot 3, 2^2 \cdot 5, 2^2 \cdot 7, \dots \end{aligned} \quad (98)$$

This is expressed by saying that Period Three Means Chaos.

#### *Chaotic Behaviour of the Non-linear Model*

If  $m$  lies between  $m_8$  and  $m_{16}$ , there will be eight 16-period stable fixed points. If a population size starts from any one of these values, it will oscillate through fifteen other values to return to the original value and this pattern will go on repeating itself. If we draw the graph, it will show rapid oscillations and will look like the graph representing a random phenomenon. Our model is perfectly deterministic, though its behaviour may *appear* to be random and stochastic.

#### *Special Features of Non-linear Difference Equation Models*

The simple model illustrates the differences in behaviour between difference and differential equation models. The problems of existence and uniqueness of solutions, of the stability of equilibrium positions are all different due to the basic fact that in spite of similarities, the Discrete and the Continuous are really different.

## **Age-Structured Population Models**

Let  $x_1(t), x_2(t), \dots, x_p(t)$  be the population sizes of  $p$  pre-reproductive age-groups at time  $t$ ;

Let  $x_{p+1}(t), x_{p+2}(t), \dots, x_{p+q}(t)$  be the population sizes of  $q$  reproductive age-groups at time  $t$ , and

Let  $x_{p+q+1}(t), x_{p+q+2}(t), \dots, x_{p+q+r}(t)$  be the population sizes of  $r$  post-reproductive age-groups at time  $t$ .

Let  $b_{p+1}, b_{p+2}, \dots, b_{p+q}$  be the birth rates i.e. the number of births per unit time per individual in the reproductive age groups.

In other age-groups, the birth rates are zero.

Let  $d_1, d_2, \dots, d_{p+q+r}$  be the death rates in the  $p + q + r$  age-groups.

Let  $m_1, m_2, \dots, m_{p+q+r}$ , be the rates of migration to the next age-groups, then we get the system of difference equations

$$\begin{aligned}
 x_1(t+1) &= b_{p+1}x_{p+1}(t) + \dots + b_{p+q}x_{p+q}(t) - (d_1 + m_1)x_1(t) \\
 x_2(t+1) &= m_1x_1(t) - (d_2 + m_2)x_2(t) \\
 &\dots \dots \dots \\
 x_{p+q+r-1}(t+1) &= m_{p+q+r-2}(t) - (d_{p+q+r-1} + m_{p+q+r-1})x_{p+q+r-1}(t) \\
 x_{p+q+r}(t+1) &= m_{p+q+r-1}x_{p+q+r-1}(t) - (d_{p+q+r})x_{p+q+r}(t)
 \end{aligned} \tag{99}$$

which can be written in the matrix form

$$X(t+1) = LX(t), \tag{100}$$

where

$$X(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_{p+q+r}(t) \end{bmatrix},$$

$$L = \begin{bmatrix}
 -(d_1 + m_1) & 0 & 0 \dots 0 & b_{p+1} & b_{p+2} & \dots & b_{p+q} & 0 & \dots & 0 & 0 \\
 m_1 & -(d_2 + m_2) & 0 \dots & 0 & 0 & \dots & 0 & 0 & \dots & 0 & 0 \\
 0 & m_2 & -(d_3 + m_3) & 0 & 0 & \dots & 0 & 0 & \dots & 0 & 0 \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 0 & 0 & 0 \dots & 0 & 0 & & 0 & 0 & \dots & m_{n-1} - d_n
 \end{bmatrix} \tag{101}$$

where  $p + q + r = n$ .

$L$  is called the Leslie matrix. All the elements of its main diagonal are negative and all the elements of its main subdiagonal are positive. In addition  $q$  elements in the first row are positive and the rest of the elements are all zero. The solution of (100) can be written as

$$X(t) = L^t X(0) \tag{102}$$

Now the Leslie matrix has the property that it has a dominant eigenvalue which is real and positive, which is greater in absolute value than any other eigenvalue and for which the corresponding eigenvector has all its components positive. If this dominant eigenvalue is greater than unity, then the populations of all age-groups will increase exponentially and if it is less than unity the population of all age-groups will die out. If this dominant eigenvalue is unity, the population can have a stable age structure.



## Mathematical Modelling through Difference Equations in Genetics

### (a) *Hardy-Weinberg Law*

Every characteristic of an individual, like height or colour of the hair, is determined by a pair of genes, one obtained from the father and the other obtained from the mother. Every gene occurs in two forms, a dominant

(denoted by a capital letter say  $G$ ) and a recessive (denoted by the corresponding small letter say  $g$ ). Thus with respect to a characteristic, an individual may be a dominant ( $GG$ ), a hybrid ( $Gg$  or  $gG$ ) or a recessive ( $gg$ ).

In the  $n$ th generation, let the proportions of dominants, hybrids and recessives be  $p_n, q_n, r_n$  so that

$$p_n + q_n + r_n = 1, \quad p_n \geq 0, q_n \geq 0, r_n \geq 0 \quad (103)$$

We assume that individuals, in this generation mate at random. Now  $p_{n+1}$  = the probability that an individual in the  $(n+1)$ th generation is a dominant ( $GG$ ) = (probability that this individual gets a  $G$  from the father)  $\times$  (probability that the individual gets a  $G$  from the mother)

$$= \left(p_n + \frac{1}{2}q_n\right)\left(p_n + \frac{1}{2}q_n\right) = \left(p_n + \frac{1}{2}q_n\right)^2$$

$$\text{or} \quad p_{n+1} = \left(p_n + \frac{1}{2}q_n\right)^2 \quad (104)$$

$$\text{Similarly} \quad q_{n+1} = 2\left(p_n + \frac{1}{2}q_n\right)\left(r_n + \frac{1}{2}q_n\right) \quad (105)$$

$$r_{n+1} = \left(r_n + \frac{1}{2}q_n\right)^2, \quad (106)$$

$$\text{so that} \quad p_{n+1} + q_{n+1} + r_{n+1} = \left(p_n + \frac{1}{2}q_n + \frac{1}{2}q_n + r_n\right)^2 = 1, \quad (107)$$

as expected. Similarly

$$\begin{aligned} p_{n+2} &= \left(p_{n+1} + \frac{1}{2}q_{n+1}\right)^2 \\ &= \left(\left(p_n + \frac{1}{2}q_n\right)^2 + \left(p_n + \frac{1}{2}q_n\right)\left(r_n + \frac{1}{2}q_n\right)\right)^2 \\ &= \left(p_n + \frac{1}{2}q_n\right)^2 \left(p_n + \frac{1}{2}q_n + \frac{1}{2}q_n + r_n\right)^2 \end{aligned}$$



$$= \left( p_n + \frac{1}{2} q_n \right)^2 = p_{n+1} \quad (108)$$

$$\text{and } q_{n+2} = q_{n+1}, \quad r_{n+2} = r_{n+1}, \quad (109)$$

so that the proportions of dominants, hybrids and recessives in the  $(n + 2)$ th generation are same as in the  $(n + 1)$ th generation.

Thus in any population in which random mating takes place with respect to a characteristic, the proportions of dominants, hybrids and recessive do not change after the first generation. This is known as Hardy-Weinberg law after the mathematician Hardy and geneticist Weinberg who jointly discovered it.

The equations (104)–(107) is a set of difference equations of the first order.

(b) *Improvement of Plants through selection*  
Suppose the recessives are undesirable and as such we do not allow the recessives in any generation to breed.

Let  $p_n, q_n, r_n$  be the proportions of dominants, hybrids and recessives before elimination of recessives and let  $p'_n, q'_n, 0$  be the populations after the elimination, then

$$\frac{p'_n}{p_n + q_n} = \frac{q'_n}{p_n + q_n} = \frac{p'_n + q'_n}{p_n + q_n} = \frac{1}{1 - r_n} \quad (110)$$

Now we allow random mating and let  $p_{n+1}, q_{n+1}, r_{n+1}$  be the proportions in the next generation before elimination of recessives, then using (104)–(108)

$$p_{n+1} = \left( p'_n + \frac{1}{2} q'_n \right)^2 \quad (111)$$

$$q_{n+1} = 2 \left( p'_n + \frac{1}{2} q'_n \right) \left( \frac{1}{2} q'_n \right) = q'_n \left( p'_n + \frac{1}{2} q'_n \right) \quad (112)$$

$$r_{n+1} = \left( \frac{1}{2} q'_n \right)^2 = \frac{1}{4} q_n'^2 \quad (113)$$

After elimination of recessives, let the new proportions be  $p'_{n+1}, q'_{n+1}, 0$ , so that

$$\frac{p'_{n+1}}{p_{n+1} + q_{n+1}} = \frac{q'_{n+1}}{p_{n+1} + q_{n+1}} = \frac{1}{p_{n+1} + q_{n+1}} = \frac{1}{1 - \frac{1}{4} q_n'^2} \quad (114)$$

$$\begin{aligned} \text{so that } q'_{n+1} &= \frac{q'_n (p'_n + \frac{1}{2} q'_n)}{1 - \frac{1}{4} q_n'^2} = \frac{q'_n (1 - \frac{1}{4} q_n'^2)}{1 - \frac{1}{4} q_n'^2} \\ &= \frac{q'_n}{1 + \frac{1}{4} q_n'^2} \end{aligned} \quad (115)$$





or 
$$P(t+1) = AP(t), \quad (122)$$

where  $P(t)$  is a probability vector and  $A$  is a matrix, all of whose elements lie between zero and unity (since these are all probabilities). Further the sum of elements of every column is unity, since the sum of elements of the  $i$ th column is  $\sum_{j=1}^n p_{ij}$  as this denotes the sum of the probabilities of the system going from the  $i$ th state to any other state and this sum must be unity.

The solution of the matrix difference equation (122) is

$$P(t) = A^t P(0) \quad (123)$$

If all the eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$  of  $A$  are distinct, we can write

$$A = SAS^{-1} \quad (124)$$

where

$$A = \begin{bmatrix} \lambda_1 & 0 & 0 & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \lambda_n \end{bmatrix} \quad (125)$$

so that

$$\begin{aligned} A^t &= (SAS^{-1})(SAS^{-1}) \dots (SAS^{-1}) \\ &= SA^t S^{-1} \\ &= S \begin{bmatrix} \lambda_1^t & 0 & 0 & \dots & 0 \\ 0 & \lambda_2^t & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \lambda_n^t \end{bmatrix} S^{-1} \end{aligned} \quad (126)$$

The probability vector will not change if  $P(t+1) = P(t)$  so that from (122)

$$(I - A)P(t) = 0 \quad (127)$$

Thus if  $P$  is the eigenvector of the matrix  $A$  corresponding to unit eigenvalue, then  $P$  does not change i.e. if the system start with probability vector  $P$  at time 0, it will always remain in this state. Even if the system starts from any other probability vector, it will ultimately be described by the probability vector  $P$  as  $t \rightarrow \infty$ .

As a special case, suppose we have a machine which can be in two states, working or non-working. Let the probability of its transition from working to non-working be  $\alpha$ , of its transition from non-working to working be  $\beta$ , then the transition probability matrix  $A$  is obtained from

$$\begin{array}{cc} & \begin{array}{cc} \text{working} & \text{non-working} \end{array} \\ \begin{array}{c} \text{working} \\ \text{non-working} \end{array} & \begin{bmatrix} 1 - \alpha & \alpha \\ \beta & 1 - \beta \end{bmatrix} \end{array} \quad (128)$$



The system of difference equations is

$$\begin{aligned} p_1(t+1) &= p_1(t)(1-\alpha) + p_2(t)\beta \\ p_2(t+1) &= p_1(t)\alpha + p_2(t)(1-\beta) \end{aligned} \quad (129)$$

or

$$\begin{bmatrix} p_1(t+1) \\ p_2(t+1) \end{bmatrix} = \begin{bmatrix} 1-\alpha & \beta \\ \alpha & 1-\beta \end{bmatrix} \begin{bmatrix} p_1(t) \\ p_2(t) \end{bmatrix} \quad (130)$$

The eigenvalues of the matrix  $A$  is given by

$$\begin{vmatrix} 1-\alpha-\lambda & \beta \\ \alpha & 1-\beta-\lambda \end{vmatrix} = 0 \text{ or } (\lambda-1)(\lambda-1-\alpha-\beta) = 0 \quad (131)$$

The eigenvector corresponding to the unit eigenvalue is  $\beta/(\alpha+\beta)$ ,  $\alpha/(\alpha+\beta)$  and as such ultimately the probability of the machines being found in working order is  $\beta/(\alpha+\beta)$  and the probability of its being found in a non-working state is  $\alpha/(\alpha+\beta)$ .

## MISCELLANEOUS EXAMPLES OF MATHEMATICAL MODELLING THROUGH DIFFERENCE EQUATIONS

Difference equations arise in economics since values of prices, quantities, national income, savings, investments at discrete intervals of time are related. These arise in genetics because proportions of dominants, hybrids and recessives in different generations are related by genetic laws. These arise in population dynamics because population sizes at discrete instants of time are related by births, deaths, immigration and emigration. These arise in finance because amounts at discrete instants of time are related by rates of interest. These arise in gambler's ruin problem because the probability of ruin (or duration of game) when gambler's capital is  $n$  is related to the probability of ruin (or duration of game) when his capital is  $n+1$ .

Similarly in geometry, difference-equations can arise because the number of compartments in which  $n$  lines or curves divide a plane or surface is related to the number of components determined by  $(n+1)$  lines or curves; in dynamics the ranges after successive rebounds of an elastic ball from a horizontal or inclined place are related; in electrical currents, the potential at

neighbouring nodes and currents in neighbouring circuits are related by Kirchhoff's laws and so on.

**POSSIBLE QUESTIONS**

**Part B (6 Marks)**

1. Give any two disciplines that difference equation arises.
2. Write about Hardy-Weinberg law.
3. Write an explanatory note on complementary function.
4. Discuss about application to actuarial science.
5. Find a solution of linear difference equation by Laplace transform.
6. Explain in detail Harrod model..

**Part C (10 Marks)**

1. Explain in detail markov chains.
2. Write about Hardy-Weinberg law.
3. Discuss in detail on particular solution.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
 (Deemed to be University Established Under Section 3 of UGC Act 1956)  
 Pollachi Main Road, Eachanari (Po),  
 Coimbatore –641 021

**UNIT-IV**

**Subject: Mathematical Modeling**

**Subject Code: 16MMP303**

**Mathematical Modeling through Difference Equations**

**Part-A(20X1=20 Marks)**

**(Question Nos. 1 to 20 Online Examinations)**

**Multiple Choice Questions**

Question	Opt 1	Opt 2	Opt 3	Opt 4	Answer
In _____ the genetic characteristics will change generation to generation and the variable representing generation is discrete variable.	Economics	Genetics	Population dynamics	None	Genetics
In genetics, the genetic characteristics will change generation to generation and the variable representing generation is _____ variable.	Discrete	Numeric	Feasible	Optimum	Discrete
In _____ the price changes are consider from year to year or month to month or week to week or day to day	Economics	Genetics	Population dynamics	None	Economics
In _____ the changes are consider in population from one age group to another and the variable representing the age group is discrete	Economics	Genetics	Population dynamics	None	Population dynamics
In population dynamics _____ the changes are consider in population from one age group to another and the variable representing the age group is _____	Discrete	Numeric	Feasible	Optimum	Discrete
No of birth and deaths are proportional to the population then the model is _____	PGM	LGM	PTM	CM	PGM
The solution of linear differential equation is of the form _____	CF+PI	CF-PI	CF*PI	CF/PI	CF+PI
CF denotes _____	Complementary function	convergen function	Conditional function	None	Complementary function
the sort form of particular integral is _____	PI	Par-Ing	Ping	None	PI
Complementary function can be obtained by _____	Matrix	Determinate	Eigen value	None	Matrix
the solution of linear differences equation can be obtained by _____ transform if t is continuous	Laplace	Z	Fourier	Gauss	Laplace
the _____ is solution of linear differences equation can be obtained by _____ transform if t is discrete	Laplace	Z	Fourier	Gauss	Z
the non linear difference equations reducible to linear equation by _____ method	Substitution	Direct	Indirect	Normal	Substitution
In difference equation _____ theory is applied	Stability	Non stability	Uniformity	Non uniformity	Stability
The Horrod model is used in the field of _____	Economics	Genetics	Population dynamics	None	Economics
The investment depends on _____ between the income of current year and last year	Addition	Difference	product	division	Difference
All the saving made are invested in the Horrod model then	$S(t) = I(t)$	$S(t) = 1/2 I(t)$	$2S(t) = I(t)$	None	$S(t) = I(t)$
In the cobweb model price of the commodity in the year denotes	Pt	qt	rt	st	pt
In the cobweb model amount of the commodity available in the market in year t denotes	Pt	qt	rt	st	qt
Amount of the commodity produced this year available for sale is a _____ function of the price of commodity	Linear	Non linear	Stable	Non stable	Linear
In the cobweb path ABCEFGI, .. And the equilibrium price and quantity are given by _____ of two straight lines	Intersection	Union	Disjunction	Conjunction	Intersection
In the cobweb path ABCEFGI, .. And the equilibrium price and quantity are given by intersection of two _____	Straight lines	Circles	Squares	Cubes	Straight lines
the actuarial science is called _____	Mathematics of finance	Mathematics of economics	Dynamics	Statics	Mathematics of finance
the actuarial science is called _____	Mathematics of investment	Mathematics of economics	Dynamics	Statics	Mathematics of investment
One-period fixed points and their stability	$Y_{t+1}=y_t$	$Y_{t+2}=y_{2t}$	$Y_{t-1}=y_t$	$y_{t+2}=y_t$	$Y_{t+1}=y_t$
Two-period fixed points and their stability	$Y_{t+2}=y_t$	$Y_{t+2}=y_{2t}$	$Y_{t-1}=y_t$	$y_{t+2}=y_{2t}$	$Y_{t+2}=y_t$
Any population in which random meeting take place with respect to a characteristic , the proportion of dominants hybrids and recessive do not change after the first generation states _____ law	Gauss	Hardy-weinberg	Fick's	Routhwelt	Hardy-weinberg
The probability of transition from state I to state j is	Markov chain	Hardy-weinberg	Fick's	Routhwelt	Markov chain






**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
(Deemed to be University Established Under Section 3 of UGC Act 1956)  
Pollachi Main Road, Eachanari (Po),  
Coimbatore –641 021  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>

---

## UNIT V

Mathematical Modeling through Graphs: Solutions that can be Modeled through Graphs – Mathematical Modeling in Terms of Directed Graphs, Signed Graphs, Weighted Digraphs and Un oriented Graphs.

### SUGGESTED READINGS

#### TEXT BOOK

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

#### REFERENCES

**R3:** Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.

# Unit-V

## **Mathematical Modelling Through Graphs**



## SITUATIONS THAT CAN BE MODELLED THROUGH GRAPHS

It has been stated that “Applied Mathematics is nothing but solution of differential equations”. This statement is wrong on many counts (i) Applied Mathematics also deals with solutions of difference, differential-difference, integral, integro-differential, functional and algebraic equations (ii) Applied Mathematics is equally concerned with inequations of all types (iii) Applied Mathematics is also concerned with mathematical modelling; in fact mathematical modelling has to precede solution of equations (iv) Applied Mathematics also deals with situations which cannot be modelled in terms of equations or inequations; one such set of situations is concerned with qualitative relations.

Mathematics deals with both quantitative and qualitative relationships. Typical qualitative relations are:  $y$  likes  $x$ ,  $y$  hates  $x$ ,  $y$  is superior to  $x$ ,  $y$  is subordinate to  $x$ ,  $y$  belongs to same political party as  $x$ , set  $y$  has a non-null intersection with set  $x$ ; point  $y$  is joined to point  $x$  by a road, state  $y$  can be transformed into state  $x$ , team  $y$  has defeated team  $x$ ,  $y$  is father of  $x$ , course  $y$  is a prerequisite for course  $x$ , operation  $y$  has to be done before operation  $x$ , species  $y$  eats species  $x$ ,  $y$  and  $x$  are connected by an airline,  $y$  has a healthy influence on  $x$ , any increase of  $y$  leads to a decrease in  $x$ ,  $y$  belongs to same caste as  $x$ ,  $y$  and  $x$  have different nationalities and so on.

Such relationships are very conveniently represented by graphs where a graph consists of a set of vertices and edges joining some or all pairs of these vertices. To motivate the typical problem situations which can be modelled through graphs, we consider the first problem so historically modelled viz. the problem of seven bridges of Königsberg.

### The Seven Bridges Problem

There are four land masses  $A, B, C, D$  which are connected by seven bridges numbered 1 to 7 across a river (Figure 7.1). The problem is to start from any point in one of the land masses, cover each of the seven bridges once and once only and return to the starting point.




Figure 7.1




Figure 7.2



There are two ways of attacking this problem. One method is to try to solve the problem by walking over the bridges. Hundreds of people tried to do so in their evening walks and failed to find a path satisfying the conditions of the problem. A second method is to draw a scale map of the bridges on paper and try to find a path by using a pencil.

It is at this stage that concepts of mathematical modelling are useful. It is obvious that the sizes of the land masses are unimportant, the lengths of the bridges or even whether these are straight or curved are irrelevant. What is relevant information is that  $A$  and  $B$  are connected by two bridges 1 and 2,  $B$  and  $C$  are connected by two bridges 3 and 4,  $B$  and  $D$  are connected by one bridge number 5,  $A$  and  $D$  are connected by bridge number 6 and  $C$  and  $D$  are connected by bridge number 7. All these facts are represented by the graph with four vertices and seven edges in Figure 7.2. If we can trace this graph in such a way that we start with any vertex and return to the same vertex and trace every edge once and once only without lifting the pencil from the paper, the problem can be solved. Again trial and error method cannot be satisfactorily used to show that no solution is possible.

The number of edges meeting at a vertex is called the degree of that vertex. We note that the degrees of  $A, B, C, D$  are 3, 5, 3, 3 respectively and each of these is an odd number. If we have to start from a vertex and return to it, we need an even number of edges at that vertex. Thus it is easily seen that Königsberg bridges problem cannot be solved.

This example also illustrates the power of mathematical modelling. We have not only disposed of the seven-bridges problem, but we have discovered a technique for solving many problems of the same type.

## Some Types of Graphs

A graph is called *complete* if every pair of its vertices is joined by an edge (Figure 7.3(a)).

A graph is called a *directed graph* or a *digraph* if every edge is directed with an arrow. The edge joining  $A$  and  $B$  may be directed from  $A$  to  $B$  or from  $B$  to  $A$ . If an edge is left undirected in a digraph, it will be assumed to be directed both ways (Figure 7.3(b)).




Figure 7.3a




Figure 7.3b






Figure 7.3c




Figure 7.3d

A graph is called a *signed graph* if every edge has either a plus or minus sign associated with it (Figure 7.3(c)).

A digraph is called a *weighted digraph* if every directed edge has a weight (giving the importance of the edge) associated with it (Figure 7.3(d)). We may also have digraphs with positive and negative numbers associated with edges. These will be called *weighted signed digraphs*.

### Nature of Models in Terms of Graphs

In all the applications we shall consider, the length of the edge joining two vertices will not be relevant. It will not also be relevant whether the edge is straight or curved. The relevant facts would be (a) which edges are joined; (b) which edges are directed and in which direction(s); (c) which edges have positive or negative signs associated with them; (d) which edges have weights associated with them and what these weights are.

## MATHEMATICAL MODELS IN TERMS OF DIRECTED GRAPHS

### Representing Results of Tournaments

The graph (Figure 7.4) shows that




Figure 7.4

- (i) Team A has defeated teams B, C, E.
- (ii) Team B has defeated teams C, E.
- (iii) Team E has defeated D.
- (iv) Matches between A and D, B and D, C and D and C and E have yet to be played.

### One-Way Traffic Problems



The road map of a city can be represented by a directed graph. If only one-way traffic is allowed from point  $a$  to point  $b$ , we draw an edge directed from  $a$  to  $b$ . If traffic is allowed both ways, we can either draw two edges, one directed from  $a$  to  $b$  and the other directed from  $b$  to  $a$  or simply draw an undirected edge between  $a$  and  $b$ . The problem is to find whether we can introduce one-way traffic on some or all of the roads without preventing persons from going from any point of the city to any other point. In other words, we have to find when the edges of a graph can be given direction in such a way that there is a directed path from any vertex to every other. It is easily seen that one-way traffic on the road  $DE$  cannot be introduced without disconnecting the vertices of the graph (Figure 7.5).




Figure 7.5(a)




Figure 7.5(b)

In Figure 7.5(a),  $DE$  can be regarded as a bridge connecting two regions of the town. In Figure 7.5(b)  $DE$  can be regarded as a blind street on which a two-way traffic is necessary. Edges like  $DE$  are called *separating edges*, while other edges are called *circuit edges*. It is necessary that on separating

edges, two-way traffic should be permitted. It can also be shown that this is sufficient. In other words, the following theorem can be established:

If  $G$  is an undirected connected graph, then one can always direct the circuit edges of  $G$  and leave the separating edges undirected (or both way directed) so that there is a directed path from any given vertex to any other vertex.

## Genetic Graphs

In a genetic graph, we draw a directed edge from  $A$  to  $B$  to indicate that  $B$  is the child of  $A$ . In general each vertex will have two incoming edges, one from the vertex representing the father and the other from the vertex representing the mother. If the father or mother is unknown, there may be less than two incoming edges. Thus in a genetic graph, the local degree of incom-

ing edges at each vertex must be less than or equal to two. This is a necessary condition for a directed graph to be a genetic graph, but it is not a sufficient condition. Thus Figure 7.6 does not give a genetic graph inspite of the fact that the number of incoming edges at each vertex does not exceed two. Suppose  $A_1$  is male, then  $A_2$  must be female, since  $A_1, A_2$  have a child  $B_1$ . Then  $A_3$  must be male, since  $A_2, A_3$  have a child  $B_2$ . Now  $A_1, A_3$  being both males cannot have a child  $B_3$ .




Figure 7.6

### Senior-Subordinate Relationship

If  $a$  is senior to  $b$ , we write  $aSb$  and draw a directed edge from  $a$  to  $b$ . Thus the organisational structure of a group may be represented by a graph like the following [Figure 7.7].




Figure 7.7

The relationship  $S$  satisfies the following properties:

- (i)  $\sim(aSa)$  i.e. no one is his own senior
- (ii)  $aSb \Rightarrow \sim(bSa)$  i.e.  $a$  is senior to  $b$  implies that  $b$  is not senior to  $a$
- (iii)  $aSb, bSc \Rightarrow aSc$  i.e. if  $a$  is senior to  $b$  and  $b$  is senior to  $c$ , then  $a$  is senior to  $c$ .

The following theorem can easily be proved: "The necessary and sufficient condition that the above three requirements hold is that the graph of an organisation should be free of cycles"

We want now to develop a *measure for the status* of each person. The status  $m(x)$  of the individual should satisfy the following reasonable requirements.



- (i)  $m(x)$  is always a whole number
- (ii) If  $x$  has no subordinate,  $m(x) = 0$
- (iii) If, without otherwise changing the structure, we add a new individual subordinate to  $x$ , then  $m(x)$  increases
- (iv) If, without otherwise changing the structure, we move a subordinate of  $a$  to a lower level relative to  $x$ , then  $m(x)$  increases.

A measure satisfying all these criteria was proposed by Harary. We define the level of seniority of  $x$  over  $y$  as the length of the shortest path from  $x$  to  $y$ . To find the measure of status of  $x$ , we find  $n_1$ , the number of individuals who are one level below  $x$ ,  $n_2$  the number of individuals who are two levels below  $x$  and in general, we find  $n_k$  the number of individuals who are  $k$  levels below  $x$ . Then the Harary measure  $h(x)$  is defined by

$$h(x) = \sum_k kn_k \quad (1)$$

It can be shown that among all the measure which satisfy the four requirements given above, Harary measure is the least.

If however, we define the level of seniority of  $x$  over  $y$  as the length of the longest path from  $x$  to  $y$ , and then find  $H(x) = \sum_k kn_k$ , we get another measure which will be the largest among all measures satisfying the four requirements. For Figure 7.8, we get

$$\begin{array}{ll} h(a) = 1.2 + 4.2 + 2.3 = 16 & H(a) = 1.1 + 3.2 + 2.3 + 2.4 = 21 \\ h(b) = 1.3 + 2.4 = 11 & H(b) = 2.1 + 2.2 + 2.3 + 1.4 = 16 \\ h(c) = 1.2 + 1.2 = 4 & H(c) = 1.1 + 1.2 + 1.3 = 6 \end{array}$$




Figure 7.8

$$\begin{array}{ll} h(d) = 1.1 & = 1 & H(d) = 1.1 & = 1 \\ h(e) = 1.3 & = 3 & H(e) = 1.2 + 2.1 & = 4 \\ h(f) = 1.1 & = 1 & H(f) = 1.1 & = 1 \\ h(g) = 1.2 & = 2 & H(g) = 1.2 & = 2 \\ h(h) & = 0 & H(h) & = 0 \\ h(i) & = 0 & H(i) & = 0 \end{array}$$



## Food Webs

Here  $aSb$  if  $a$  eats  $b$  and we draw a directed edge from  $a$  to  $b$ . Here also  $\sim(aSa)$  and  $aSb \Rightarrow \sim(bSa)$ . However the transitive law need not hold. Thus consider the food web in Fig. 7.9. Here fox eats bird, bird eats grass, but fox does not eat grass.




Figure 7.9

We can however calculate measure of the status of each species in this food web by using (1)  $h(\text{bird}) = 2$ ,  $h(\text{fox}) = 4$ ,  $h(\text{insect}) = 1$ ,  $h(\text{grass}) = 0$ ,  $h(\text{deer}) = 1$ .

## Communication Networks

A directed graph can serve as a model for a communication network. Thus consider the network given in Figure 7.10. If an edge is directed from  $a$  to  $b$ , it means that  $a$  can communicate with  $b$ . In the given network  $e$  can communicate directly with  $b$ , but  $b$  can communicate with  $e$  only indirectly through  $c$  and  $d$ . However every individual can communicate with every other individual.

Our problem is to determine the importance of each individual in this network. The importance can be measured by the fraction of the messages on an average that pass through him. In the absence of any other knowledge, we can assume that

if an individual can send message direct to  $n$  individuals, he will send a message to any one of them with probability  $1/n$ . In the present example, the communication probability matrix is




Figure 7.10

$$\begin{matrix}
 & \begin{matrix} a & b & c & d & e \end{matrix} \\
 \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} & \begin{bmatrix} 0 & 1/2 & 1/2 & 0 & 0 \\ 1/2 & 0 & 1/2 & 0 & 0 \\ 1/3 & 1/3 & 0 & 1/3 & 0 \\ 0 & 0 & 1/2 & 0 & 1/2 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix}
 \end{matrix} \quad (2)$$

No individual is to send a message to himself and so all diagonal elements are zero. Since all elements of the matrix are non-negative and the sum of elements of every row is unity, the matrix is a stochastic matrix and one of its eigenvalues is unity. The corresponding normalised eigenvector is  $[11/45, 13/45, 3/10, 1/10, 1/15]$ . In the long run, these fractions of messages will pass through  $a, b, c, d, e$  respectively. Thus we can conclude that in this network,  $c$  is the most important person.

If in a network, an individual cannot communicate with every other individual either directly or indirectly, the Markov chain is not ergodic and the process of finding the importance of each individual breaks down.

### Matrices Associated with a Directed Graph

For a directed graph with  $n$  vertices, we define the  $n \times n$  matrix  $A = (a_{ij})$  by  $a_{ij} = 1$  if there is an edge directed from  $i$  to  $j$  and  $a_{ij} = 0$  if there is no edge directed from  $i$  to  $j$ . Thus the matrix associated with the graph of Figure 7.11 is given by

$$A = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix} \end{matrix} \quad (3)$$

We note that (i) the diagonal elements of the matrix are all zero (ii) the number of non-zero elements is equal to the number of edges (iii) the number of non-zero elements in any row is equal to the local outward degree of the vertex corresponding to the row (iv) the number of non-zero elements in a column is equal to the local inward degree of the vertex corresponding to the column. Now




Figure 7.11

$$A^2 = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 2 & 1 & 1 & 0 \\ 1 & 2 & 1 & 0 \\ 1 & 1 & 2 & 0 \\ 1 & 2 & 1 & 0 \end{bmatrix} \end{matrix} = (a_{ij}^{(2)}) \quad (4)$$



The element  $a_{ij}^{(2)}$  gives the number of 2-chains from  $i$  to  $j$ . Thus from vertex 2 to vertex 1, there are two 2-chains viz. via vertex 3 and vertex 4. We can generalise this result in the form of a theorem viz. "The element  $a_{ij}^{(2)}$  of  $A^2$  gives the number of 2-chains i.e. the number of paths with two-edges from vertex  $i$  to vertex  $j$ ".

The theorem can be further generalised to "The element  $a_{ij}^{(m)}$  of  $A^m$  gives the number of  $m$ -chains i.e. the number of paths with  $m$  edges from vertex  $i$  to vertex  $j$ ". It is also easily seen that "The  $i$ th diagonal element of  $A^2$  gives the number of vertices with which  $i$  has symmetric relationship".

From the matrix  $A$  of a graph, a symmetric matrix  $S$  can be generated by taking the elementwise product of  $A$  with its transpose so that in our case

$$S = A \times A^T = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix} \times \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 1 & 0 \\ & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (5)$$

$S$  obviously is the matrix of the graph from which all unreciprocated connections have been eliminated. In the matrix  $S$  (as well as in  $S^2, S^3, \dots$ ) the elements in the row and column corresponding to a vertex which has no symmetric relation with any other vertex are all zero.

### Application of Directed Graphs to Detection of Cliques

A subset of persons in a socio-psychological group will be said to form a clique if (i) every member of this subset has a symmetrical relation with every other member of this subset (ii) no other group member has a symmetric relation with all the members of the subset (otherwise it will be included in the clique) (iii) the subset has at least three members.

If other words a clique can be defined as a maximal completely connected subset of the original group, containing at least three persons. This subset should not be properly contained in any larger completely connected subset.

If the group consists of  $n$  persons, we can represent the group by  $n$  vertices of a graph. The structure is provided by persons knowing or being connected to other persons. If a person  $i$  knows  $j$ , we can draw a directed edge from  $i$  to  $j$ . If  $i$  knows  $j$  and  $j$  knows  $i$ , then we have a symmetrical relation between  $i$  and  $j$ .



With this interpretation, the graph of Figure 7.11 shows that persons 1, 2, 3 form a clique. With very small groups, we can find cliques by carefully observing the corresponding graphs. For larger groups analytical methods based on the following results are useful: (i)  $i$  is a member of a clique if the  $i$ th diagonal element of  $S^3$  is different from zero. (ii) If there is only one clique of  $k$  members in the group, the corresponding  $k$  elements of  $S^3$  will be  $(k-1)(k-2)/2$  and the rest of the diagonal elements will be zero. (iii) If there are only two cliques with  $k$  and  $m$  members respectively and there is no element common to these cliques, then  $k$  elements of  $S^3$  will be  $(k-1)(k-2)/2$ ,  $m$  elements of  $S^3$  will be  $(m-1)(m-2)/2$  and the rest of the elements will be zero. (iv) If there are  $m$  disjoint cliques with  $k_1, k_2, \dots, k_m$  members, then the trace of  $S^3$  is  $\frac{1}{2} \sum_{i=1}^m k_i(k_i-1)(k_i-2)$ . (v) A member is non-cliquical if only if the corresponding row and column of  $S^2 \times S$  consists entirely of zeros.

## MATHEMATICAL MODELS IN TERMS OF SIGNED GRAPHS

### Balance of Signed Graphs

A signed (or an algebraic) graph is one in which every edge has a positive or negative sign associated with it. Thus the four graphs of Figure 7.16 are signed graphs. Let positive sign denote friendship and negative sign denote enmity, then in graph (i) A is a friend of both B and C and B and C are




Figure 7.16

also friends. In graph (ii) A is friend of B and A and B are both jointly enemies of C. In graph (iii), A is a friend of both B and C, but B and C are enemies. In graph (iv) A is an enemy of both B and C, but B and C are not friends.

The first two graphs represent normal behaviour and are said to be balanced, while the last two graphs represent unbalanced situations since if A is a friend both B and C and B and C are enemies, this creates a tension in the system and there is a similar tension when B and C have a common enemy A, but are not friends of each other.



We define the sign of a cycle as the product of the signs of component edges. We find that in the two balanced cases, this sign is positive and in the two unbalanced cases, this is negative.

We say that a cycle of length three or a triangle is balanced if and only if its sign is positive. A complete algebraic graph is defined to be a complete graph such that between any two edges of it, there is a positive or negative sign. A complete algebraic graph is said to be balanced if all its triangles are balanced. An alternative definition states that a complete algebraic graph is balanced if all its cycles are positive. It can be shown that the two definitions are equivalent.

A graph is locally balanced at a point  $a$  if all the cycles passing through  $a$  are balanced. If a graph is locally balanced at all points of the graph, it will obviously be balanced. A graph is defined to be  $m$ -balanced if all its cycles of length  $m$  are positive. For an incomplete graph, it is preferable to define it to be balanced if all its cycles are positive. The definition in terms of triangle is not satisfactory, as there may be no triangles in the graph.

## Structure Theorem and Its Implications

**Theorem.** The following four conditions are equivalent:

- (i) The graph is balanced i.e. every cycle in it is positive.
- (ii) All closed line-sequences in the graph are positive i.e. any sequence of edges starting from a given vertex and ending on it and possibly passing through the same vertex more than once is positive.
- (iii) Any two line-sequences between two vertices have the same sign.
- (iv) The set of all points of the graph can be partitioned into two disjoint sets such that every positive sign connects two points in the same set and every negative sign connects two points of different sets.

The last condition has an interesting interpretation with possibility of application. It states that if in a group of persons there are only two possible relationships viz. liking and disliking and if the algebraic graph representing these relationships is balanced, then the group will break up into two separate parties such that persons within a party like one another, but each person of one party dislikes every person of the other party. If a balanced situation is regarded as stable, this theorem can be interpreted to imply that a two-party political system is stable.



**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
(Deemed to be University Established Under Section 3 of UGC Act 1956)  
Pollachi Main Road, Eachanari (Po),  
Coimbatore –641 021  
**DEPARTMENT OF MATHEMATICS**

---

<b>Subject: Mathematical Modeling</b>	<b>Subject Code: 16MMP303</b>	<b>L T P C</b>
---------------------------------------	-------------------------------	----------------

<b>Class:II M.Sc</b>	<b>Semester:III</b>	<b>4 1 0 4</b>
----------------------	---------------------	----------------

---

### **UNIT V(Cont)**

Mathematical Modeling through Graphs: Solutions that can be Modeled through Graphs – Mathematical Modeling in Terms of Directed Graphs, Signed Graphs, Weighted Digraphs and Un oriented Graphs.

### **SUGGESTED READINGS**

#### **TEXT BOOK**

**T1:** J.N. Kapur, (2015). Mathematical Modeling, Wiley Eastern Limited, New Delhi.

#### **REFERENCES**

**R3:** Frank. R. Giordano, Maurice. D.Weir, WilliamP. Fox, (2003). A first course in Mathematical Modelling, Vikash Publishing House, UK.



## MATHEMATICAL MODELLING IN TERMS OF WEIGHTED DIGRAPHS

### Communication Networks with Known Probabilities of Communication

In the communication graph of Figure 7.10, we know that  $a$  can communi-

cate with both  $b$  and  $c$  only and in the absence of any other knowledge, we assigned equal probabilities to  $a$ 's communicating with  $b$  or  $c$ . However we may have a priori knowledge that  $a$ 's chances of communicating with  $b$  and  $c$  are in the ratio 3:2, then we assign probability .6 to  $a$ 's communicating with  $b$  and .4 to  $a$ 's communicating with  $c$ . Similarly we can associate a probability with every directed edge and we get the weighted digraph (Figure 7.19) with the associated matrix




Figure 7.19

$$B = \begin{matrix} & \begin{matrix} a & b & c & d & e \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} & \begin{bmatrix} 0 & 0.6 & 0.4 & 0 & 0 \\ 0.5 & 0 & 0.5 & 0 & 0 \\ 0.4 & 0.3 & 0 & 0.3 & 0 \\ 0 & 0 & .3 & 0 & 0.7 \\ 0 & 1.0 & 0 & 0 & 0 \end{bmatrix} \end{matrix} \quad (6)$$

We note that the elements are all non-negative and the sum of the elements of every row is unity so that  $B$  is a stochastic matrix and unity is one of its eigenvalues. The eigenvector corresponding to this eigenvalues will be different from the eigenvector found in Section 7.2.6 and so the relative importance of the individuals depends both on the directed edges as well as on the weights associated with the edges.

**Weighted Digraphs and Markov Chains**

A Markovian system is characterised by a transition probability matrix. Thus if the states of a system are represented by  $1, 2, \dots, n$  and  $p_{ij}$  gives the probability of transition from the  $i$ th state to  $j$ th state, the system is characterised by the transition probability matrix (t.p.m)








Figure 7.20

In this example  $d$  is an absorbing state or a state of equilibrium. Once a system reaches the state  $d$ , it stays there for ever.

It is clear from Figure 7.20, that in whichever state, the system may start, it will ultimately end in state  $d$ . However the number of steps that may be required to reach  $d$  depends on chance. Thus starting from  $c$ , the number of steps to reach  $d$  may be 1, 2, 3, 4, ...; starting from  $b$  the number of steps to reach  $d$  may be 2, 3, 4, ... and starting for  $a$ , the number of steps may be 3, 4, 5, ... In each case, we can find the probability that the number of steps required is  $n$  and then we can find the expected number of steps to reach it.

Thus for the matrix

$$\begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} a \\ b \end{matrix} & \begin{bmatrix} 1 & 0 \\ 1/3 & 2/3 \end{bmatrix} \end{matrix} \quad (12)$$

$a$  is an absorbing state. Starting from  $b$ , we can reach  $a$  in 1, 2, 3, ...,  $n$  steps with probabilities  $(1/3)$ ,  $(1/3)(2/3)$ ,  $(1/3)(2/3)^2$ , ...,  $(1/3)(2/3)^{n-1}$ , ..., so that the expected number of steps is

$$\sum_{n=1}^{\infty} n \frac{1}{3} \left(\frac{2}{3}\right)^{n-1} = 3 \quad (13)$$

## General Communication Networks

So far we have considered communication networks in which the weight associated with a directed edge represents the probability of communication along that edge. We can however have more general networks e.g.

(a) for communication of messages where the directed edge represents the channel and the weight represents the capacity of the channel say in bits per second

(b) for communication of gas in pipelines where the weights are the capacities, say in gallons per hour

(c) communication roads where the weights are the capacities is cars per hour.

An interesting problem is to find the maximum flow rate, of whatever is being communicated, from any vertex of the communication network to any other. Useful graph-theoretic algorithms for this have been developed by Elias, Feinstein and Shannon as well as by Ford and Fulkerson.

## Signal Flow Graphs

The system of algebraic equations

$$\begin{aligned} x_1 &= 4y_0 + 6x_2 - 2x_3 \\ x_2 &= 2y_0 - 2x_1 + 2x_3 \\ x_3 &= 2x_1 - 2x_2 \end{aligned} \quad (14)$$

can be represented by the weighted digraph in Figure 7.22. For solving for  $x_1$ , we successively eliminate  $x_3$  and  $x_2$  to get the graphs in Figure 7.23 and finally we get

$$x_1 = 4y_0$$

We can similarly represent the solution of any number of linear equations graphically.




Figure 7.22




Figure 7.23



## Weighted Bipartitic Digraphs and Difference Equations

Consider the system of difference equations

$$\begin{aligned} x_{t+1} &= a_{11}x_t + a_{12}y_t + a_{13}z_t \\ y_{t+1} &= a_{21}x_t + a_{22}y_t + a_{23}z_t \\ z_{t+1} &= a_{31}x_t + a_{32}y_t + a_{33}z_t \end{aligned} \quad (15)$$

This can be represented by a weighted bipartitic digraph (Figure 7.24). The weights can be positive or negative.




Figure 7.24

## MATHEMATICAL MODELLING IN TERMS OF UNORIENTED GRAPHS

### Electrical Networks and Kirchoffs' Laws

For more than a hundred years after Euler solved the Königsberg problem in 1736, graph theory continued to deal with interesting puzzles only. It was in 1849 that Kirchoffs' formulation of his laws of electrical currents in graph-theoretic terms led to interest in serious applications of graph theory

An electrical circuit (Figures 7.25a, b) consists of resistors  $R_1, R_2, \dots$ , inductances  $L_1, L_2, \dots$ , capacitors  $C_1, C_2$  and batteries  $B_1, B_2$ , etc.




Figure 7.25



The network diagram represents two independent aspects of an electrical network. The first gives the interconnection between components and the second gives voltage-current relationship of each component. The first aspect is called network topology and can be modelled graphically. This aspect is independent of voltages and currents. The second aspects involves voltages and current and is modelled through differential equations.

For topological purposes, lengths and shapes of connections are not important and graphs of Figures 7.25(a), 7.25(b) and 7.25(c) are isomorphic.

For stating Kirchhoff's laws, we need two incidence matrices associated with the graph. If  $v$  and  $e$  denote the number of vertices and edges respectively, we define the *vertex* or *incidence* matrix  $A = [a_{ij}]$  as follows:

$$\begin{aligned} a_{ij} &= 1 && \text{if the edge } j \text{ is incident at vertex } i \\ a_{ij} &= 0 && \text{if the edge } j \text{ is not incident at vertex } i \end{aligned}$$

This consists of  $v$  rows and  $e$  columns. For graph 7.25,  $A$  is given by

$$A = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \left[ \begin{array}{cccccc} 0 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{array} \right] \end{matrix} \quad (22)$$

We note that every column has two non-zero elements.

Similarly we define the circuit matrix  $B = [b_{kj}]$  as follows

$$\begin{aligned} b_{kj} &= 1 && \text{if element } j \text{ is in circuit } k \\ &= 0 && \text{if element } j \text{ is not in circuit } k \end{aligned}$$

The matrix  $B$  contains as many rows as there are circuits and it has  $e$  columns. In our case

$$B = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \left[ \begin{array}{cccccc} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 & 1 \end{array} \right] \end{matrix} \quad (23)$$

Now Kirchoff's laws can be written in the matrix form as follows:

$$AI = 0 \text{ (Kirchoff's current law)} \quad (24)$$

$$BV = 0 \text{ (Kirchoff's voltage law)} \quad (25)$$

where  $I$  is an  $e \times 1$  column matrix giving the  $e$  currents and  $V$  is  $e \times 1$  column matrix giving  $e$  voltages.

Matrices  $A$  and  $B$  depend on the graph only, matrices  $I$  and  $V$  depend on currents and voltages only.  $A$  and  $B$  can be written independently of  $I$  and  $V$ . Now an important question is as to how many of the components of the current and voltage vectors are independent.

It can be proved that the rank of  $A$  is  $v - 1$  and the rank of  $B$  is  $e - v + 1$ . Thus  $v - 1$  and  $e - v + 1$  are the numbers of linearly independent Kirchoff's current and voltage equations.

The graph-theoretic methods can now be used to (i) establish the validity of the circuit and vertex equations and find their generalisations (ii) conditions under which unique solutions of these equations exist (iii) justify the duality procedures used in network theory (iv) develop short-cut methods for writing equations (v) develop techniques for network synthesis.

## Map-Colouring Problems

The four colour problem that every plane map, however complex, can be coloured with four colours in such a way that two neighbouring regions get different colours, challenged and fascinated mathematicians for over one hundred years till it was finally solved by Appel and Haken in 1976 by using over 1000 hours of computer time. The problem is essentially




Figure 7.26

graph-theoretic since the sizes and shapes of regions are not important. That four colours are necessary is easily seen by considering the simple graph in Figure 7.26. It was the proof of the sufficiency that took more than hundred years. However the efforts



to solve this problem led to the development of many other graph-theoretic models.

Similar map-colouring problems arise for colouring of maps on surface of a sphere, a torus or other surfaces. However many of these were solved even before the simpler-looking four-colour problem was disposed of.

## Planar Graphs

In printing of T.V. and radio circuits; we want that the wires, all lying in a plane, should not intersect. In the graph of Figure 7.27a wires appear to intersect, but we can find an isomorphic graph in Figure 7.27(b) in which edges do not intersect. A graph which is such that we can draw a graph isomorphic to it in which edges do not intersect is called a planar graph.




Figure 7.27 (a)




Figure 7.27 (b)

A complete graph with five vertices is not planar (Figure 7.28a). We can draw nine of the edges so that these do not intersect (Figure 7.28b) but however we may draw, we cannot draw all the ten edges without at least two of them intersecting. The proof of this depends on Jordan's theorem that every simple closed curve divides the plane into two regions, one inside the curve and one outside the curve.  $ABCDE$  in Figure 7.28(b) is a closed Jordan curve and we cannot draw three edges either inside it or outside it without intersecting.




Figure 7.28 (a)




Figure 7.28 (b)

## Euler's Formula for Polygonal Graphs

A polygonal graph with  $n$  vertices and  $n$  straight or curved edges has  $n$  vertices,  $n$  edges and two faces (one inside and one outside) so that for this graph



$$V - E + F = 2 \quad (26)$$

If we add on one edge, another polygonal region of  $r$  vertices, we increase the number of vertices by  $r - 2$ , the number of edges by  $r - 1$  and the

number of faces by 1, so that the net increases in  $V - E + F$  is zero and the formula (26) remains valid. It can be shown by using the principle of induction that (26) is valid for any polygonal graph with any number of regions.

To draw the dual graph  $G^*$  of  $G$ , we take a point inside each region and draw an edge through it intersecting one of the edges of the region. It is obvious that for this dual graph the number of vertices, edges and faces is given by

$$V^* = F, \quad E = E^*, \quad F^* = V, \quad (30)$$

so that

$$V^* - E^* + F^* = F - E + V = 2, \quad (31)$$

as expected.

## Regular Solids

A polygonal graph  $G$  is said to be completely regular if both  $G$  and its dual  $G^*$  are regular i.e. if the degree of each vertex of  $G$  is the same (say  $\rho$ ) and the degree of each vertex of  $G^*$  is the same (say  $\rho^*$ ). From this definition, it follows

$$2E = \rho V = \rho^* F \quad (32)$$

$$\text{or} \quad E = \frac{1}{2} \rho V, \quad F = \frac{\rho}{\rho^*} V \quad (33)$$

Substituting (33) in (26)

$$V - \frac{1}{2} \rho V + \frac{\rho}{\rho^*} V = 2 \quad (34)$$

$$\text{or} \quad V(2\rho + 2\rho^* - \rho\rho^*) = 4\rho^* \quad (35)$$

Since  $V, \rho, \rho^*$  are positive integers

$$2\rho + 2\rho^* - \rho\rho^* > 0 \quad \text{or} \quad (\rho - 2)(\rho^* - 2) < 4 \quad (36)$$

If  $\rho > 2, \rho^* > 2$ , the only solutions of the inequality (36) are

$$\rho = 3, \quad \rho^* = 3; \quad \rho = 3; \quad \rho^* = 4; \quad \rho = 3, \quad \rho^* = 5; \quad \rho = 4, \quad \rho^* = 3; \\ \rho = 5, \quad \rho^* = 3.$$

Substituting in (35) and (33), we get the table and graphs

	$\rho$	$V$	$E$	$F$	$\rho^*$	$V^*$	$E^*$	$F^*$
(i)	3	4	6	4	3	4	6	4
(ii)	3	8	12	6	4	6	12	8
(iii)	3	20	30	12	5	12	30	20
(iv)	4	6	12	8	3	8	12	6
(v)	5	12	30	20	3	20	30	12

The corresponding graphs are given in Figure 7.29(a)-(e). It is obvious that tetrahedron graph is dual to itself, cube is dual of octahedron and Dodecahedron and Icosahedron are duals of each other.




Figure 7.29

These five graphs corresponding to five Platonic regular solids (Figure 7.30).




Figure 7.30




Figure 7.31

**There is another solution of (36) viz.  $\rho = 2$ ,  $\rho^* = 2, 3, 4, \dots$ . The corresponding graphs  $G$  and  $G^*$  are shown in Figure 7.31.**

### **POSSIBLE QUESTIONS**

#### **Part B (6 Marks)**

- 1.Explain in detail senior-subordinate relationship.
- 2.Write an explanatory note on planar graphs.
- 3.Discuss in detail weighted digraphs and markov chains.
- 4.Write a note on the following
  - i) Signal flow graphs
  - ii) Map-colouring problems
  - iii) Planar graphs
  - iv) Euler's formula for polygonal graphs
- 5.Explain about one-way traffic problems.
- 6.Discuss in detail about communication networks
- 7.Write a note on seven bridges problem.
- 8.Give a brief note on Genetic graphs.

#### **Part C (10 Marks)**

- 1.Discuss in detail about communication networks.
- 2.Give a detailed note on electrical networks and Kirchoff's laws.
- 3.Give a brief note on Genetic graphs.





**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
(Deemed to be University Established Under Section 3 of UGC Act 1956)  
Pollachi Main Road, Eachanari (Po),  
Coimbatore –641 021

## UNIT-V

Subject: Mathematical Modeling

Subject Code: 16MMP303

## Mathematical Modeling through Graphs

## Part-A(20X1=20 Marks)

(Question Nos. 1 to 20 Online Examinations)

## Multiple Choice Questions

Question	Opt 1	Opt 2	Opt 3	Opt 4	Answer
Mathematics deals with both quantative and _____ relationship	Qualitative	Numeric	Decimal	Integer	Qualitative
Apply mathematics deals with solution of _____	Difference	Numeric	Decimal	Integer	Difference
Apply mathematics deals with solution of _____	Integral	Numeric	Decimal	Integer	Integral
Apply mathematics deals with solution of _____	Functional	Numeric	Decimal	Integer	Functional
Apply mathematics deals with solution of _____	Algebraic	Numeric	Decimal	Integer	Algebraic
In graphical model the problem of 7 bridges is called _____	Fick's	Routhwelt	Konigsberg	Gauss	Konigsberg
A graph is called _____ if every pair of vertices is joined by an edge	Complete	Incomplete	Digraph	Continuous	Complete
A graph is called _____ if every edge is directed with an arrow	Complete	Incomplete	Digraph	Continuous	Digraph
A graph is called _____ if every edge has either + or - sign associated with it	Complete	Incomplete	Digraph	signed graph	signed graph
A digraph is called _____ if every directed edge has a weight associated wit it	Weighted digraph	signed graph	Digraph	Complete	Weighted digraph
A graph is called _____ if each of its vertices has same degree r.	regular	irregular	solid	unsolid	regular
If traffic is allowed from point a to b the edge can draw _____ from a to b.	undirected	directed	complete	incomplete	directed
If G is undirected connected graph then one can always direct _____ edge of G	Non circuit	Vertex	Non vertex	Circuit	Circuit
In genetic graph, the local degree of incoming edges at eac vertex must be less then or equal to _____	1	2	3	4	2
The necessary condition for a directed graph is to be _____	one way traffic	Two way traffic	Genetic	Nature	Genetic
The measure m(x) is always a _____ number	Real	complex	whole	natural	Whole
If x has no subordinates then measure m(x) equals _____	1	0	2	3	0
If witout oterwise changing the struture we move subordinate of a to a lower level relative to x then _____	Increases	decreases	stable	unstable	increases
If witout oterwise changing the struture we add a new individual subordinate to x then m(x) _____	Increases	decreases	stable	unstable	increases
In communication network a _____ graph can serve as a model	undirected	stable	directed	unstable	directed
An individual can send message direct to n individuals with propability _____	n	1/n	2n	3n	1/n
In a matrix representation an individual can send message to himself then _____ elements are	row	column	digonal	all	digonal
All the elements of matrix are non negative and the sum of elements of every row is unity, the matrix is _____	stochastic	propabilistic	direct	ergodic	stochastic
All the elements of matrix are non negative and the sum of elements of every row is _____, the matrix is _____	2	unity	3	4	unity
The markov chain is not _____	stochastic	propabilistic	direct	ergodic	ergodic
A subset is to form clique if every member of subset has a _____ relation with other member	symmetrical	non symmetrical	stable	unstable	symmetrical
A subset of persons in a socio - psychological group will set to form a _____	queue	clique	line	None	clique
The subset has atleast _____ member	1	2	3	4	3
If the group consists of n persons then can represents the group by _____ verties of graph	n+1	n-1	n	2n	n
For each communication netwrok can set up the corresponding _____ propability matrix	Digonal	Unit	row	Transition	Transition
A _____ graph Is one in which every edge has positive or negative sign	direct	undirect	signed	unsigned	signed
A _____ graph Is one in which every edge has positive or negative sign	direct	undirect	algebraic	unsigned	algebraic
The graph is balanced, every cycle in it is _____	Positive	negative	both	none	positive
All closed line sequences in the graph is _____	Positive	negative	both	none	positive
Any two lines sequence between two verties have the same _____	sign	number	constant	variable	sign
Any two lines sequence between two verties have the _____ sign	same	different	equal	none	same
The set of all points of graph can be partitioned into _____ disjoints sets	1	2	4	3	2
Every negative sign connects _____ points of differents set	1	2	3	4	2
Every positive sign connects _____ points of differents set	1	2	3	4	2
An algebraic grap is set to be antibalanced if every cycle in it has _____ no of positive edges	even	odd	real	distinct	even

Reg. No -----

(16MMP303)

**KARPAGAM ACADEMY OF HIGHER EDUCATION**

**Karpagam University**

**COIMBATORE-21**

**DEPARTMENT OF MATHEMATICS**

**Third Semester**

**I INTERNAL TEST- JUL '17**

**MATHEMATICAL MODELLING**

**Date: . 07.17 ( )**

**Time:2 hours**

**Class: IIM.Sc(Mathematics)**

**Maximum: 50 Marks**

---

**PART- A (20x1 = 20 Marks)**

**ANSWER THE FOLLOWING**

- 1.If there is one dependent continuous variable and a number of independent continuous variables then system is called  
a)ODE                      b)PDE c)LDE                      d)HDE
- 2.If there is immigration into the population from outside at a rate \_\_\_\_\_ to the population size  
a)Logically   b)exponentially   c)inversely   d) proportionally
- 3.If  $P(t)$  price of commodity and its rate of change is proportional to the \_\_\_\_\_ between demand and supply  
a)Addition   b)Difference   c)Division   d)Multiplication
- 4.Two chemical substances combined in the ratio \_\_\_\_\_ to form the third substances Z  
a) a:bb)a:2b                      c)2a:b                      d)a:3b
- 5.In population growth model b and d denotes  
a) Birth & death                      b) Business & death  
c)birth & decrease                      d)birth&increase

- 6.If there is immigration into the population from outside at a rate \_\_\_\_\_ to the population size  
a)Logically                      b) exponentially  
c)inversely                      d) proportionally
- 7.In radio geology the of age solar system is used to estimate \_\_\_\_\_  
a)Radio active   b)Diffusion   c)Decay                      d)immigration
- 8.In the model 'change of price of commodity  $S(t)$  denotes  
a)System   b)Supply                      c) Size                      d)model
- 9.\_\_\_\_\_ law is used in the model 'Diffusion'  
a)Fick's                      b) Hooke's                      c) Newton's                      d) Gauss
- 10.In the model 'change of price of commodity  $p_e$  denotes  
a)Equilibrium price                      b) Eligible price  
c) Essential price                      d) Evaluation price
- 11.If there are no prey the \_\_\_\_\_ species will decline at a rate proportional to the population.  
a)Prey                      b) Predator                      c) permanent                      d) persuieng
- 12.The population of  $x=0$  and  $y=0$  is called \_\_\_\_\_ position.  
a)zero                      b) equilibrium                      c) unit                      d)value
- 13.The real parts of all the eigenvalues of the matrix  $[c_{ij}]$  is negative are called  
a)Rout-Herwitz                      b) Fick's                      c) Newtons                      d) Gauss
- 14.In simple epidemic mode limit  $t$  tends to infinity of  $I(t)$  denotes \_\_\_\_\_  
a)n                      b)  $n+1$                       c)  $n-1$                       d)  $2n$
- 15.The predator species increases and the prey species \_\_\_\_\_ at a rate proportional to the product of two populations.

a)increases b)decreases c)uniformly d)stable

16.The initial populations of prey and predator species are

a)p/q and a/b b)a/b and p/q c)a/b d) p/q

17. $x_1(t), x_2(t), \dots, x_n(t)$  represent the populations of n species states\_\_\_\_\_ model.

a)multi-species b)single-species c)prey d)predator

18.A susceptible person can infected at a rate proportional to

a)SI b)SIS c) SHM d) MOC

19.The population of  $x=0$  and  $y=0$  is called \_\_\_\_\_ position.

a)zero b) equilibrium c)unit d)null

20.The rate of growth of each species \_\_\_\_\_ due to the presence of the other.

a)increases b)decreases c)uniformly d)stable

#### **PART- B (3 x 2 = 6 Marks)**

#### **ANSWER ALL THE QUESTIONS**

21.Explain diffusion in linear growth and decay.

22.In equation of logistic law,  $k=0.007, R=1000, N(0)=50$ , find  $N(10)$ .

23.Find curves for which tangent at a point is always perpendicular to the line joining the point to the origin.

#### **PART- C (3 x 8 = 24 Marks)**

#### **ANSWER ALL THE QUESTIONS**

24.a ) Discuss about logistic law of population growth.

(OR)

b) Explain about population growth models

25. a) Suppose the population of the world now is 4 billion and its doubling period is 35 years, what will be the population of the world after 350 years?

(OR)

b) Explain about prey-predator models.

26. a)Derive a Simple Epidemic Model.

(OR)

b)Discuss in detail Domar Macro models.



Reg. No -----

(16MMP303)

**KARPAGAM UNIVERSITY**  
**KARPAGAM ACADEMY OF HIGHER EDUCATION**  
**COIMBATORE-21**

**DEPARTMENT OF MATHEMATICS**

**Third Semester**

**II INTERNAL TEST- SEP '17**

**MATHEMATICAL MODELLING**

**Date: . 09.17 ( )**

**Time: 2 hours**

**Class: II M.Sc(Mathematics)**

**Maximum: 50 Marks**

---

**PART- A (20x1 = 20 Marks)**

**ANSWER THE FOLLOWING**

1. In a model Motion under inverse square law, the conic is parabola then  
a)  $e < 1$       b)  $e = 1$       c)  $e > 1$       d)  $e = 0$
2. In \_\_\_\_\_ the genetic characteristics will change generation to generation and the variable representing generation is discrete variable.  
a) Economics      b) Genetics  
c) Population dynamics      d) Mechanics
3. \_\_\_\_\_ no of kepler's law are in planetary motion  
a) 1      b) 2      c) 3      d) 4
4. Complementary function can be obtained by \_\_\_\_\_  
a) Matrix      b) Determinate  
c) Eigen value      d) Eigen Vector
5. The solution of linear differences equation can be obtained by \_\_\_\_\_ transform if  $t$  is continuous

- a) Laplace    b) Z    c) Fourier    d) Gauss
6. A graph is called \_\_\_\_\_ if every pair of vertices is joined by an edge  
a) Complete      b) Incomplete  
c) Digraph      d) Continuous
  7. In graphical model the problem of 7 bridges is called \_\_\_\_\_  
a) Fick's      b) Routhwelt  
c) Konigsberg      d) Gauss
  8. The markov chain is not \_\_\_\_\_  
a) Stochastic      b) propabilistic  
c) direct      d) ergodic
  9. Any two lines sequence between two verties have the \_\_\_\_\_ sign  
a) Same    b) different    c) equal    d) one
  10. In \_\_\_\_\_ the price changes are consider from year to year or month to month or week to week or day to day  
a) Economics      b) Genetics  
c) Population dynamics      d) Population growth
  11. The solution of linear differential equation is of the form  
a)  $CF + PI$     b)  $CF - PI$     c)  $CF * PI$     d)  $CF / PI$
  12. In difference equation \_\_\_\_\_ theory is applied  
a) Stability      b) Non stability  
c) Uniformity      d) Non uniformity
  13. In the cobweb model price of the commodity in the year denotes  
a)  $pt$       b)  $qt$       c)  $rt$       d)  $st$
  14. If  $G$  is undirected connected graph then one can always direct \_\_\_\_\_ edge of  $G$   
a) Non circuit      b) Vertex  
c) Non vertex      d) Circuit
  15. The measure  $m(x)$  is always a \_\_\_\_\_ number  
a) Real      b) complex    c) whole      d) natural

16. In communication network a \_\_\_\_\_ graph can serve as a model  
 a) undirected    b) stable    c) directed    d) unstable
17. In a matrix representation an individual can send message to himself then \_\_\_\_\_ elements are zero.  
 a) row    b) column    c) diagonal    d) all
18. If the group consists of  $n$  persons then can represents the group by \_\_\_\_\_ vertices of graph  
 a)  $n$     b)  $n+1$     c)  $n-1$     d)  $2n$
19. The necessary condition for a directed graph is to be \_\_\_\_\_  
 a) one way traffic    b) Two way traffic  
 c) Genetic    d) Nature
20. If without otherwise changing the structure we add a new individual subordinate to  $x$  then  $m(x)$  \_\_\_\_\_  
 a) Increases    b) decreases    c) stable    d) unstable

**PART- B (3 x 2 = 6 Marks)**

**ANSWER ALL THE QUESTIONS**

21. Write the components of velocity and acceleration vectors along radial and transverse directions.
22. Explain formula for Laplace Transform.
23. Explain Food Webs.

**PART- C (3 x 8 = 24 Marks)**

**ANSWER ALL THE QUESTIONS**

24. a) Explain about the catenary.  
 (OR)  
 b) Explain in detail Kepler's law of planetary motion
25. a) Find a solution of linear difference equation by Laplace

transform.

(OR)

- b) Explain in detail Harrod model.
26. a) Discuss in detail weighted digraphs and markov chains.  
 (OR)  
 b) Discuss in detail about communication networks.

[15MMP306]

**KARPAGAM UNIVERSITY**  
Karpagam Academy of Higher Education  
(Established Under Section 3 of UGC Act 1956)  
COIMBATORE – 641 021  
(For the candidates admitted from 2015 onwards)

**M.Sc., DEGREE EXAMINATION, NOVEMBER 2016**  
Third Semester

**MATHEMATICS**

**MATHEMATICAL MODELING**

Time: 3 hours

Maximum : 60 marks

**PART – A (20 x 1 = 20 Marks) (30 Minutes)**  
**(Question Nos. 1 to 20 Online Examinations)**

**(Part - B & C 2 ½ Hours)**

**PART B (5 x 6 = 30 Marks)**  
**Answer ALL the Questions**

21. a) Discuss about logistic law of population growth.  
Or  
b) Give a brief note on diffusion or a medicine in the blood stream.
22. a) Derive a Simple Epidemic Model.  
Or  
b) Show that national income, investment and savings increase exponentially.
23. a) Explain about the catenary.  
Or  
b) Explain in detail Kepler's law of planetary motion.
24. a) Write an explanatory note on complementary function.  
Or  
b) Explain in detail markov chains.
25. a) Discuss in detail weighted digraphs and markov chains.  
Or  
b) Discuss in detail about communication networks.

**PART C (1 x 10 = 10 Marks)**  
**(Compulsory)**

26. Explain about motion of a projectile.